



University of
Salford
MANCHESTER

Using Compassion Focused Therapy as an adjunct to Trauma-Focused CBT for fire service personnel suffering with symptoms of trauma

Beaumont, EA, Durkin, M, McAndrew, SL and Martin, C

<http://dx.doi.org/10.1017/S1754470X16000209>

Title	Using Compassion Focused Therapy as an adjunct to Trauma-Focused CBT for fire service personnel suffering with symptoms of trauma
Authors	Beaumont, EA, Durkin, M, McAndrew, SL and Martin, C
Type	Article
URL	This version is available at: http://usir.salford.ac.uk/40757/
Published Date	2016

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: usir@salford.ac.uk.

Table 1: Some of the treatment interventions incorporated into both conditions

Trauma-focused CBT (Ehlers et al., 2005)	Compassion-focused therapy (Gilbert, 2010)
<p>Identifying relevant appraisals, memory characteristics and triggers</p> <p>Identifying behavioural and cognitive strategies that maintain PTSD</p> <p>Examining “hot spots”</p> <p>Socratic questioning</p> <p>Identifying an alternative new appraisal – e.g., by adding it to a written account or by using imaginal reliving</p> <p>Revisiting the scene of the trauma to: - (1) obtain evidence that helps explain why or how an event occurred. This is helpful for FSP who have appraisals such as “I could have prevented this from happening” and (2) focusing on what was different between “then” and “now”</p> <p>Reclaiming work – reintroducing social and behavioural activities that have been avoided or given up following the trauma</p> <p>Develop a narrative account - starting before the trauma and ending after the individual is safe again. Events are placed in the past</p> <p>Cognitive restructuring - focusing on the personal meanings of the trauma and its sequelae</p> <p>Examination of maintaining strategies - rumination, hypervigilance and/or safety behaviours</p>	<p>Developing sympathy, acceptance and insight into one’s own difficulties through self-reflection and mindfulness</p> <p>Learning to notice and experience physiological and psychological reactions with compassion, empathy and kindness</p> <p>Developing breathing techniques – e.g., Soothing Rhythm Breathing</p> <p>Creating an imaginary safe place in the mind’s eye that provides a sense of calm and peace</p> <p>Imagining and using acting skills to experience a compassionate self</p> <p>Experiencing compassion as a flow which can flow in three ways: - (1) from other people to oneself, (2) from oneself to other people and (3) from and to self</p> <p>Using thought records to explore the role played by self-critical rumination</p> <p>Learning to respond compassionately to the ‘bully within’</p> <p>Thinking about and responding to the anxious, sad, angry and critical self</p> <p>Compassionate letter writing which focuses on being kind, supportive and nurturing as opposed to being self-critical.</p> <p>Creating a ‘step by step’ approach to cope with trauma symptoms such as avoidance</p>

Table 2: Pre and Post-therapy Mean Scores and Standard Deviations for the CBT only and the combined group

	Therapy Type											
	CBT Group (<i>n</i> = 8)						CBT + CFT Group (<i>n</i> = 9)					
	Pre		Post		Diff		Pre		Post		Diff	
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>
HADS												
Anxiety	10.3	2.7	4.4	1.9	6.0	2.6	14.8	4.5	5.3	1.1	9.5	4.7
Depression	10.6	3.5	4.9	2.0	5.7	3.0	15.9	3.3	5.9	1.4	10.0	2.6
IES												
Avoidance	19.7	6.2	7.7	3.4	12.0	4.4	22.1	5.0	5.0	3.5	17.1	6.3
Hyper-arousal	15.4	5.1	5.1	3.3	10.3	2.9	13.2	5.6	3.4	2.9	9.8	4.4
Intrusion	20.9	5.0	7.4	3.9	13.5	5.2	23.0	5.9	6.9	4.6	16.1	5.9
Total IES	56.0	7.5	20.2	6.4	35.8	7.7	54.0	15.3	15.3	9.1	38.7	13.0
SCS												
	1.9	0.5	3.1	0.4	1.3	0.7	2.2	0.8	3.9	0.6	1.7	0.9

HADS = Hospital Anxiety and Depression Scale; IES = Impact of Events Scale; SCS = Self-Compassion Scale