



University of
Salford
MANCHESTER

Attending a workplace : its contribution to volume and intensity of physical activity

Rafferty, D, Dolan, C and Granat, MH

<http://dx.doi.org/10.1088/0967-3334/37/12/2144>

Title	Attending a workplace : its contribution to volume and intensity of physical activity
Authors	Rafferty, D, Dolan, C and Granat, MH
Type	Article
URL	This version is available at: http://usir.salford.ac.uk/40894/
Published Date	2016

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: usir@salford.ac.uk.

Title:

Attending a workplace: its contribution to volume and intensity of physical activity.

Authors:

Mr Daniel Rafferty¹, Ms Catriona Dolan¹, Professor Malcolm Granat².

Affiliations:

¹ Glasgow Caledonian University, Institute for Applied Health Research, Cowcaddens Road, Glasgow, G4 OBA, UK.

² University of Salford, Centres for Health Sciences Research, The University of Salford, The Crescent, Salford, M5 4WT, UK.

Corresponding Author:

Mr Daniel Rafferty. Glasgow Caledonian University, Institute for Applied Health Research, Cowcaddens Road, Glasgow, G4 OBA, UK.

Email: d.raffety@gcu.ac.uk

Telephone: ++ 141 331 8138

Fax: ++ 141 331 8110

Abstract

Purpose

Understanding the contribution that attending a workplace has in accumulating physical activity (PA) may help inform strategies used to increase PA. This study explores the influence that attending work has on the total number of steps taken and the time spent in moderate to vigorous activity (MVPA).

Methods

A global position system (GPS) was used to identify the geographical domain of the participant. An activity monitor (activPAL, PALtechnologies Ltd, Glasgow, UK) was employed to measure the number of steps taken and the cadence of those steps. Both devices were worn for seven consecutive days and 5 work days extracted post data collection. The data from the two devices were synchronised allowing domain, volume and intensity of PA to be explored. The distance from the home domain to the workplace was used to establish if there was any relationship between commute distance and number of steps accumulated and time in MVPA.

Results

Twenty-six office workers [17F; mean age 38 (range 23–65)] were recruited. The number of steps taken per day on average for the group was 11,008 (SD \pm 2,999) with time spent in MVPA per day being 32.7 (SD \pm 17.1) minutes. The commute accounted for 32% or 3,550 (SD \pm 1664) of the steps taken and 68% or 22.0 (SD \pm 14.1) minutes of MVPA. No statistically significant correlations with distance from home to the workplace for either variable were found.

Conclusion

This work explores the contribution that attending work makes to PA, combining data from a GPS system and an objective activity monitor. The commute to works accounts for more than two-thirds of the MVPA accumulated per day. This provides meaningful insight into the volume and intensity of individuals' activity and also its context.

Key words:

Objective activity monitoring, GPS, active travel, commute to work, Occupational activity.

Introduction

The health benefits of physical activity (PA) have been well documented (38). Current guidelines recommend that adults accumulate at least 30 minutes a day of moderate to vigorous activity (MVPA) where MVPA is defined as movement characterised as having a metabolic equivalent (METs) of between 3 and 6 (39). Individuals may accumulate their MVPA in a variety of contexts and physical locations which are usually related to their occupation, daily routines and preferred leisure pursuits (29). Understanding where and at what volume and intensity PA occurs may help to inform public health messages and strategies to increase PA. Focusing on increasing leisure-time PA alone may result in targeting groups less in need of intervention, whilst missing opportunities for increasing PA in the wider population (17). Physical activity leads to increased energy expenditure, with total PA considered the only modifiable variable in relation to total energy expenditure. This can be dichotomised between work and non-work related PA for a working population (6). However, whilst commonly used subjective measures of PA, such as the International Physical Activity Questionnaire (IPAQ) (8) attempt to categorise activity into domains of occupational, transportation, household and recreational domains, little has been done to objectively categorise PA in this way.

For working adults with increasingly sedentary occupations (18,23), technology that facilitates less movement (23,28), and longer working hours (12) present challenges for increasing daily PA. A number of workplace interventions have been successful at increasing PA in office workers', in the short-term. For example, encouraging employees to engage in active breaks which involve performing moderate intensity exercises with co-workers away from their desks (2,33). Promoting active travel is another way of attempting to increase PA

in the working population (3,37). Strategies for promoting active travel include; cycle to work schemes that reduce the cost of bicycles and related accessories, encouraging employees to get off public transport a stop earlier and walk the extra distance, work-based competitions surrounding use of active travel modes, as well as policy changes such as reduced parking and improved infrastructure (9). The Osaka health study found that, regardless of the chosen mode of transportation, increasing the amount of walking to/from work had an independent effect on reducing the risk of hypertension (14).

In addition to lack of PA, workers also face the health risks associated with too much sedentary behaviour (SB). This carries an increased risk of ill health, including cancer, cardiovascular disease, diabetes, and obesity, independent to the amount of physical activity an individual may perform (22,34). Research into effective interventions to reduce occupational SB is still in its infancy(32), and, to add complexity, a recent meta-analysis concluded that interventions that contained both a component to increase PA and to reduce SB were less effective (24).

Commuting distances to our workplace have increased (20) as residential areas become further away from the workplace. Having a longer commute could result in greater opportunities for PA, however it could be that longer commutes increase the likelihood of non-active transportation being chosen.

Both workplace interventions and active travel employ different strategies to increase PA, resulting in the literature being multi-faceted. However, more needs to be understood about the contribution of workplace activity (occurring at the place of work) and commuting

activity (travelling to/from place of work) to total MVPA. In particular, the volume and intensity of PA should be put in the context of where and why the PA is being undertaken. Currently, much of our understanding about workplace and commuting activity is based on self-report or travel survey information(10,35). The use of objective activity monitors allows more accurate measurement of the volume and intensity of activity. Combining these data with information from a Global Positioning System (GPS) allows it be categorised in terms of physical domain and context. Combined GPS and activity monitoring data has been previously used to explore changes in transportations modes (1,3,19,27). However, differences in MVPA categorised into home or commute or work place has yet to be explored in this way.

This aim of this study was to use activity monitors to objectively measure the volume and intensity of PA in a group of office workers and combine this data with GPS data to categorise MVPA in terms of the domain and context of activity. This would allow an understanding of the contribution of workplace and commuting MVPA to total daily MVPA facilitating a greater understanding of the activity profile of those employed in sedentary occupations. The study also explored the relationship between commuting distance, steps accumulated (volume), and time in MVPA (intensity). Such insights will be valuable in determining the focus of future PA interventions targeted at working adults.

Methods

Study design and participants

This was a descriptive observational study with participants recruited by email using a convenience sample of employees at Glasgow Caledonian University. Participants had to be aged 18 years or older, be in full-time employment (35hrs/week) with normal working days of Monday through to Friday, and have a job that they generally considered being seated at a desk (e.g. academic, administrative, technical, full-time PhD student). Participants were excluded from taking part if they had impaired mobility due to medical reasons or if an unaccustomed change in activity was anticipated during the data collection period (e.g. vacations). A suitable data collection period was arranged with each participant to ensure a target of 7 consecutive days of data were collected. Subject's demographics were recorded. All subjects gave informed, written consent and the study was approved by Glasgow Caledonian University ethics committee. Work days were identified and extracted from the data.

To help understand the PA levels of those taking part, participants were categorised as being sedentary or inactive (< 3,000 steps per day), low active (3,000-7,499), somewhat active (7,500-9,999), active (10,000-12,499), or highly active (>12,500) (16).

MVPA

MVPA was defined as a period of walking with a cadence of greater than 109 steps/minute (4). This value was chosen as it has been previously demonstrated in a small treadmill study that healthy young males walking with a MET value of 5 adopt a cadence of 123.6 ± 4.9 steps per minute (36). Therefore in order to include all stepping which might be within this

moderate intensity range a value of the mean cadence less three standard deviations (109 steps/min) would include 99% of the population assuming a normal distribution.

PA Measurement

Activity data was collected using an activPAL, a thigh-worn accelerometer-based activity monitor (PALtechnologies Ltd, Glasgow, UK). The activPAL has been demonstrated to be accurate for measuring step count and cadence (13). The device was sealed in a neoprene sleeve, attached to the participant's thigh using activPAL stickies and then covered with Tegaderm tape. The device was worn continuously except when bathing or showering. The data was downloaded from the device and exported to Excel in 15s epochs (PalExcel files). Time spent walking in each epoch and the number of steps taken in that epoch is reported as output variables. Cadence was calculated as the number of steps taken divided by time walking in seconds in the 15s epoch then multiplying by 60. Time in MVPA was calculated by summing the time where the cadence was above 109 steps/min. Time in MVPA and number of steps was summed for each domain, determined by the GPS data, on each day and average values per day calculated and averaged for all participants. Additionally, the total number of steps and total time in MVPA per day was also calculated.

Domain Measurement

The domain data was collected using the Amod AGL3080 GPS data-logger device (AMOD Technology, Taipei City, Taiwan). The GPS device was of a suitable size to be placed in a bag or pocket and was carried every time participants went outdoors. The latitude and longitude of the device's domain, when outdoors, was provided on a second-by-second basis using GPS Utility software (<http://www.gpsu.co.uk>). The GPS Utility

(<http://www.gpsu.co.uk/>) was then used to convert this to text files for use in Excel (GPSExcel files). To remove noise and spurious data the GPS domain data was averaged over a 15s period. When domain information was unavailable the device was considered to be at the last domain identified using Google Maps until domain data from the device became visible again.

Combining PA and domain measurement

The domain of where the person was could be determined by the GPS data and Google maps. Six possible domains were identified (Table 1):

Domain	Definition
Work	in the work place
Commute	leaving home to arriving at the work place or vice versa
Work excursion	at work but being outside going to or coming back from another domain e.g. going out for lunch
Home	in the home
Home excursion	at home but being outside, going to or coming back from another domain e.g. in the garden, going to or coming from a restaurant/cinema
Other	in another domain e.g. shopping malls, restaurants, cinema

Table 1 the definition to define each domain

The time stamped PAExcel and the GPSExcel files were then synchronised in Excel using the 15s data generated from both the activity monitor and the GPS device allowing the

domain where the activity took place to be identified. These were then expressed as a percentage of the total number of steps taken and total time in MVPA.

Statistical analysis

As this was an observational study no inferential statistics were applied. All data presented are averages and standard deviations for the collection period.

Spearman's rho correlation was conducted on the distance from home to the workplace, the total number of steps taken and the time in MVPA. Depending on the distribution of the data either a Spearman's rho or a Pearson correlation was conducted for the time in MVPA and the number of steps taken in each domain.

Results

Twenty-six office workers [17F; mean age 38 (range23–65)] were recruited. On average participants lived 19.4 km (SD \pm 19.5, max 64.1, min 0.8) from their place of work.

Physical activity levels of participants are shown in Table 2.

Activity level	Number of participants
(Steps per day)	
Inactive	1
(<5000)	
Low active	3
(5000-7499)	
Somewhat active	5
(7500-9999)	
Active	11
(10000-12500)	
Highly active	6
(>12500)	

Table 2: Participants physical activity levels

Eighteen of the 26 participants (69%) achieved more than an average of 30 minutes of MVPA per day, with 5 of those achieving more than 30 minutes of MVPA (19%) in their commute alone. In no other domains did any participant achieve more than 30 minutes of MVPA.

The total number of steps and time in MVPA and the correlations between time in MVPA and the steps taken in each domain are presented in Table 3.

	Steps per day	Time in MVPA (mins)	r^2	p
	Average(\pm SD)	Average(\pm SD)		
Total ^a	11008 (2999)	32.7 (17.1)	0.636	<0.001
Commute ^a	3550 (1664)	22.0 (14.1)	0.636	<0.001
Work excursion	890 (811)	4.1 (5.0)	0.537	0.005
Work ^a	3564 (1599)	3.5 (3.8)	0.334	0.096
Home	1610 (1032)	0.3 (0.7)	-0.162	0.428

Home	425	1.3	0.184	0.396
excursion	(1053)	(2.9)		
Other	970	1.5	0.143	0.485
	(808)	(2.1)		

Table 3: Group averages (\pm SDs) for total steps and time in MVPA for each domain and correlations of total number of steps taken with time in MVPA in each domain. ^a denotes a Pearson's product correlation, all other correlations were a Spearman's rho.

The percentage of total steps taken and the time in MVPA are displayed in Figure 1(a&b)




Figure 1(a)




Figure 1(b)

Figure 1(a&b) Percentages of steps taken and time in MVPA in each domain.

No significant correlations were found when the distance from the workplace data was explored for either time in MVPA or total number of steps taken ($p>0.05$) nor steps taken during the commute or time in MVPA ($p>0.05$).

Discussion

The results suggest that commuting, walking in the work place, and making excursions whilst at work leads to the accumulation of the majority of both an individual's daily step count and MVPA, independent of commuting distance. In this study, the commute, work excursion and steps in the workplace accounted for 8,004 (72%) of the average 11,008 steps per day, with 29.6 min (91%) of time in MVPA being accrued when either commuting to work, at work, or during excursions made during work time. This is notable in a population whose occupation is predominantly desk-based and, therefore considered sedentary and at most risk of the negative health consequences of inactivity (5,15).

The use of GPS monitoring alongside accelerometer data is increasing in PA research, as the importance of the environment in which PA occurs is being realised (21,30). Recent studies using such methods have focused on the impact of the built environment on PA (26,31) and the home-school commute of school children (7). A study by Oliver, et al. (21) noted the potential for using a combination of GPS and accelerometer data to understand travel related PA and the potential for using a combination of GPS and accelerometer data to understand changes in domain and PA. As far as the authors are aware, this is the first study to use these methods to try and understand the impact of the work-home-commute on overall PA.

Sixty-eight percent of work day MVPA was accumulated during the commute and it was the only domain in which more than 30 minutes of MVPA was recorded (n=5). Both MVPA and step count during the commute was independent of the distance commuted ($p>0.05$). These results, therefore, suggest that commuting distance is not a barrier to accumulating MVPA through travel to and from the workplace. Workplace schemes aimed at encouraging

employees to commute using active travel modes e.g. cycle to work schemes, walk-to-work week, should therefore not neglect those whose distance or circumstances does not facilitate active travel for the entire journey. There could be merit in providing information to employees on how to increase their MVPA during commuting such as by walking briskly to/from public transport stations or considering increasing the distance between motorised transport and work (e.g. by getting off the bus one stop early, parking further away).

Whilst commuting accounted for the highest proportion of work day MVPA, time spent at work, including leaving the workplace during work time, together represented 23% of daily MVPA. This demonstrates that it is possible to incorporate MPVA even whilst working in a predominantly sedentary occupation. Therefore, ways to encourage employees to further increase their activity during work hours could be worthy of further consideration. Workplace interventions aimed at increasing PA through increasing step counts (11,25) and encouraging active breaks (2,33) have proved successful at increasing workplace PA. However, the impact of such interventions on PA accumulated during the commute or leisure time is not known. Interventions to increase MVPA during work hours, should therefore also aim to ensure that PA during commuting and leisure time is maintained.

Work excursion marginally represented the largest in-work MVPA at 12.4% and was significantly correlated with the overall time in MVPA. There are therefore health benefits to encouraging employees to leave the workplace during their lunch break rather than eat lunch at their desks. This could be as simple as publicising short walking routes in the neighbourhood or information on discounts for nearby facilities. Further research is needed to investigate the impact of such promotions on the PA patterns of employees.

There are a number of limitations to this study. The sample size is relatively small and is drawn from a University population. The University estate consists of 105,000m² of buildings and 86,000m² of grounds making the campus relatively large and could, therefore, have led to an overestimation in MVPA actual undertaken in the workplace in comparison to employers with smaller sites. However, this does not affect the contribution of commutes to the proportion of total MVPA. Self reported measures may have provided additional information. However there was considerable discrepancies between the technology and self reported. For the purposes of this work, the technology measures were considered more robust.

Conclusion

Commuting to work can lead to the accumulation of the majority of an individual's MVPA, independent of commuting distance. This study highlights the benefits of investigating linked GPS and accelerometer data to increase understanding of how PA is accumulated during daily living. Seventy-two percent of daily step count was accumulated either commuting to (32%), in work (32%) or during workplace excursions(8%). This is encouraging as trends suggest that commuting distances are increasing (20). Strategies should be developed to ensure that active travel remains a component of work commuting trips by promoting its benefits to workers and providing them with information on how to increase this element of their trip.

Acknowledgments

The authors would like to acknowledge the participants in this work and their diligence in complying with the data collection methods. Ms Gillian McLellan for the effort made to recruit, conduct the data collection and preparation of the data for analysis. There are no funding sources to report.

Conflict of interest

Professor Malcolm Granat is a co-inventor of the activPAL physical activity monitor and is currently a director of PAL Technologies Ltd.

References

- (1) Almanza E, Jerrett M, Dunton G, Seto E, Pentz MA. A study of community design, greenness, and physical activity in children using satellite, GPS and accelerometer data. *Health Place* 2012;18(1):46-54.
- (2) Bort-Roig J, Martin M, Puig-Ribera A, et al. Uptake and factors that influence the use of 'sit less, move more' occupational intervention strategies in Spanish office employees. *International Journal of Behavioral Nutrition and Physical Activity* 2014;11(1):152-160
- (3) Chaix B, Kestens Y, Duncan S, et al. Active transportation and public transportation use to achieve physical activity recommendations? A combined GPS, accelerometer, and mobility survey study. *The international journal of behavioral nutrition and physical activity* 2014;11:124.
- (4) Chastin SFM, Dall PM, Tigbe WW, et al. Compliance with physical activity guidelines in a group of UK based postal workers using an objective monitoring technique. *European Journal of Applied Physiology* 2009;106(6):893--899.
- (5) Chau JY, Grunseit AC, Chey T, et al. Daily sitting time and all-cause mortality: a meta-analysis. *PloS one* 2013;8(11):e80000.
- (6) Church TS, Thomas DM, Tudor-Locke C, et al. Trends over 5 decades in U.S. occupation-related physical activity and their associations with obesity. *PloS one* 2011;6(5):e19657.
- (7) Collins P, Al-Nakeeb Y, Lyons M. Tracking the commute home from school utilizing GPS and heart rate monitoring: establishing the contribution to free-living physical activity. *Journal of physical activity & health* 2015;12(2):155.

- (8) Craig CL, Marshall AL, Sjöström M, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 2003;35(8):1381-1395.
- (9) de Nazelle A, Nieuwenhuijsen MJ, Antó JM, et al. Improving health through policies that promote active travel: A review of evidence to support integrated health impact assessment. *Environ Int* 2011;37(4):766-777.
- (10) Department for Transport. National Travel Survey England 2015. 2016:1--54. Available from:
https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/551437/national-travel-survey-2015.pdf [accessed 29/09/2016]
- (11) Gilson ND, Puig-Ribera A, McKenna J, Brown WJ, Burton NW, Cooke CB. Do walking strategies to increase physical activity reduce reported sitting in workplaces: a randomized control trial. *The international journal of behavioral nutrition and physical activity* 2009;6(1):43-43.
- (12) Goetzel R, Ozmlnkowski R. The health and cost benefits of work site health-promotion programs. *Annu Rev Public Health* 2008;29:303-323.
- (13) Grant PM, Ryan CG, Tigbe WW, Granat MH. The validation of a novel activity monitor in the measurement of posture and motion during everyday activities. *Br J Sports Med* 2006;40(12):992-997.
- (14) Hayashi T, Tsumura K, Suematsu C, Okada K, Fujii S, Endo G. Walking to Work and the Risk for Hypertension in Men: The Osaka Health Survey. *Ann Intern Med* 1999;131(1):21.

- (15) Healy GN, Lawler S, Thorp A, et al. Reducing prolonged sitting in the workplace. An evidence review: full report. 2012;ISBN:978-1-921822-06-3.
- (16) Hills AP, Mokhtar N, Byrne NM. Assessment of physical activity and energy expenditure: an overview of objective measures. *Frontiers in nutrition* 2014;1:5.
- (17) Khaing Nang EE, Khoo EYH, Salim A, Tai ES, Lee J, Van Dam RM. Patterns of physical activity in different domains and implications for intervention in a multi-ethnic Asian population: a cross-sectional study. *BMC Public Health* 2010;10(1):644-644.
- (18) King GA, Fitzhugh EC, Bassett Jr DR, McLaughlin JE, Strath SJ, Swartz AM, et al. Relationship of leisure-time physical activity and occupational activity to the prevalence of obesity. *Int J Obes* 2001;25(5):606-612.
- (19) Lachowycz K, Jones AP, Page AS, Wheeler BW, Cooper AR. What can global positioning systems tell us about the contribution of different types of urban greenspace to children's physical activity? *Health Place* 2012;18(3):586.
- (20) Office for National Statistics. 2011 Census Analysis - Distance Travelled to Work. 2014. Available from: <http://www.ons.gov.uk/ons/rel/census/2011-census-analysis/distance-travelled-to-work/index.html> [accessed 01 February 2016]
- (21) Oliver M, Badland H, Mavoa S, Duncan MJ, Duncan S. Combining GPS, GIS, and accelerometry: methodological issues in the assessment of location and intensity of travel behaviors. *Journal of physical activity & health* 2010;7(1):102.
- (22) Owen N, Healy GN, Matthews CE, Dunstan DW. Too Much Sitting: The Population-Health Science of Sedentary Behavior. *Exerc Sport Sci Rev* 2010 07;38(3):105-113.

(23) Parry S, Straker L. The contribution of office work to sedentary behaviour associated risk. *BMC Public Health* 2013 03/26;13:296-296.

(24) Prince SA, Saunders TJ, Gresty K, Reid RD. A comparison of the effectiveness of physical activity and sedentary behaviour interventions in reducing sedentary time in adults: a systematic review and meta-analysis of controlled trials. *Obesity reviews : an official journal of the International Association for the Study of Obesity* 2014;15(11):905-919.

(25) Rashid M, Kampschroer K, Wineman J, Zimring C. Spatial layout and face-to-face interaction in offices a study of the mechanisms of spatial effects on face-to-face interaction. *Environment and Planning B: Planning and Design* 2006;33:825--844.

(26) Rodriguez DA, Brown AL, Troped PJ. Portable Global Positioning Units to Complement Accelerometry-Based Physical Activity Monitors. *Medicine and Science in Sports and Exercise* [H.W.Wilson - EDUC] 2005;37(11):S572.

(27) Rodriguez DA, Cho G, Elder JP, et al. Identifying walking trips from GPS and accelerometer data in adolescent females. *Journal of physical activity & health* 2012;9(3):421.

(28) Saidj M, Oppert J, Menai M, et al. Descriptive study of sedentary behaviours in 35,444 French working adults: cross-sectional findings from the ACTI-Cités study. *BMC Public Health* 2015;15(1):1-10

(29) Samitz G, Egger M, Zwahlen M. Domains of physical activity and all-cause mortality: systematic review and dose-response meta-analysis of cohort studies. *Int J Epidemiol* 2011;40(5):1382-1400.

- (30) Schoeppe S, Duncan MJ, Badland H, Oliver M, Curtis C. Associations of children's independent mobility and active travel with physical activity, sedentary behaviour and weight status: A systematic review. *Journal of Science and Medicine in Sport* 2013 7;16(4):312-319.
- (31) Seeger CJ, Welk GJ, Erickson S. Using Global Position Systems and physical activity monitors to assess the built environment. *URISA Journal* 2008;20(2):5.
- (32) Shrestha N, Ijaz S, Kukkonen-Harjula KT, Kumar S, Nwankwo CP. Workplace interventions for reducing sitting at work. *The Cochrane database of systematic reviews* 2015;1:CD010912.
- (33) Taylor WC, King KE, Shegog et al. Booster Breaks in the workplace: participants' perspectives on health-promoting work breaks. *Health Educ Res* 2013 JUN 2013;28(3):414-425.
- (34) Thorp AA, Owen N, Neuhaus M, Dunstan DW. Sedentary Behaviors and Subsequent Health Outcomes in Adults. *Am J Prev Med* 2011;41(2):207-215.
- (35) Transport Scotland. Transport and Travel in Scotland 2014. 2015:1--60. Available from: http://www.transport.gov.scot/sites/default/files/documents/rrd_reports/uploaded_reports/j389989/j389989.pdf [accessed 29/09/2016]
- (36) Tudor-Locke C, Sisson SB, Collova T, Lee SM, Swan PD. Pedometer-determined step count guidelines for classifying walking intensity in a young ostensibly healthy population. *Canadian Journal of applied physiology* 2005;30(6):666--676.
- (37) Wanner M, Götschi T, Martin-Diener E, Kahlmeier S, Martin BW. Active transport, physical activity, and body weight in adults: a systematic review. *Am J Prev Med* 2012;42(5):493.

(38) Warburton DER, Nicol CW, Bredin SSD. Health benefits of physical activity: the evidence. CMAJ : Canadian Medical Association journal = journal de l'Association medicale canadienne 2006;174(6):801-809.

(39) World Health Organisation. Global Strategy on Diet, Physical Activity and Health. 2016; Available at: http://www.who.int/dietphysicalactivity/physical_activity_intensity/en/.

Captions

Figure 1(a&b) Percentages of steps taken and time in MVPA in each domain.