



University of
Salford
MANCHESTER

Exploring the evidence base for acupuncture in the treatment of Meniere's Syndrome — A systematic review

Long, AF, Xing, M, Morgan, K and Brettle, A

<http://dx.doi.org/10.1093/ecam/nep047>

Title	Exploring the evidence base for acupuncture in the treatment of Meniere's Syndrome — A systematic review
Authors	Long, AF, Xing, M, Morgan, K and Brettle, A
Type	Article
URL	This version is available at: http://usir.salford.ac.uk/id/eprint/12662/
Published Date	2011

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: usir@salford.ac.uk.

Original Article

Exploring the Evidence Base for Acupuncture in the Treatment of Ménière's Syndrome—A Systematic Review

Andrew F. Long¹, Mei Xing², Ken Morgan³ and Alison Brettle⁴

¹School of Healthcare, University of Leeds, Leeds, LS2 9UT, UK, ²School of Community Health Sciences and Social Care, University of Salford, Salford, UK, ³Acupuncturist, Bolton, UK and ⁴School of Nursing, University of Salford, Salford, UK

Ménière's syndrome is a long-term, progressive disease that damages the balance and hearing parts of the inner ear. To address the paucity of information on which evidence-based treatment decisions should be made, a systematic review of acupuncture for Ménière's syndrome was undertaken. The method used was a systematic review of English and Chinese literature, from six databases for randomized, non-randomized and observational studies. All studies were critically appraised and a narrative approach to data synthesis was adopted. Twenty-seven studies were included in this review (9 in English and 18 in Chinese languages): three randomized controlled trials, three non-randomized controlled studies and four pre-test, post-test designs. All but one of the studies was conducted in China. The studies covered body acupuncture, ear acupuncture, scalp acupuncture, fluid acupuncture point injection and moxibustion. The studies were of varying quality. The weight of evidence, across all study types, is of beneficial effect from acupuncture, for those in an acute phase or those who have had Ménière's syndrome for a number of years. The review reinforces the importance of searching for studies from English and Chinese literature. The transferability of the findings from China to a Western context needs confirmation. Further research is also needed to clarify questions around the appropriate frequency and number of treatment/courses of acupuncture. The weight of evidence suggests a potential benefit of acupuncture for persons with Ménière's disease, including those in an acute phase and reinforces the importance of searching for published studies in the Chinese language.

Keywords: acupuncture – Ménière's syndrome – systematic review – Chinese literature

Introduction

Ménière's syndrome is a long-term, progressive disease that damages the balance and hearing parts of the inner ear. It is most common between the ages of 40 and 50 and rare in children and onset after the age of 60. Its incidence is about 1 in 1000, equally distributed between men and women and well documented in Caucasian, African-American and Asian races (1). The

etiology and treatment of Ménière's disease is not fully understood. In a literature-based, clinical review of the diagnosis and treatment of Ménière's disease, Saeed (2) commented that 'currently, the treatment of Ménière's disease is empirical. As yet, no treatment has prospectively modified the clinical course of the condition and thereby prevented the progressive hearing loss.' Indeed, Thorp *et al.* (3) in a critical review of studies of medical and surgical approaches queried whether any evidence-based medicine existed in the treatment of Ménière's disease.

Conventional biomedical treatments include drugs, diet and surgery. Drugs often form the first line of treatment.

For reprints and all correspondence: Andrew F. Long, School of Healthcare, University of Leeds, Room 3.10, Baines Wing, Leeds, LS2 9UT, UK. Tel: +44-113-343-6250; Fax: +44-113-343-7560; E-mail: a.f.long@leeds.ac.uk

© 2009 The Author(s).

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/2.0/uk/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

These include: diuretics (to reduce the amount of sodium in the body); drugs to block symptoms of motion sickness, nausea and vomiting, and anxiety and vertigo; systemic or local corticosteroids to reduce inflammation within the inner ear and to stop any immune reactions; antibiotics; and drugs to improve blood flow in the inner ear (4). The most commonly recommended dietary treatment is a low sodium diet. Patients have sometimes been helped by limiting certain components of their diet such as sugar, monosodium glutamate, caffeine and alcohol (4). Surgical treatment is the last resort of conventional medicine for Ménière's disease, the most controversial (2) and of greatest risk.

As with other chronic diseases where treatment may have serious side-effects (5), or is variably experienced by patients as beneficial (6), or as with Ménière's disease there is limited high quality evidence as to the most appropriate treatments (3), some who suffer from the condition may explore the potential of complementary and alternative medicine (CAM) therapies to alleviate their symptoms. However, there is little published information on which CAM therapies might help those with Ménière's disease. To begin to address this evidence gap, and following anecdotal comments about the potential benefits of one CAM therapy—traditional Chinese medicine (TCM) acupuncture—a systematic review was undertaken of published evidence on the effectiveness of acupuncture in treating the symptoms of Ménière's disease, drawing on both English and Chinese language sources. This article reports the findings from that review.

Methods

Searching

As there was no previous review of published evidence in this area, and thus to ensure comprehensive coverage of the literature, any published English or Chinese language study, which used a quantitative outcome measure and had a sample size of 10 or more, was included. The criterion for sample size was set low in anticipation that current literature in this field might have a relatively small sample size; an *a priori* prescribed higher limit might unduly restrict the located evidence. The search was undertaken in two parts. The first was based on a search of English language literature, undertaken in 2004/5 and forming the focus for a final year dissertation of an undergraduate student (K.M.) studying for a BSc in Traditional Chinese Medicine (Acupuncture) at the University of Salford; this was updated in 2009. The second comprised an opportunistic search of Chinese language literature as part of a study visit of one of the authors (M.X.) to China in the winter of 2003. The inclusion and exclusion criteria are summarized in Table 1.

Table 1. Inclusion and exclusion criteria

<i>Inclusion criteria:</i>	
Study on patients with a diagnosis of Ménière's disease	
All types of TCM acupuncture or moxibustion	
Study using acupuncture or moxibustion in addition to other TCM e.g. herbal medicine	
Studies of any controlled type and case series if the sample size was ≥ 10	
Studies in the English or Chinese languages	
<i>Exclusion criteria:</i>	
Studies with a sample size of <10 or single case reports or reports of opinion	
Studies which treated dizziness, tinnitus or other symptoms of Ménière's disease without a diagnosis of Ménière's	
Other forms of Traditional Chinese Medicine without acupuncture or moxibustion	

Six databases were searched: MEDLINE via OVID interface (1966 to January 2009); The Cochrane Controlled Trials Register (on 18 January 2009), EMBASE (1974 to January 2009); CINAHL (1982 to January 2009); AMED (1985 to January 2009); and, for the Chinese language review, the Chinese database of Science and Technology (1993 to August 2003). A search of *eCAM* was also undertaken (on 6 January 2009). Hand searching of Chinese journals related to acupuncture and moxibustion was undertaken in China for the months of September to December 2003. Search terms of the electronic databases focused on terms related to *acupuncture* and *Ménière's disease*. Free text search terms were used and controlled subject headings were exploded so that any subcategories were included (e.g. endolymphatic hydrops, acupuncture therapy or electroacupuncture). See Appendix 1 for an illustrative search strategy.

Appraisal and Analysis

Titles and abstracts were screened by one member (K.M. or M.X., respectively) and 27 papers were included in the review (Fig. 1). All included studies were critically appraised. For the English language studies, an established evaluation tool (7) was used, suitably adapted to add in appropriate search-specific and acupuncture-related aspects (8). For the Chinese language literature, data were extracted and simultaneously translated into English, using a standard template which included the following categories: study design; sample; treatment method; duration of treatment; outcome measurement; results; commentary on the appropriateness of acupuncture; and, overall comments/critique of the paper. Each English language paper was read and appraised by two members of the team independently (K.M. and M.X./A.F.L.) and any discrepancies resolved through discussion. The Chinese summaries were critically appraised (M.X. and A.F.L.) and where necessary further data extracted from the primary source.




Figure 1. Flowchart literature search process.

A narrative approach to data synthesis was taken, with greater weight given to studies of greatest quality (minimizing bias, accuracy of diagnosis, appropriateness of acupuncture treatment and use of appropriate outcome assessment). A meta-analysis was inappropriate due to difference in study designs and outcome measures and the limited amount of quantitative data for individual patients. Restriction of the analysis to only controlled trials would have substantially narrowed the available evidence base and was judged inappropriate in light of the lack of previous attempts to draw together evidence on acupuncture for Ménière's disease.

Results

Nature of the Evidence

Of the 27 studies included in the review, 9 were studies in the English language and 18 in the Chinese language. All but one of the studies was conducted in China.

The studies comprised: three randomized controlled trials (RCTs) (9–11); three non-randomized controlled studies (12–14); four pre-test, post-test designs (15–18); nine post-test designs (19–27) and eight case series reports (28–35). Together they comprised a total of 1888 patients in the acupuncture treatment arm. Seventeen of the studies followed the patients up for at least a year after the end of treatment.

Particular symptoms formed the focus of a number of studies, including vertigo (21) and dizziness (11,18,32). The patients had been suffering from Ménière's disease for various periods of time, ranging from 1 to 24 years. Thirteen of the studies included patients at an acute phase, within 1–10 days of an acute attack. Three studies (9,15,30) explicitly focused on the effects of acupuncture for acute symptom relief. Sample sizes varied from 15 (11) to 286 (33), and 5 had a sample size of less than 30 (18,19,25,26,30). The six controlled studies had total sample sizes ranging from 36 to 189 in the acupuncture arm. Nine of the studies had 50 or fewer participants.

The majority of studies employed a graded outcome measurement approach, differentiating three to four categories: 'cured', 'outstandingly effective', 'effective/improved' or 'not effective'. 'Cured' had the common meaning of 'dizziness and other symptoms having disappeared' or 'all symptoms disappeared', 'able to return to work/resume normal activities' and 'no recurrence within a 1 or 2 year period' (depending on the study's follow-up time). 'Outstandingly effective' had a similar meaning but with recurrence of symptoms 'occasionally', by, for example, 6 months. 'Effective/improved' related to 'relief of symptoms'. Particular symptoms were often mentioned as part of the achieved outcomes, in particular dizziness and vertigo.

Nature of Acupuncture

The studies covered five types of acupuncture: body acupuncture; ear acupuncture; scalp acupuncture; fluid acupuncture point injection; and moxibustion. All such types are classifiable as falling within the TCM style, ensuring the 'de-qi' sensation. Half of the studies involved an individualized TCM prescription approach to the treatment and half used a pre-set prescription, itself based on TCM principles and approaches to treating particular symptoms commonly experienced by persons with Ménière's disease, in particular, symptoms of dizziness, vertigo, nausea and vomiting. While none of studies mentioned the background of the acupuncturist, acupuncturists in China are all qualified medical doctors and at least 4 years training is required to work in a hospital.

No single, uniform number of courses or duration of treatment within each course was apparent. In two controlled studies (10,14), once daily acupuncture was undertaken for 10 days; in another (13), no indication of the number of treatments or how often the treatment was given; in yet another (30), patients continued with the treatment until symptom relief. Across the studies, a treatment 'once a day' for a course of 'up to 10 sessions' with the possibility of a second (or more) course was common.

Quality of the Evidence

A number of weaknesses were evident in the studies. First, as only six of the studies had a comparison group and four others also included pre-treatment measurement, any observed improved outcomes could have arisen because of other factors, including symptom remission. Secondly, only eight of the studies either included audiometric testing (9,19,21) or recruited Ménière's disease patients according to established criteria for Ménière's (10,18,23,24,26). Only two studies included audiometric testing as part of their outcome measurement (9,27). Thirdly, most of the Chinese language studies did

not provide detail on the inclusion/exclusion criteria. Fourthly, very limited detail was provided on the choice of study participants or setting of the studies. Fifthly, the length of follow-up (and thus final outcome measurement point) varied; eight studies had a follow-up time of 2 years and nine had 1 year, while for seven no or little detail was provided.

Evidence of Effectiveness

The weight of evidence, across all study types, is one of the beneficial effects from five types of acupuncture—body, ear or scalp acupuncture, fluid acupuncture point injection, or moxibustion (Table 2). The three randomized controlled trials (9–11) demonstrate a statistically significant benefit of body or scalp acupuncture against Western medicine and vitamins, with a mean difference in the total effectiveness percentage of 14% in favor of acupuncture ($P < 0.01$). Two non-randomized controlled trials, comparing body acupuncture against Western medicine (12) or Chinese herbal medicine (13), show a similar mean difference in total effectiveness percentage (19%, $P < 0.001$). The evidence from the four pre-test, post-test studies reports even higher success rates (90–100%) for acupuncture, including moxibustion on acupoint Du-20, either on its own (18) or combined with body (15), and/or ear acupuncture (16) or herbal medicine (17). High success rates are also reported from the post-test and case-series studies. As 13 studies included patients in an acute phase of Ménière's disease, the evidence suggests beneficial effects for both those in an acute phase and who have had Ménière's for a number of years. There is, however, insufficient evidence to recommend one, or another, type of acupuncture, nor the number of courses of treatment that might be needed for beneficial effect.

Discussion

This is the first systematic review of evidence on the potential benefits of acupuncture for the treatment of Ménière's disease and draws on both English and Chinese language sources. It demonstrates beneficial effects from up to five types of acupuncture (body, ear or scalp acupuncture, fluid acupuncture point injection, or moxibustion) and for both those in an acute phase and who have had Ménière's disease for a number of years. At the same time, it is important to note that the quality of evidence varies and is drawn from studies with varying potentiality for bias. In the five studies with a non-acupuncture comparison group (irrespective of the mode of allocation), the differences in total effectiveness percentage in favor of acupuncture were of the order of 14–19%.

Table 2. Evidence table (English and Chinese language studies)

Study, study type and country	Treatment, sample size and follow-up time	Confirmed diagnosis and time with Ménière's disease	Appropriateness of treatment	Key findings	Summary evaluative comments and overall quality
Randomized controlled trials Zhang <i>et al.</i> (9) China	Body acupuncture vs. Western medicine 15-day treatment <ul style="list-style-type: none"> • Acupuncture: once a day for 2–3 days; if symptom relief, then every other day; • Western medicine: two drugs (Serc and Vitamin B₃) and Vitamin B₆ ($n = 76$; $I = 39$; $C = 37$) Follow-up: 15 days (end of treatment period)	MD patients attending hospital for acute attack Unknown duration	No TCM pattern diagnosis used; set acupoints; TCM rationale provided	Total effectiveness rate: 74% vs. 49% ($P < 0.05$) 69% vs. 43% symptom control ($P < 0.05$) Small, audiometrically confirmed, hearing change	Strengths of the study include: confirmed MD, random allocation to treatment groups, detailed overview and rationale of the treatments, graded outcome measurements and audiometric testing. The short length of follow-up limits the generalization of the results. Overall quality judgment: Good
Gao and Ni (10) China	Scalp acupuncture and Western medicine vs. Western medicine and vitamin B ₁ and B ₁₂ <ul style="list-style-type: none"> • Acupuncture (or injected vitamins) once a day for 10 days as one course, total of three courses • Western medicine: once a day for 10 days, total of three courses ($n = 132$; $I = 58$; $C = 74$) Follow-up: 2 years	MD confirmed by two medical committees Up to 7 years	Set prescription without individualized diagnosis and treatment	Total effectiveness rate: 98% vs. 89% ($P < 0.05$) 89% vs. 27% symptom control ($P < 0.001$) 84% vs. 49% hearing improvement ($P < 0.001$)	Strengths of the study include: confirmed MD, random allocation to treatment groups, detailed overview of the treatments, and graded outcome measurement. The chosen treatment approach is experimental, rather than a traditional TCM one. Overall quality judgment: Good
Fang (11)* China	Acupuncture and drug treatment vs. drug treatment <ul style="list-style-type: none"> • Acupuncture—key points; once daily (acute stage), once every 3 days (remission stage). • Drug treatment (sibelum, 10 mg once daily for 30 days) ($n = 65$; $I = 36$; $C = 29$; of which Ménière's $I = 11$; $C = 4$) Follow-up: 30 days	Unclear if the MD cases confirmed MD 1 day to 3 months (from dizziness)	Individualized diagnosis and prescription	Total effectiveness rate for Ménière's group: 91% vs. 50% (NS, $P > 0.05$) 7 out of 11 vs. none of controls dizziness cured	Strengths of the study include: random allocation to treatment groups, individualized diagnosis, detailed overview of the treatments, and graded (and quantitative) outcome measurement. Weakness in very small number of MD sample size, non-confirmation of MD diagnosis and short follow-up time. Overall quality judgment: Fair
Non-randomized controlled trials Yú and Shi (12) China	Acupuncture vs. Western medicine and vitamin <ul style="list-style-type: none"> • Acupuncture—key and specific points. • Western medicine (vitamin C and Luminal (an anti-convulsive drug injection)) No details given on daily frequency or number of sessions of acupuncture ($n = 168$; $I = 86$; $C = 82$) Follow-up: 2 years	Unclear if confirmed MD 1–7 years	Treatment group has TCM diagnosis and individualized treatment; control group has set acupoints.	Total effectiveness rate: 100% vs. 74% ($P < 0.01$) 60% vs. 39% cured and no recurrence at 2 years ($P < 0.01$) 30% vs. 8% outstanding improvement and no recurrence at 1 year	Strengths include details over outcome measurement; length of follow-up; use of TCM pattern differentiation. Weaknesses include: lack of clarity over how patients are allocated to the two treatment groups; MD not confirmed; and limited detail over treatments (form or frequency). Overall quality judgment: Fair

(continued)

Table 2. Continued

Study, study type and country	Treatment, sample size and follow-up time	Confirmed diagnosis and time with Ménière's disease	Appropriateness of treatment	Key findings	Summary evaluative comments and overall quality
Yan (13)* China	Body acupuncture vs. Chinese herbal medicine (for vertigo) <ul style="list-style-type: none"> Acupuncture: Two courses of 15 sessions 1 p day, with 5 day rest Chinese herbal medicine: Two courses of 15 sessions 1 p day, with 5 day rest <i>n</i> = 257; <i>Ac</i> = 189; <i>Hb</i> = 68 Follow-up: 1 year	Patients with MD 102 patients had MD for ≤1 year, 135 1–5 years and 20 > 5 years	No TCM pattern diagnosis used to choose acupuncture Set acupuncture prescription	Total effectiveness rate: 99% vs. 88% (<i>P</i> < 0.01) 93% vs. 60% 'complete' response (<i>P</i> < 0.001) 99% vs. 88% 'complete' or 'partial' response 11% vs. 49% recurrence rate (<i>P</i> < 0.001) [when not indicated]	Short report, but adequate details on MD confirmation, treatment points, appropriate 2:1 case: comparison subject ratio and explanation of set acupuncture prescription. Lack of detail over how patients were allocated to the treatment and comparison group (non-random).
Qin and Jia (14) China	Scalp and ear acupuncture vs. body acupuncture <ul style="list-style-type: none"> Scalp and ear acupuncture once per day for 10 days, plus patients (taught to) self-treat for 5 days. Body acupuncture once a day for 10 days (<i>n</i> = 102; <i>I</i> = 72; <i>C</i> = 30) Follow-up: none	Confirmed MD by hospital Duration—no information	Set prescription without individualized diagnosis and treatment	Total effectiveness rate: 94% vs. 73% (<i>P</i> < 0.01) 53% vs. 33% cured (<i>P</i> < 0.01) 27% vs. 23% marked improvement	Strengths include details over acupuncture treatment and their rationales and outcome measurement. Weaknesses include: lack of follow-up; and lack of clarity over how patients are allocated to the treatment groups. Overall quality judgment: Fair
Pre-test, post-test design Li and Li (15) China	Acupuncture Once a day for 20 days <i>n</i> = 56 Follow-up: 1 year	Patients with MD in acute phase (all previously treated with Western medicine without effect) Up to 8 years	Set prescription, needing one point (Gang Shen), based on clear rationale	Total effectiveness rate: 100% 77% acute symptom control after one treatment; 16% after two. Most patients needed only 7 sessions to experience benefit.	Extensive details over treatment procedure and rationale are provided (Gang Shen is an experimental point for MD). But lack of detail over choice of participants and outcome measurement scale. Overall quality judgment: Fair
Li (16) China	Herbal medicine and ear acupuncture, for 2–60 days <i>n</i> = 90 Follow-up: 1 year	Patients with MD according to explicit criteria Up to 20 years	Individualised treatment following TCM principles for herbal medicine; ear acupuncture points set in conjunction with herbal medicine	Total effectiveness rate: 90% 72% cured 18% improved	Strengths include: clear diagnostic criteria, appropriate TCM principles and outcome measurement in relation to these principles and descriptions. But lack of detail over treatment duration or choice of participants. Note: this is predominantly a herbal approach with ear acupuncture as an adjunct. Overall quality judgment: Good
Dong and Zhou (17) China	Scalp acupuncture plus herbal medicine Once daily acupuncture for 7 days; simple herbal tea (to continue tea for 1 year) <i>n</i> = 180 Follow-up: 1 year	MD acute stage (hospitalized following acute attack) Up to 10 years	TCM individualized diagnosis and treatment, with herbal remedy adjusted	Total effectiveness rate: 97% 70% cured 18% outstanding effect 8% improved	Strengths include: clarity over outcome measurement, TCM diagnosis and treatment and length of follow-up. Overall quality judgment: Good
Sun and Li (18) China	Moxibustion at Du-20 Twice a day for 15 days <i>n</i> = 20 Follow-up: 1 year	Patients with clinically confirmed MD Up to 2 years	TCM individualized diagnosis with set acupoint (Du-20)	Total effectiveness rate: 100% 75% cured 25% outstandingly improved	Use of standard diagnosis criteria and outcome measurement criteria for MD; appropriate acupoint with TCM rationale. (Note: treatment also included advice on diet and emotion) Overall quality judgment: Good

Post-test design Dai and Liang (19)* China	Electro-acupuncture, acupuncture and moxibustion One treatment daily for up 6 days (30 min application) n=23 Follow-up: 1 year	MD patients who had auditory vertigo symptoms Up to 17 years	One set point for acupuncture, another for moxibustion; other points added according to TCM diagnosis	Total effectiveness rate: 100% 70% 'cured' 18% 'excellent'	Targets only cases with auditory vertigo syndrome. Detailed description of other Ménière's symptoms of cases; full explanation of potential role of acupuncture in treating Ménière's.
Liu (20) China	Acupuncture 10 sessions for 1 month n=51 Follow-up: 2 year	MD—query over criteria Duration—no information	Individualized treatment following TCM principles	Total effectiveness rate: 90% 22% cured 69% improved	Strengths include: appropriate TCM principles. But lack of detail over treatment duration, choice of participants and diagnostic criteria for MD. Overall quality judgment: Poor
Chao (21) China	Moxibustion at Du-20 (over 2 hours), for up to three sessions n=32 Follow-up: 2 years	Confirmed MD (Otol. Dept) average 9 years	Du-20 is a good experiential point for dizziness. Not diagnosed according to TCM theory. Note: moxibustion only	Total effectiveness rate: 100% 100% symptom relief (8 from one session, 10 from two, and 14 after three). No recurrence after 2 years.	Strengths include: MD confirmed by Otolaryngology Department, use of standard treatment duration, use of an appropriate (Du-20) treatment point and length of follow-up. Weaknesses include: unclear sampling criteria, sole focus on one MD symptom (dizziness) and lack of control group. Overall quality judgment: Fair
Tian (22)* China	Electro-acupuncture Two to three courses of 10 sessions daily, with 2 days rest between course (10–20 min application) n=72 Follow-up period: 1 year	Patients with MD Fifty had MD for 1–5 years, 22 for >5 years	Treatment according to TCM pattern diagnosis	Total effectiveness rate: 74% 50% 'marked improvement' 24% 'improved' Treatment more effective in cases of shorter duration	Short article; in consequence, limited details on methods, except on acupuncture treatment. Overall quality judgment: Fair
Bo (23) China	Acupoint injection at Du-20 Once a day for 10 days, 1 day break, then another course of treatment (3–30 sessions in total) n=88 Follow-up: not stated	Patients with MD according to 'book' of common diseases Up to 3 years	Set, single point used, with TCM rationale	Total effectiveness rate: 100% 75% cured 23% improved	Clear diagnostic criteria with detailed treatment procedure. But duration of follow-up is not stated and lack of clarity over outcome measurement (causes of dizziness disappeared). Overall quality judgment: Fair
Wang and Chen (24) China	Acupuncture points injection with Dansheng liquid on one side at one time plus scalp acupuncture Acupuncture with injection once a day, with scalp acupuncture every other day, for 10 days. 5-day break, second course of treatment n=50 Follow-up: 2 years	Patients with MD Up to 14 years	Set prescription but based on good TCM treatment principles	Total effectiveness rate: 92% 62% cured 30% outstandingly improved	Combined traditional and scalp needling plus points injection seems a potentially good treatment method, and based on TCM principles. While diagnosis is based on TCM principles, treatment is not individualized. Good detail on treatment procedures. Overall quality judgment: Fair
Zhang (25)* China	Body acupuncture Courses of 10 sessions with a 2-day rest between (30 min application) n=18 Follow-up period: 2 years	Patients with MD Up to 6 years.	Treatment according to TCM pattern diagnosis	Total effectiveness rate: 94% 72% cured 22% marked effect	Study report is from an abstract, translated from the original Chinese paper. Brief detail on methods is provided, with extensive detail on needling and TCM rationale. Overall quality judgment: Fair

(continued)

Table 2. Continued

Study, study type and country	Treatment, sample size and follow-up time	Confirmed diagnosis and time with Ménière's disease	Appropriateness of treatment	Key findings	Summary evaluative comments and overall quality
Zhou (26) China	Acupuncture 3–15 sessions, once per day (average of 9) <i>n</i> = 20 Follow-up: 1 year	MD—query over criteria Up to 8 years	TCM pattern differentiation	Total effectiveness rate: 100% 80% cured 20% improved	Lack of detail over choice of participants or diagnostic criteria for MD. While diagnosis is individualized, the treatment points are only for one TCM diagnosis pattern. It is unclear if the treatment is individualized. Overall quality judgment: Poor
Zhang (27) China	Acupuncture Once a day for 10 days, 5 day break, another course <i>n</i> = 60 Follow-up: 2 years	MD—query over criteria Up to 10 years	Set acupoints based on explicit TCM rationale for point choice	Total effectiveness rate: 92% 60% cured 30% outstandingly improved	Strengths include: explicit rationale for set acupoint prescription, based on TCM theory, and extensive details over treatment methods. Lack of clarity over how many courses were provided and diagnostic criteria. Overall quality judgment: Poor
Case series					
Steinberger and Pansini (28)* ?Yugoslavia	Body acupuncture, ear acupuncture and moxibustion (and, rarely, scalp acupuncture) Five treatments, once a day [30 min application] <i>n</i> = 34 Follow-up period: unclear	Patients with MD—with confirmed auditory tests	Treatment according to TCM pattern diagnosis	Total effectiveness rate: 100% 100% success for vertigo after 3 treatments	Very brief report on cases over 5-year period, with limited detail on outcome measurement (focus on vertigo). Strengths include: the use of audiometric tests (for auditory acuity) and extensive detail on needling and TCM treatment rationale. Overall quality judgment: Poor
Xu and Ge (29)* China	Body acupuncture Courses of 10 sessions, once a day (20–30 min application) <i>n</i> = 75 Follow-up period: 1 year	Patients with MD according to clinical symptoms 18 had had MD for <1 year, 57 for ≥ 1 year	Set acupoint prescription, with extra points if tinnitus and deafness were severe	Total effectiveness rate: 93% 39% cured 33% 'marked' improvement 21% improved On average, needed two treatment courses (range 1–5)	Reporting on 18-year case series of clinically confirmed MD cases. Short report, with limited but adequate detail on the acupuncture treatment. Overall quality judgment: Poor
Tian (30)* China	Body acupuncture 10–15 treatments <i>n</i> = 21 Follow-up period: unstated	Patients with MD, with a 'sudden onset' of symptoms	Treatment according to TCM pattern diagnosis	Total effectiveness rate: 100% 100% 'cured' [Eight had alleviation of symptoms after four treatments; 13 after 7]	Brief report on cases treated since 1975 with very limited detail, except on rationale and approach to treatment (and ways to improve the treatment). Overall quality judgment: Poor

Song and Yi (31) China	Acupuncture Once a day until all symptoms disappear (mean = 5; range 2 to >10) n = 152 Follow-up: unclear	MD—query over criteria Up to 15 years	Treatment following TCM principles, but only for the one symptom of dizziness	Total effectiveness rate: 99% 91% cured over short term 8% improved	Strengths include: treatment based on appropriate TCM principles. But lack of detail over choice of participants or diagnostic criteria for MD, treatment not individualized, no standard course of treatment, and unclear length of follow-up Overall quality judgment: Poor
Zhu (32) China	Acupuncture n = 51 Follow-up: not indicated	Patients with MD at an acute stage Up to 6 years	No TCM diagnosis differentiation or individualized treatment	Total effectiveness rate: 98% 64% cured 32% outstanding effect	Following appropriate TCM principles for treating two symptoms (dizziness and vomiting) of MD, but limited detail and no explicit follow-up mentioned. Overall quality judgment: Poor
Zhang and Shang (33) China	Acupuncture Once a day for 3 days (as one course) n = 286 Follow-up: 1 year	Patients with MD Up to 24 years	Appropriate acupuncture at single point. No TCM pattern differentiation	Total effectiveness rate: 90% 57% cured 28% outstanding effect	The study reports on a large group of patients; the treatment provided is an integrated treatment for MD, but there is a lack of detail over the number of courses given. Overall quality judgment: Fair
Lu (34)* China	Body acupuncture and moxibustion One treatment per day [30 min application]; 2–14 daily treatments given (mean of 7.4) n = 30 Follow-up: 1 year	Patients with MD Twenty-two patients had MD ≤10 years; eight for >10	Treatment according to TCM pattern diagnosis	Total effectiveness rate: 100% 87% cured 13% effective 30% symptom free after 1 year; 47% re-occurrence within 6 months	Short report on cases treated over a number of years. Argues that Ménière's syndrome belongs to the category of 'dizziness' in TCM. Strengths include one year follow-up and clear rationale for treatment. Overall quality judgment: Fair
Wang (35) China	Acupuncture plus moxibustion at Du-20 Once a day for 7 days as one course of treatment n = 30 Follow-up: 2 years	MD—query over criteria Up to 10 years	TCM pattern differentiation and treatment	Total effectiveness rate: 100% 97% cured	Reporting on cases treated since 1988 for TCM diagnosis of dizziness. Overall quality judgment: Fair

Asterisk indicated studies located from English language search. RCT: randomized controlled trial; CT: controlled trial; MD: Ménière's Disease; I: Intervention; C: Comparison/Control.

Given that the weight of evidence is from China, further research within a Western context is required. There is also a need to clarify questions around the appropriate frequency and number of treatments or courses of acupuncture. This is especially important where persons have to pay for acupuncture treatment. Exploring possible sets of pre-defined acupuncture points is also important, given that not all acupuncturists in the UK or Europe adopt a TCM diagnosis and treatment approach (36).

The review differs in relation to others in the CAM area, as it draws on both English and Chinese literature. Such an approach led to the inclusion of 18 additional studies and, in principle (irrespective of study quality), strengthens the conclusions of the review. A similar conclusion is drawn by Yuan *et al.* (37) who explored treatment regimens of acupuncture for low back pain, accessing English and Chinese literature, the latter located by Chinese experts. Indeed, their systematic review identified substantive differences in the treatment frequency between practice in China and the West.

The inclusion of English and Chinese language papers does not however avoid the issue of publication bias; publications in other languages would still be excluded. Neither does it avoid publication bias arising from the greater likelihood of the reporting of positive results (i.e. the non-reporting of negative studies). As Moher *et al.* (38) argued, substantial bias may occur in the results of a CAM systematic review if languages other than English are excluded. Extending the searching process to the Chinese language is an important step, given the lengthy historical tradition and use of acupuncture within China. Indeed, accessing research literature published in the Chinese language perhaps needs to become a *sine qua non* in this type of literature review. This is challenging, requiring access to Chinese language readers (39,40). It was possible in this study only because one of the project team members (M.X.) was native Chinese. Optimal methods of accessing and including Chinese language literature however need further exploration.

Generalizing or transferring the results of studies undertaken in another country and cultural context is not straightforward. All but one of the studies took place in China. Citizens' attitudes to TCM and acupuncture in a Chinese cultural context, where there is a long tradition, recognition and establishment within the health care sector (36), along with established education and training approaches, are likely to be very different than in a Western European context. This has important consequences for patients deciding when to seek acupuncture. In China, acupuncture may be sought early or as a routine treatment. In the West, where the first line of treatment for someone with Ménière's disease would likely be biomedical, acupuncture may be sought only when the condition becomes chronic. There are also differences in education and training of, and access to,

acupuncturists (traditional TCM trained practitioners or Western acupuncture (41), thus the recognized importance of reporting studies using the STRICTA framework (8). This strongly suggests the need for a controlled trial of acupuncture, for both persons in an acute phase and those who have had Ménière's disease for a number of years.

Any such study needs to be appropriately designed. As there is no definitive, demonstrated bio-medical curative approach for Ménière's disease, choice of comparison group is not straightforward. A placebo design would seem both inappropriate and ethically problematic, as it is highly improbable that potential study participants will not be taking some medication or other approach to enable them to cope. While arguing for the need for greater and more rigorous evidence, Thorp *et al.* (3) make a similar comment in the context of conventional medicine. In Europe, an appropriate comparison might be drug treatment. In China, an appropriate contrast might be a (Western) drug approach or different forms of TCM approaches (e.g. types of acupuncture and/or herbal medicine). Indeed, the former form of comparison (acupuncture vs. a drug or herbal approach) was used in four of the included controlled studies (9–12), and the latter (different forms of acupuncture or Chinese herbal medicine) in two others (13,14). It is important to note that choice of comparison group should be culturally determined.

Whatever choice is made, it is important that full methodological details are provided in the research report, utilizing the STRICTA recommendations (8) both as a guide to reporting but also study design. The review suggests the potential of exploring treatment 'once a day' for a course of 'up to 10 sessions' with the possibility of a second (or more) course. Such research should measure both the short- and longer-term symptom effect, based upon patient reports of symptom benefit and (time before any) recurrence and extent of severity/symptom relief at any recurrence, general health and well-being effects and a potential greater resolve to cope with the illness.

There are some limitations to this review. First, the Chinese literature search accessed only one database and limited hand searching. More studies might be generated from a wider search. Secondly, it is notable that the studies from China report positive effects of acupuncture. This may be due to publication bias, (42) or be a true effect, demonstrating that frequent and a large number of treatment sessions, evident in the Chinese studies, leads to better outcomes (37). In this regard it is interesting to note one of the findings from a systematic review of clinical trials on acupuncture in the Japanese literature (43); studies undertaken pre-1990 tended to focus on the appropriate choice of acupuncture technique, whereas post-1990 the focus tended to be on evaluation of acupuncture efficacy. This finding has some

resonance within the Chinese literature in this review, especially within the case-series studies and their presentation and the depth of explanation about technique. Thirdly, inclusion of non-randomized and uncontrolled studies in the review could be argued to be problematic (44). Such studies were used here as a means to support, or otherwise, findings from more powerful studies. In addition, such studies provided additional information on the maintenance of beneficial effects. Fourthly, in a recent report relating specifically to Cochrane systematic reviews of TCM, where only 5 of the 28 described RCTs could be authenticated as RCTs, Wu *et al.* (45) argue for the need for review authors to verify with study authors for claims that their RCTs were in fact randomized. This was not undertaken in this study. Fifthly, no attempt is made to draw together any of the TCM explanations indicated in the included papers about the mechanisms that might underpin acupuncture effecting the symptoms of Ménière's disease or other factors which may be associated with such symptoms.

Conclusion

The aim of this systematic review was to locate and critically appraise evidence for acupuncture as a treatment for Ménière's disease, drawing from both English and Chinese language literature. Despite the range in the quality of the located evidence, the overall conclusion is of the potential benefit of acupuncture for persons with Ménière's disease, including those in an acute phase. The review also demonstrates the importance of searching for studies in the Chinese language for such a therapy as acupuncture, given its lengthy historical tradition within Chinese medicine. As all but one of the studies took place within China, further research is needed in a Western health care context and to examine the frequency and number of treatment or courses of acupuncture.

Acknowledgements

The authors gratefully acknowledge the comments of Tony Bennett, as expert patient.

Funding

Research Development Fund of the University of Salford.

References

- Ferrari M and PDXMD. *PDXMD Ear, Nose & Throat Disorders*. Philadelphia: Elsevier Science, 2003.
- Saeed SR. Diagnosis and treatment of Ménière's Disease. *Br Med J* 1998;316:368–72.
- Thorp MA, Shehab ZP, Bance ML, Rutka JA. Does evidence-based medicine exist in the treatment of Ménière's disease? A critical review of the last decade of publications. *Clin Otolaryngol* 2000;25:456–60.
- Salt A. *Compilation of Treatments for Ménière's Disease*. [Online] Washington University Cochlear Fluids Research Laboratory: St. Louis, 2004. Available at URL: <http://oto.wustl.edu/men/mtreat.htm> [Accessed July 2008].
- Vincent C, Furnham A. Why do patients turn to complementary medicine. *J Clin Psychol* 1996;35:37–48.
- Thorne S, Paterson B, Russell C, Schultz A. Complementary/alternative medicine in chronic illness as informed self-care decision making. *Int J Nurs Stud* 2002;39:671–83.
- Reisch JS, Tyson JE, Mize SG. Aid to the evaluation of therapeutic studies. *Pediatrics* 1989;85:815–27.
- MacPherson H, White A, Cummings M, Jobst K, Rose K, Niemtow R. Standards for reporting interventions in controlled trials of acupuncture: the STRICTA recommendations. *Acupunct Med* 2002;20:22–5.
- Zhang ZF, Xue F, He ZD, Hou S, Li H. Clinical study on acupuncture and moxibustion for treatment of acute onset stage of Ménière's syndrome. *Shanghai J Acupunct Moxi* 1983;2: 28–31.
- Gao XP, Ni HH. Observation on therapeutic effect of scalp-acupuncture on Ménière's Disease. *Chin Acupunct Moxi J* 2002;22:583–4.
- Fang G. Observation on treatment of dizziness mainly by acupuncture. *J Trad Chin Med* 2007;27:16–8.
- Yu S and Shi X (1997) Acupuncture for the treatment of 86 cases of Ménière's syndrome. *Shanghai Acupunct J* 2002; 16: 21.
- Yan SM. Acupuncture for Ménière's syndrome: short- and long-term observation of 189 cases. *Int J Clin Acupunct* 1999;10: 303–4.
- Qin YX, Jia XC. Combination of scalp acupuncture and ear acupuncture for the treatment of 78 cases of Ménière's syndromes. *Mod Chin Med J* 2003;6:52–3.
- Li CX, Li CZ. Clinical observation of acupuncture for the treatment of 56 cases of Ménière's syndrome. *J Acupunct Clin Pract* 1993;9:10.
- Li XH. Acupuncture in combination of herbal medicine for the treatment of 90 Ménière's Syndrome. *Liaoning J Trad Chin Med* 1999;26:220.
- Dong JJ, Zhou Z. Dizziness decoction combine with scalp acupuncture for the treatment of 180 cases of Ménière's syndrome. *Sanxi J Chin Med* 2001;22:470–1.
- Sun LJ, Li AX. Pressing moxibustion on D20 for the treatment of Ménière's Syndrome. *Chin Acupunct Moxi* 2001;21:10.
- Dai J, Liang S. Treatment of Ménière's disease by needling Ear-Shenmen and moxibustion at Baihui. *Int J Clin Acupunct* 1993;4:281–4.
- Liu Y. Acupuncture for the treatment of 51 cases of Ménière's Syndrome. *Shanghai J Acupunct* 1995;14:258.
- Chao LY. Moxibustion on Du-20 for the treatment of 32 cases of Ménière's syndrome. *Xing Jiang J Chin Med* 1996;15:28.
- Tian ZM. Treatment by electro-acupuncture – Ménière's disease. *J Chin Med* 1999;61:28–9.
- Bo SL. Injecting D20 for the treatment of 88 cases of Ménière's syndrome. *Chin Acupunct Moxi* 2002;22:72.
- Wang QQ, Chen HL. Combination of acupuncture points injection with scalp acupuncture for the treatment of Ménière's Syndrome. *J Sichuan Trad Chin Med* 2002;20:73.
- Zhang Z. The treatment of 18 cases of Ménière's disease mainly with the points Bai Hui (GV-20) joined to Hou Ding (GV-19). *Ji Lin Zhong Yi Yao (Jilin Chin Med Medicinals)* 2002; 9: 48. Précis in English, Wolfe H. Acupuncture for Ménière's disease. *Townsend Lett* 2003;244:146–7.
- Zhou FH. Acupuncture for the treatment of 20 Ménière's syndrome. *J Chin Acupunct Moxi* 2002;18:26.
- Zhang XJ. Acupuncture for the treatment of 60 cases of Ménière's syndrome. *Sanxi Chin Med J* 2003;24:545.
- Steinberger A, Pansini M. The treatment of Ménière's disease by acupuncture. *Am J Chin Med* 1983;11:102–105.
- Xu B, Ge S. Treatment of Ménière's disease by acupuncture: report of 75 cases. *J Trad Chin Med* 1987;7:69–70.
- Tian C. Therapeutic effect of Ménière's syndrome treated by acupuncture. *Chin J Acupunct Moxi* 1991;4:5–6.
- Song JJ, Yi SY. Acupuncture for the treatment of 152 cases of Ménière's syndrome. *Acupunct Chin* 1992;6:152.

32. Zhu XL. Treatment of Ménière's syndrome with acupuncture. *Jilin Chin Med J* 1995;1:11.
33. Zhang LS, Shang SP. Injection to the acupuncture points for the treatment of 286 cases of Ménière's syndrome. *Shanghai Acupunct Moxi J* 1996;15:22.
34. Lu ZH. Ménière's syndrome treated with acupuncture: a report of 30 cases. *Int J Clin Acupunct* 1997;8:295–7.
35. Wang DF. Moxibustion for the treatment of 30 cases of Ménière's syndrome. *Acupunct Chin* 1999;3:163.
36. Hughes JG, Goldbart J, Fairhurst E, Knowles K. Exploring acupuncturists' perceptions of treating patients with rheumatoid arthritis. *Complement Ther Med* 2007;15:101–8.
37. Yuan J, Kerr D, Park J, Liu XH, McDonough S. Treatment regimens of acupuncture for low back pain – a systematic review. *Complement Ther Med* 2008;16:295–304.
38. Moher D, Pham B, Lawson ML, Klassen TP. The inclusion of reports of randomised trials published in languages other than English in systematic reviews. *Health Tech Assess* 2003;7:1–77.
39. Pilkington K, Richardson J. Exploring the evidence: the challenges of searching for research on acupuncture. *J Altern Complement Med* 2004;10:587–90.
40. Shekelle PG, Morton SC, Suttrop MJ, Buscemi N, Friesen C. Challenges in systematic reviews of complementary and alternative medicine topics. *Ann Int Med* 2005;142:1042–7.
41. Acupuncture Regulatory Working Group. *The Statutory Regulation of the Acupuncture Profession: the Report of the Acupuncture Regulatory Working Group*. London: The Prince of Wales Foundation for Integrated Health, 2003.
42. Vickers A, Goyal N, Harland R, Rees R. Do certain countries produce only positive results? A systematic review of controlled trials. *Control Clin Trials* 1998;19:159–66.
43. Tsukayama H, Yamashita H. Systematic review of clinical trials on acupuncture in the Japanese literature. *Clin Acupunct* 2002;3:105–13.
44. Linde K, Scholz M, Melchart D, Willich SN. Should systematic reviews include non-randomized and uncontrolled studies? The case of acupuncture for chronic headache. *J Clinl Epidem* 2002;55:77–85.
45. Wu T, Zhuo Q, Liu J, Yang X, Moher D. Assessing the quality of Cochrane systematic reviews of Traditional Chinese Medicine. *XV Cochrane Colloquium* Oct 23–27, Sao Paulo: Brazil, 2007.

Received September 24, 2008; accepted May 5, 2009

Appendix 1. Search Strategy for MEDLINE via OVID interface

1. meniere\$.mp.
2. (endolymphatic adj hydrops).mp.
3. exp Endolymphatic Hydrops/
4. exp Meniere Disease/
5. 1 or 2 or 3 or 4
6. exp Acupuncture Therapy/
7. (acupuncture or electro?acupuncture or auriculotherapy or (auricular adj needles) or (needling adj therapy) or (electric adj needling)).mp.
8. exp Medicine, Chinese Traditional/
9. 6 or 7 or 8
10. 9 and 5

Key:
 mp = title, original title, abstract, name of substance word, subject heading word
 exp = explode subject heading
 / = subject heading (MESH) search
 \$ = truncation symbol
 ? = wildcard symbol
 adj = adjacent to
