



University of  
**Salford**  
MANCHESTER

# Photoreactivity of biologically active compounds. XIX: Excited states and free radicals from the antimalarial drug primaquine

Kristensen, S, Edge, R, Tønnesen, HH, Bisby, RH and Navaratnam, S

<http://dx.doi.org/10.1016/j.jphotobiol.2008.11.006>

<b>Title</b>	Photoreactivity of biologically active compounds. XIX: Excited states and free radicals from the antimalarial drug primaquine
<b>Authors</b>	Kristensen, S, Edge, R, Tønnesen, HH, Bisby, RH and Navaratnam, S
<b>Type</b>	Article
<b>URL</b>	This version is available at: <a href="http://usir.salford.ac.uk/id/eprint/1739/">http://usir.salford.ac.uk/id/eprint/1739/</a>
<b>Published Date</b>	2009

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: [usir@salford.ac.uk](mailto:usir@salford.ac.uk).

## Photoreactivity of biologically active compounds. XIX: Formation and reactivity of excited states and free radicals from primaquine

Solveig Kristensen<sup>\*a</sup>, Ruth Edge<sup>b,c,#</sup>, Hanne Hjorth Tønnesen<sup>a</sup>, Roger Hugh Bisby<sup>d</sup>, Suppiah Navaratnam<sup>b,d</sup>

<sup>a</sup>University of Oslo, School of Pharmacy, Department of Pharmacy, PO Box 1068 Blindern, NO-0316 Oslo, Norway.

E-mail addresses: [solveig.kristensen@farmasi.uio.no](mailto:solveig.kristensen@farmasi.uio.no); [h.h.tonnesen@farmasi.uio.no](mailto:h.h.tonnesen@farmasi.uio.no)

<sup>b</sup>STFC Daresbury Laboratory, Warrington WA4 4WD, UK.

E-mail address: [s.navaratnam@dl.ac.uk](mailto:s.navaratnam@dl.ac.uk)

<sup>c</sup>School of Physical and Geographical Sciences (Chemistry), Keele University, Keele, Staffs, ST5 5BG, UK

<sup>d</sup>Biomedical Sciences Research Institute, Peel Building, University of Salford, Salford M5 4WT, UK

E-mail address: [R.H.Bisby@salford.ac.uk](mailto:R.H.Bisby@salford.ac.uk)

<sup>#</sup> Present Address: School of Chemistry, University of Manchester, Oxford Road, Manchester, M13 9PL, UK.

E-mail address: [Ruth.Edge@manchester.ac.uk](mailto:Ruth.Edge@manchester.ac.uk)

\*Corresponding author. Address: University of Oslo, School of Pharmacy, Department of Pharmacy, PO Box 1068 Blindern, NO-0316 Oslo, Norway. Tel.: +47 22854218; fax: +47 22854402; E-mail address: [solveig.kristensen@farmasi.uio.no](mailto:solveig.kristensen@farmasi.uio.no)

### Abstract

The formation and reactivity of excited states and free radicals from primaquine was studied in order to evaluate the primary photochemical reaction mechanisms. The excited primaquine triplet was not detected, but is likely to be formed with a short lifetime ( $< 50$  ns) and with a triplet energy  $< 250$  kJ/mol as the drug is an efficient quencher of the fenbufen triplet and the biphenyl triplet, and forms  $^1\text{O}_2$  by laser flash photolysis ( $^{\text{PQ}}\Phi_{\Delta} = 0.025$ ). Primaquine photoionises by a biphotonic process and also forms the monoprotonated cation radical ( $\text{PQH}^{2+\bullet}$ ) by one electron oxidation by  $\text{OH}\cdot$  ( $k_q = 6.6 \cdot 10^9 \text{ M}^{-1} \text{ s}^{-1}$ ) and  $\text{Br}_2\cdot^-$  ( $k_q = 4.7 \cdot 10^9 \text{ M}^{-1} \text{ s}^{-1}$ ) at physiological pH, detected as a long-lived transient decaying essentially by a second order process ( $k_2 = 7.4 \cdot 10^8 \text{ M}^{-1} \text{ s}^{-1}$ ).  $\text{PQH}^{2+\bullet}$  is scavenged by  $\text{O}_2$ , although at a limited rate ( $k_q = 1.0 \cdot 10^6 \text{ M}^{-1} \text{ s}^{-1}$ ). The reduction potential ( $E^\circ$ ) of  $\text{PQH}^{2+\bullet} / \text{PQH}^+$  is  $< +1015$  mV. Primaquine also forms  $\text{PQH}^{2+\bullet}$  at pH 2.4, by one electron oxidation by  $\text{Br}_2\cdot^-$  and proton loss ( $k_q = 2.7 \cdot 10^9 \text{ M}^{-1} \text{ s}^{-1}$ ). The non-protonated cation radical ( $\text{PQ}^+\cdot$ ) is formed during one electron oxidation with  $\text{Br}_2\cdot^-$  at alkaline conditions ( $k_q = 4.2 \cdot 10^9 \text{ M}^{-1} \text{ s}^{-1}$  at pH 10.8). The estimated  $\text{pK}_a$ -value of  $\text{PQH}^{2+\bullet} / \text{PQ}^+\cdot$  is  $\text{pK}_a \sim 7$ -8. Primaquine is not a scavenger of  $\text{O}_2\cdot^-$  at physiological pH. Thus self-sensitization by  $\text{O}_2\cdot^-$  is eliminated as a degradation pathway in the photochemical reactions. Impurities in the raw material and photochemical degradation products initiate photosensitized degradation of primaquine in deuterium oxide, prevented by addition of the  $^1\text{O}_2$  quencher sodium azide. Photosensitized degradation by formation of  $^1\text{O}_2$  is thus important for the initial photochemical decomposition of primaquine, which also proceeds by free radical reactions. Formation of  $\text{PQH}^{2+\bullet}$  is expected to play an essential part in the photochemical degradation process in a neutral, aqueous medium.

### Keywords

Primaquine, photoreactivity, free radicals, excited states, pulse radiolysis, laser flash photolysis

## Abbreviations

Abs	Absorption at given wavelength
BP	Biphenyl
Em.	Emission wavelength
Ex.	Excitation wavelength
$E^\circ$	Reduction potential
FEN	Fenbufen
$k_1$	First order rate constant
$k_2$	Second order rate constant
$k_{\text{obs}}$	Observed rate constant
$k_q$	Bimolecular quenching rate constant
$k_{\text{ref}}$	Rate constant of reference compound
M	Molar per litre
PQ	Primaquine
PQ <sub>ox</sub>	Oxidized degradation products of primaquine
${}^{\text{PQ}}\Phi_{\Delta}$	Quantum yield of ${}^1\text{O}_2$ -formation sensitized by primaquine
${}^{\text{PN}}\Phi_{\Delta}$	Quantum yield of ${}^1\text{O}_2$ -formation sensitized by perinaphtenone
Sens	Photosensitizing impurities
TRPH	Tryptophan
TYR	Tyrosine
$\epsilon$	Molar absorption coefficient
$\lambda$	Wavelength
$\lambda_{\text{max}}$	Absorption maximum
$\tau$	Lifetime of transient
$\Phi_{\text{F}}$	Fluorescence quantum yield
$\tau_{\text{F}}$	Fluorescence lifetime

## 1. Introduction

The 8-aminoquinoline primaquine is an antimalarial compound that acts as a tissue schizontocide. The drug is mainly used curatively to eliminate latent liver forms of *Plasmodium ovale* and *P. vivax*, since severe haematological adverse effects (methaemoglobinaemia and haemolysis) prohibit use for general prophylaxis [1]. Primaquine is a photolabile compound, with a ground state absorption maximum at 350 nm and extension of the absorption spectrum towards 430 nm under physiological conditions. The drug decomposes photochemically into several degradation products at physiological pH, by oxygen dependent reaction mechanisms [2-4]. Primaquine forms an intramolecular hydrogen bond, which seems to be essential for chemical and biological properties, including photochemical stability and fluorescence. Photostability in protic media is well correlated with fluorescence properties, and quenching by breaking of the intramolecular bond and protonation of the quinoline nitrogen in the excited singlet state appears to be an important factor influencing photochemical stability of primaquine in pure, protic media [5].

Primaquine is an *in vitro* photosensitizer, inducing photohaemolysis, photopolymerization of lens proteins and photoreduction of cytochrome C [3, 6, 7]. Free radicals ( $\text{OH}\cdot$  and  $\text{O}_2\cdot^-$ ) that are formed in the photochemical reactions are important mediators also in the *in vitro* oxidation of haemoglobin induced by primaquine [8, 9]. The haematological adverse effects observed after medication with primaquine can (partly) be ascribed to photochemical reactions of the drug *in vivo*, taking into account that Caucasian skin transmits optical radiation above 320 nm [10].

The present work was undertaken to identify and quantitatively characterize the primary transient excited states and free radicals from primaquine, with the aim to obtain further knowledge on the kinetics and mechanisms of primaquine photodecomposition.

## 2. Materials and methods

Primaquine diphosphate (98-99 % pure) was purchased from Sigma-Aldrich. Primaquine-base was produced by extraction with  $\text{CHCl}_3$  from an alkaline aqueous solution of the phosphate salt. The water used was deionised and purified.  $\text{D}_2\text{O}$  (99.9 % atom D) was purchased from Sigma-Aldrich. All other chemicals and organic solvents were of p.a. grade.

### 2.1. Preparation of samples for laser flash photolysis and pulse radiolysis

Solutions were saturated with high purity  $\text{N}_2$ ,  $\text{N}_2\text{O}$ ,  $\text{O}_2$  or air as appropriate by flushing with gas for at least 30 min prior to measurements. Phosphate buffer was prepared using  $\text{NaH}_2\text{PO}_4$  and  $\text{Na}_2\text{HPO}_4$ .

For detection of the primaquine-triplet by laser flash photolysis studies, primaquine diphosphate (0.2 – 0.4 mM) was dissolved in aqueous solutions at pH 7.4 and 9.3 saturated with  $\text{N}_2$  or air in the presence and absence of phosphate buffer (10 mM); in  $\text{N}_2$ -saturated methanol; and in  $\text{O}_2$ -saturated  $\text{D}_2\text{O}$  at pH 8-9. Primaquine-base ( $9.1\cdot 10^{-5}$  -  $3.1\cdot 10^{-4}$  M) was dissolved in toluene saturated with  $\text{N}_2$  or  $\text{O}_2$  and in  $\text{N}_2$ -saturated acetonitrile. Benzophenone dissolved in acetonitrile was used as an actinometer [11].

Primaquine-base was dissolved in O<sub>2</sub>-saturated toluene ( $^{PQ}Abs_{355} = 0.25$ ) for measurements of <sup>1</sup>O<sub>2</sub> quantum yield by laser flash photolysis. Perinaphthenone in air-saturated toluene ( $^{PN}Abs_{355} = 0.25$ ) was used as reference [12].

Primaquine-base (0.1-0.3 mM) was dissolved in N<sub>2</sub>-saturated acetonitrile in the presence of fenbufen (6 mM) for the study of energy transfer by laser flash photolysis. Fenbufen (6 mM) in acetonitrile was used as reference [13].

Primaquine-base (93 μM) was dissolved in N<sub>2</sub>-saturated toluene in the presence of biphenyl (10 mM) for the study of energy transfer by pulse radiolysis. Biphenyl (10 mM) in N<sub>2</sub>-saturated toluene was used as reference [14].

Primaquine diphosphate (5.0 μM - 0.2 mM) was dissolved in N<sub>2</sub>O-saturated 10 mM phosphate buffer pH 7.4 for investigation of the reaction with the hydroxyl radical OH• by pulse radiolysis. Samples were added 10 mM KBr to verify the formation of PQH<sup>2+</sup>• and characterize the radical. Samples were saturated with O<sub>2</sub> or N<sub>2</sub>O, and mixed to concentration ranges of O<sub>2</sub>:N<sub>2</sub>O ~ 0:100, 20:80 and 75:25, to evaluate the lifetime of PQH<sup>2+</sup>• as a function of sample oxygen concentration. The pKa-value and cation radicals of primaquine diphosphate (10 μM and 0.5 mM) formed at various degree of protonation were studied in N<sub>2</sub>O-saturated samples at pH 2.3 - 11.5. Perchloric acid and NaOH were used for pH adjustments.

The reduction potential under physiological conditions was evaluated by pulse radiolysis of primaquine diphosphate (0.5 mM) in N<sub>2</sub>O-saturated 10 mM phosphate buffer pH 7.4 containing 0.1 M KBr, and 5 mM tryptophan (TRPH) or 2 mM tyrosine (TYR).

In order to study the reduction of primaquine by superoxide anions at physiological pH, primaquine diphosphate (0.2 mM) was studied by pulse radiolysis in N<sub>2</sub>-saturated 10 mM phosphate buffer pH 7.4 containing 1% t-butanol.

## 2.2. *Laser flash photolysis*

The laser flash photolysis experiments were carried out with 15 ns pulses from a J.K. Laser System 2000 Neodymium/YAG oscillator with a Neodymium/glass amplifier. The power dependence measurements used 5 ns pulses from a Continuum Surelight II-10 Nd-YAG laser. Both lasers delivered up to 100 mJ of 355 nm radiation in unfocussed single pulses. The detection system consisted of a water cooled Xe arc lamp and a pulsing unit, high radiance monochromator and quartz optics (Applied Photophysics Ltd.). Optical transmissions (10 mm path length) at various wavelengths selected with the monochromator (bandwidths 2-20 nm) were measured as a function of time before and after the pulse using photoelectric detection. The output of the photomultiplier (Hamamatsu R928) was displayed on a Tektronix TDS 380 digitizing oscilloscope. Data processing was performed on a Dan PC using software developed in house. The sample cell was constructed from Spectrosil quartz and formed part of a flow-through system, introducing fresh solution prior to each laser shot [14, 15]. Benzophenone in acetonitrile was used as a reference.

The luminescence (1270 nm) from excited <sup>1</sup>O<sub>2</sub> (<sup>1</sup>Δ<sub>g</sub>) produced by photoexcitation of primaquine in oxygenated solution was detected by a liquid N<sub>2</sub>-cooled Germanium Detector / Amplifier (Applied Detector Corporation 403HS) closely coupled to the laser photolysis cell in right-angle geometry [16, 17]. A 1mm thick (20 mm diameter) piece of AR-coated silicon

(II-IV Inc.) was placed between the diode and cell to act as a cut-off filter for light  $< 1100$  nm. The 403HS power supply bias voltage was operated at  $-450$  V. The amplifier output was AC-coupled to the digitizer.

### 2.3. *Pulse radiolysis*

The pulse radiolysis experiments were carried out with a 12 MeV RDL, 3GHz electron linear accelerator with a pulse duration from  $0.22 - 2$   $\mu$ s and with a peak current of about 30 mA. It is normally operated at 10 pulses per second but the single pulse mode is achieved by modifying the pulses to the gun [18]. The detection system, similar in principle to the laser flash photolysis apparatus, employed a Kratos monochromator, EMI 9558Q photomultiplier and Tektronix TDS 754C digitizing oscilloscope. The sample cell (Spectrosil quartz) had an optical path length of 25 mm [19]. Radiation doses were measured using the thiocyanate actinometer where the absorption of  $(\text{SCN})_2^{\bullet-}$  formed by pulsing an air saturated aqueous solution of 10 mM KSCN is measured taking  $\epsilon_{500} (\text{SCN})_2^{\bullet-} = 7.1 \cdot 10^3 \text{ M}^{-1} \text{ cm}^{-1}$  and  $G(\text{SCN})_2^{\bullet-} = 2.9 \text{ heV}^{-1}$  [20].

### 2.4. *Photochemical degradation*

Photodegradation products from primaquine diphosphate ( $40 \mu\text{M}$  in  $\text{D}_2\text{O}$ ) were formed by irradiation for 5 min. or 15 min. in the Suntest CPS sun simulating exposure unit (1.8 kW Xe-burner; Heraeus GmbH, Hanau, Germany), equipped with a quartz-glass dish with IR reflective coating and a glass filter (transmission 310-800 nm). The energy output was  $765 \text{ W/m}^2$ , quantified by the XenoCal Sensor equipped with XenoSoft Software (Atlas, Germany) calibrated against filtered radiation (300-800 nm) of a Xe discharge lamp. The samples were irradiated in quartz cells (optical path length 10 mm) during continuous stirring. The degradation was followed by HPLC and UV-vis spectrophotometry. The degradation of primaquine ( $40 \mu\text{M}$ ) in  $\text{D}_2\text{O}$  was compared to the analogue degradation in water, and the influence of molecular oxygen was investigated by flushing with  $\text{N}_2$ -gas for 10 min. prior to irradiation.

Photosensitized degradation of primaquine in  $\text{D}_2\text{O}$  was studied using a 900 W Xe arc lamp fitted with a Monochromator  $f$  3.4 (Applied Photophysics Ltd., Surrey, England). The samples were irradiated at  $455 \pm 20$  nm up to 60 min. in quartz sample cells (optical path length 10 mm) under continuous stirring at ambient temperature. The adjusted energy at the surface of the container was 72 mV, quantified by a Thermopile voltmeter (200 mV, Applied Photophysics Ltd.) calibrated against a blackbody radiator. The samples contained non-irradiated primaquine diphosphate ( $40 \mu\text{M}$ ) or primaquine diphosphate ( $40 \mu\text{M}$ ) irradiated in the Suntest for 5 min. or 15 min. as described above. The samples that were irradiated for 15 min. were added to extra primaquine diphosphate ( $40 \mu\text{M}$ ) prior to monochromatic irradiation due to extensive decomposition in the Suntest. Additionally, sodium azide ( $\text{NaN}_3$  10 mM) was added as a  $^1\text{O}_2$  quencher prior to monochromatic irradiation in samples that were pre-irradiated for 5 min. Primaquine diphosphate ( $20 \mu\text{M}$  in  $\text{D}_2\text{O}$ ) was irradiated at  $350 \pm 20$  nm (72 mV) for 0-60 min. as a control of the sensitized reaction. Degradation of primaquine was quantified by use of HPLC and followed by UV-vis spectrophotometry.

### 2.5. *Quantification by HPLC*

The concentration of primaquine was quantified by reverse phase HPLC, as previously described [4] (System 2).

## 2.6. UV-visible absorption spectra

UV-visible absorption spectra (190-700 nm) were recorded by a Shimadzu UV-2401PC UV-vis recording spectrophotometer. The pH of aqueous solutions was adjusted when appropriate by addition of NaOH or HClO<sub>4</sub>, and controlled by use of a pH-electrode.

## 2.7. Fluorescence measurements

Fluorescence spectra were measured on a PTI modular Fluorescence System using Felix™ for Windows software. The excitation source was a 75 W xenon lamp. The monochromators were Model 101 (f/4 0.2 m Czerny-Turner configuration).

Fluorescence lifetimes (Ex. 337 nm; Em. 527 nm) of primaquine diphosphate in water (Abs<sub>337</sub> = 0.31), primaquine diphosphate in D<sub>2</sub>O (Abs<sub>337</sub> = 0.39) and primaquine diphosphate in D<sub>2</sub>O (40 μM) irradiated in the Suntest for 20 min. (resulting in Abs<sub>337</sub> = 0.08) were measured. Riboflavin in purified water (Abs<sub>337</sub> < 0.1) was used as reference (Ex. 337 nm; Em. 538 nm).

Fluorescence quantum yields (Φ<sub>f</sub>) of primaquine diphosphate in water and in D<sub>2</sub>O (Ex. 350 nm, Em. 370-700 nm; Abs<sub>350</sub> ≤ 0.115) and of primaquine diphosphate in D<sub>2</sub>O (40 μM) irradiated in the Suntest for 20 min. (Ex. 455 nm, Em. 470-700 nm, Abs<sub>455</sub> = 0.086) were calculated. Fluorescence quantum yields were obtained by using quinine sulphate dissolved in 0.05 M H<sub>2</sub>SO<sub>4</sub> as reference (Φ<sub>f</sub> = 0.55) [21].

The formation of <sup>1</sup>O<sub>2</sub> sensitized by primaquine diphosphate in D<sub>2</sub>O (Ex. 455 nm) and photoproducts of primaquine diphosphate in D<sub>2</sub>O (40 μM) irradiated in the Suntest for 20 min. was investigated by direct detection of <sup>1</sup>O<sub>2</sub> luminescence at 1270 nm in a steady state mode. The detector was an EQ-817 Germanian Detector System operated under liquid N<sub>2</sub> operations. Rose Bengal in D<sub>2</sub>O was used as reference (Ex. 548 nm).

## 3. Results and discussion

Primaquine is an aminoquinoline with pK<sub>a</sub> values = 3.2 (quinoline N) and 10.4 (amine group) [22]. The pK<sub>a</sub> value of the aniline N is very low (< -1), likely due to delocalization of the lone-pair electrons of this group into the aromatic ring. At physiological pH primaquine exists as the monocation PQH<sup>+</sup> (Scheme 1). A summary of the photophysical properties and reaction rates of primaquine transients are presented in Tab.1.

### 3.1. Studies of the excited state of primaquine by laser flash photolysis and pulse radiolysis

Following 355 nm laser flash photolysis of a nitrogen saturated aqueous solution of primaquine (0.1 mM), transient absorption bands centred at 410 and 720 nm were observed at the end of pulse, the 720 nm band decayed with a first order rate of 2•10<sup>6</sup> s<sup>-1</sup> while that at 410 nm was found to be long lived and decayed by a second order process (Fig. 1). Moreover, the long wavelength absorption band was quenched within the pulse in the presence of nitrous oxide. On the basis of the absorption spectrum and the reaction with nitrous oxide, the long wavelength absorption band is assigned to the hydrated electron formed presumably by reaction (1):



On saturating the solution with oxygen, it was found that the long wavelength band rapidly decayed ( $\tau < 0.2 \mu\text{s}$ ) while that centred at 410 nm showed only a very slight increase in the decay rate. On the basis of the above we have assigned this band (410 nm) to the cation radical of primaquine. This assignment was confirmed by pulse radiolysis (see section 3.2) and is in agreement with the previous work by Viola et al. [23]. Figure 1B shows the difference absorption spectrum of the long lived species in air saturated aqueous solution of primaquine. The difference between the 5  $\mu\text{s}$  spectrum (Fig. 1A) and 1B (in 370-600 nm wavelength region) is explicable in terms of the absorption due to the anion radical of primaquine formed by the reaction of the photoejected hydrated electron with primaquine in 1A where as the electron is quenched by oxygen in 1B. Primaquine anion radical is shown to have a broad absorption band with maximum at 490 nm [24]. Furthermore, this shows that the superoxide anion radical formed by the reaction of hydrated electron with oxygen does not reduce primaquine as will be shown in section 3.3 by pulse radiolysis.

In order to determine whether the photoionisation is monophotonic or not, we determined the effect of laser intensity on the transient formation. It was found that both 720 and 410 nm absorption increased non-linearly with the laser intensity (see inset in Fig. 1B), indicating that the photoionisation is not a monophotonic process. From the log plot (inset (b) Fig. 1B) we obtained a value of 1.6 instead of 2 due to a biphotonic process. This may indicate a mixed mono and bi- photonic process; we have failed to observe significant transient absorbance at either 700 or 410 nm under experimental conditions that otherwise produce normal one-photon excitation in past experiments. Stevenson et al. [25] note that the observed two photon power dependence is less with an experimental setup in which the laser and analyzing light beams are perpendicular (as in our experiment) than in an arrangement with collinear beams. In addition it is noted that at the powers and concentrations used in our experiments, saturation of absorption is likely. These factors may explain why we observe power dependencies of less than 2 in experiments such as those illustrated in Figure 1. From the above experiments no absorption due to the triplet state could be detected in the UV/ vis region. It may be that the triplet state is too short lived and/or its absorption coefficient is too small for our detection system to detect. Various media were used for a thorough investigation but no transient attributable to the primaquine triplet was detected.

The luminescence at 1270 nm from excited  $^1\text{O}_2$  ( $^1\Delta_g$ ) sensitized by photoexcitation (355 nm) of primaquine ( $^{\text{PQ}}\text{Abs}_{355} = 0.25$ ) in  $\text{O}_2$ -saturated toluene was then measured as an indirect detection of the primaquine triplet (Fig. 2). Perinaphthenone in air-saturated toluene ( $^{\text{PN}}\text{Abs}_{355} = 0.25$ ) was used as reference. For any given laser energy, the number of photons absorbed by the two solutions will be the same since both have the same absorbance at the wavelength of excitation (355 nm). The quantum yield of  $^1\text{O}_2$  formation sensitized by primaquine ( $^{\text{PQ}}\Phi_{\Delta}$ ) is 0.025, as calculated from

$$^{\text{PQ}}\Phi_{\Delta} / ^{\text{PN}}\Phi_{\Delta} = ^{\text{PQ}}I_e / ^{\text{PN}}I_e \quad \text{Eq. 1}$$

where  $^{\text{PN}}\Phi_{\Delta}$  is taken as 0.98 [12];  $^{\text{PQ}}I_e$  and  $^{\text{PN}}I_e$  are initial emission intensities for primaquine and perinaphthenone, respectively, at any given laser intensity and are obtained by the slopes of the lines in Fig. 2 ( $^{\text{PQ}}I_e = 1.06$ ,  $R^2 = 0.977$ ;  $^{\text{PN}}I_e = 42.25$ ,  $R^2 = 0.999$ ). This is in agreement with a value of 0.01 in air saturated benzene [26]. However, Motten et al. [27] did not detect any significant  $^1\text{O}_2$  formation induced by primaquine in  $\text{D}_2\text{O}$  at pD 7 ( $^{\text{PQ}}\Phi_{\Delta} < 0.005$ ).



At pulse energy of 9.50 mJ the initial intensity of the emission is reduced from 10.2 mV to 5.4 mV by shaking the O<sub>2</sub>-saturated sample of primaquine in toluene with air prior to excitation. Moreover, <sup>PQ</sup>I<sub>e</sub> is doubled (factor 2.2) on increasing the primaquine concentration from 9.1•10<sup>-5</sup> M to 6.4•10<sup>-4</sup> M. These observations verify the formation of <sup>1</sup>O<sub>2</sub> during photoexcitation of primaquine. It seems likely that <sup>1</sup>O<sub>2</sub> is formed by energy transfer from the short lived primaquine triplet (<sup>3</sup>PQ\*) in accordance with reaction (2):



Sensitization experiments were performed to further confirm the existence of the primaquine triplet. Energy transfer was studied by laser flash photolysis at 355 nm of primaquine (0.3 mM) in N<sub>2</sub>-saturated acetonitrile in the presence of fenbufen (6 mM). Fenbufen in acetonitrile shows transient absorption centred at 420 nm which is assigned to its triplet state [13]. The lifetime of the fenbufen triplet is 10±0.5 μs, calculated by the first order decay of the transient at 420 nm (k<sub>ref</sub> = 9.9±0.4•10<sup>4</sup> s<sup>-1</sup>, n = 2). Primaquine reduces this lifetime by about 90%. But no absorption due to the primaquine triplet could be detected. On reducing the concentration of primaquine to 0.1 and 0.2 mM, the fenbufen triplet was quenched at the rate of k<sub>obs</sub> = 4•10<sup>5</sup> and 7•10<sup>5</sup> s<sup>-1</sup> respectively. From these data the bimolecular rate constant for quenching of fenbufen triplet by primaquine was calculated to be k<sub>q</sub> = 3•10<sup>9</sup> M<sup>-1</sup> s<sup>-1</sup>. The quenching rate constant is calculated from

$$k_q (\text{M}^{-1}\text{s}^{-1}) = k_{\text{obs}} (\text{s}^{-1}) - k_{\text{ref}} (\text{s}^{-1}) / [\text{primaquine}] (\text{M}) \quad \text{Eq. 2}$$

Thus primaquine is an efficient quencher of the fenbufen triplet. It seems likely that, as the fenbufen triplet decays, the primaquine triplet is formed in accordance with reaction (3):



Sensitization of primaquine (93 μM) in N<sub>2</sub>-saturated toluene was further investigated by pulse radiolysis in the presence of biphenyl (10 mM). The biphenyl triplet is formed by pulse radiolysis during energy transfer from excited toluene [14]. The first order decay of the biphenyl triplet in toluene at 360 nm (k<sub>ref</sub> = 2.2•10<sup>4</sup> s<sup>-1</sup>; n = 4; RSD = 20 %) is increased in the presence of primaquine (k<sub>obs</sub> = 3.8•10<sup>5</sup> s<sup>-1</sup>; n = 3; RSD = 10 %). Primaquine is an efficient quencher of the biphenyl triplet with a bimolecular quenching rate constant k<sub>q</sub> = 3.9•10<sup>9</sup> M<sup>-1</sup>s<sup>-1</sup>, as calculated by Eq. 2.

As the biphenyl triplet decays, it seems likely that the primaquine triplet is formed in accordance with reaction (4):



Existence of a short lived primaquine triplet (< 50 ns) is likely, based on the sensitization experiments, formation of singlet oxygen and on the effect of oxygen concentration on the yield of singlet oxygen. Failure to observe the triplet-triplet absorption spectrum of primaquine under our conditions may be due to either the short lifetime and/or very small molar absorption coefficient in the wavelength region analysed.

### 3.2. Identification and characterizations of the cation radicals of primaquine by pulse radiolysis

As seen in the previous section primaquine is photoionised in a biphotonic process to form the cation radical. We have used the techniques of pulse radiolysis to determine and characterise the cation radical.

The reaction of the hydroxyl radical OH• with monoprotinated primaquine (PQH<sup>+</sup>) was studied by pulse radiolysis of an N<sub>2</sub>O-saturated solution of primaquine (0.2 mM) in 10 mM phosphate buffer pH 7.4. Under these conditions the initially produced hydrated electrons are scavenged by N<sub>2</sub>O, leaving only OH• as the reactive radical [14]. Fig. 3 shows the transient difference absorption spectra observed 7 μs and 191 μs after the pulse. Under these conditions a broad band centred at 400 nm and a very broad band in the visible region are observed at the end of OH• reaction (7 μs). However at 191 μs the band centred at 400 nm narrows and shifts to 410 nm. OH• radicals are known to react with organic compounds by addition to double bonds, hydrogen abstraction, and electron abstraction by formation of cation radicals as in the case of the antimalarial mefloquine [17]. The bimolecular quenching rate constant of OH• by primaquine  $k_q = 6.6 \cdot 10^9 \text{ M}^{-1} \text{ s}^{-1}$ , calculated as the slope of the line ( $R^2 = 0.957$ ) when the pseudo first order rate for the growth of transient absorption at 410 nm was plotted as a function of primaquine concentration at 5.0-10.0 μM ( $n \leq 3$ ). The rate of reaction is in the order of magnitude expected for the addition of OH• radicals to, or abstraction of an electron / H-atom from, aromatic compounds [28]. Hence, the spectrum shown in Fig.3 could be either the primaquine monoprotinated cation radical and/or other products, such as an OH• - addition product, as illustrated in reaction (5):



The formation of PQH<sup>2+•</sup> was confirmed and the monoprotinated cation radical further characterized by pulse radiolysis of primaquine (0.2 mM) in N<sub>2</sub>O-saturated 10 mM phosphate buffer pH 7.4 containing 10 mM KBr. The one electron oxidizing radical Br<sub>2</sub>•<sup>-</sup> is produced when N<sub>2</sub>O-saturated aqueous solutions of KBr are subject to pulse radiolysis [14]. Fig. 4 shows the transient difference absorption spectra observed at 5 μs and 51 μs after the pulse. The transient absorption maxima at 410 nm and 560 nm are due to formation of PQH<sup>2+•</sup> during oxidation of primaquine by the Br<sub>2</sub>•<sup>-</sup> radicals at neutral pH, in accordance with reaction (6):



Both 410 and 560nm absorption decayed by a second order process with  $k_2 = 7.4 \pm 0.6 \cdot 10^8 \text{ M}^{-1} \text{ s}^{-1}$ , hence both bands are due to the same species, PQH<sup>2+•</sup>. It is clear, from a comparison of the spectra in Figs. 3 and 4, that less than half of OH• reaction with primaquine results in the formation of PQH<sup>2+•</sup>.

Br<sub>2</sub>•<sup>-</sup> shows characteristic transient absorption at 360 nm. By plotting the pseudo first order rate for the decay of transient absorption at this wavelength as a function of primaquine concentration at zero,  $1.0 \cdot 10^{-5} \text{ M}$ ,  $2.5 \cdot 10^{-5} \text{ M}$  and  $5.0 \cdot 10^{-5} \text{ M}$  ( $n \leq 3$ ), the bimolecular quenching rate constant was calculated as the slope of the line ( $R^2 = 0.998$ );  $k_q = 4.7 \cdot 10^9 \text{ M}^{-1} \text{ s}^{-1}$ .

Primaquine forms oxygenated photoproducts in a complex degradation pattern which is highly dependent on the oxygen content of the medium [2, 4]. Thus the lifetime of PQH<sup>2+•</sup> as a function of oxygen concentration was studied by pulse radiolysis of primaquine (0.2 mM) in 10 mM phosphate buffer pH 7.4 containing 10 mM KBr. Since PQH<sup>2+•</sup> decayed essentially by

a second order process, the dose was kept as low as possible (~ 2.5 Gy) and adjusted to ensure the initial concentration of  $\text{PQH}^{2+\bullet}$  remained the same for this set of experiments. Samples were saturated with  $\text{O}_2$  or  $\text{N}_2\text{O}$ , and mixed to concentration ranges of  $\text{O}_2:\text{N}_2\text{O} \sim 0:100, 20:80$  and  $75:25$ . The lifetime of  $\text{PQH}^{2+\bullet}$  was calculated by the pseudo first order decay of the transient absorption at 410 nm ( $n \leq 4$ ;  $R^2 = 0.980$ ) as a function of per cent oxygen content (Fig. 5).  $\text{O}_2$  will influence degradation of the drug by direct reactions with  $\text{PQH}^{2+\bullet}$  according to reaction (7):



where PQox is oxidized degradation product(s).

The bimolecular quenching rate constant of  $\text{PQH}^{2+\bullet}$  by  $\text{O}_2$ ,  $k_q = 1.0 \cdot 10^6 \text{ M}^{-1}\text{s}^{-1}$ , was calculated from the slope of the plot of the pseudo first order rate of the transient decay at 410 nm ( $\text{PQH}^{2+\bullet}$ ) vs. the molar oxygen concentration ( $n \leq 4$ ;  $R^2 = 0.962$ ). The oxygen content in water under pure oxygen at 25 °C was taken as  $1.29 \cdot 10^{-3} \text{ M}$ , according to Connors et al. [29].

The reduction potential of primaquine under physiological conditions was further evaluated by pulse radiolysis of primaquine (0.5 mM) in  $\text{N}_2\text{O}$ -saturated 10 mM phosphate buffer pH 7.4 containing 5 mM tryptophan (TRP) and 0.1 M KBr. Fig. 6 shows the transient difference absorption spectra observed at 7  $\mu\text{s}$ , 27  $\mu\text{s}$  and 69  $\mu\text{s}$  after the pulse. The maximum at 520 nm is due to formation of the deprotonated cation radical of tryptophan by reaction (8a & 8b) in the presence of  $\text{Br}_2\bullet^-$ :



The corresponding decays of  $\text{TRP}\bullet$  (520 nm) and increase in  $\text{PQH}^{2+\bullet}$  (410 nm) shows that primaquine is a quencher of  $\text{TRP}\bullet$  according to reaction (9):



The bimolecular quenching rate constant of  $\text{TRP}\bullet$  by  $\text{PQH}^+$ ,  $k_q = 2.5 \pm 0.3 \cdot 10^7 \text{ M}^{-1}\text{s}^{-1}$ , was determined from the pseudo first order rate constant for the decay of  $\text{TRP}\bullet$  absorption at 520 nm in the absence of primaquine ( $k_{\text{ref}} = 1.6 \pm 0.6 \cdot 10^3 \text{ s}^{-1}$ ,  $n = 4$ ) and in the presence of 0.5 mM primaquine ( $k_{\text{obs}} = 1.4 \pm 0.1 \cdot 10^4 \text{ s}^{-1}$ ,  $n = 2$ ), and calculated according to Eq. 2. The pseudo first order rate constant for the decay of  $\text{TRP}\bullet$  when quenched by  $\text{PQH}^+$  ( $k_{\text{obs}} = 1.4 \pm 0.1 \cdot 10^4 \text{ s}^{-1}$ ,  $n = 2$ ) as measured at 520 nm corresponds well with the rate constant for the increase of  $\text{PQH}^{2+\bullet}$  measured at 410 nm ( $k_1 = 1.6 \cdot 10^4 \text{ s}^{-1}$ ,  $n = 1$ ). This is expected according to reaction (9), and verifies that primaquine ( $\text{PQH}^+$ ) acts as a quencher of  $\text{TRP}\bullet$ . The reduction potential  $E^\circ$  ( $\text{PQH}^{2+\bullet} / \text{PQH}^+$ ) at physiological pH is thus lower than +1015 mV, which is the value for  $E^\circ$  ( $\text{TRP}\bullet / \text{TRPH}$ ) at neutral pH [30].

The reduction potential of  $\text{PQH}^{2+\bullet} / \text{PQH}^+$  was also evaluated by pulse radiolysis of primaquine (0.5 mM) in  $\text{N}_2\text{O}$ -saturated 10 mM phosphate buffer pH 7.4 containing 2 mM tyrosine (TYR) and 0.1 M KBr. Tyrosine will not form radicals in reactions with  $\text{Br}_2\bullet^-$  at these concentrations due to a much faster reaction rate between  $\text{PQH}^+$  and  $\text{Br}_2\bullet^-$ . The apparent first order decay of  $\text{PQH}^{2+\bullet}$  absorption detected at 410 nm ( $k_1 = 3.0 \pm 0.7 \cdot 10^3 \text{ s}^{-1}$ ,  $n = 3$ ) is not changed in the presence of tyrosine ( $k_1 = 3.1 \pm 0.7 \cdot 10^3 \text{ s}^{-1}$ ,  $n = 2$ ). The tyrosine radical, detected

in a separate experiment at 410 nm in the absence of primaquine, has a faster decay ( $k_1 = 1.8 \cdot 10^4 \text{ s}^{-1}$ ,  $n = 1$ ). The reduction potential  $E^\circ$  ( $\text{PQH}^{2+\bullet} / \text{PQH}^+$ ) is thus likely to be even lower than +930 mV, which is the reduction potential  $E^\circ$  ( $\text{TYR}^\bullet / \text{TYR}$ ) at neutral pH [30]. However, it can not be excluded that the quenching of  $\text{PQH}^{2+\bullet}$  by tyrosine is not observed due to a low reaction rate.

The long-lived nature of the 390-400 nm / 550-560 nm absorbing species formed in the reaction between  $\text{PQH}^+$  and  $\text{OH}^\bullet / \text{Br}_2^{\bullet-}$ , its relatively slow reactivity with oxygen ( $k_q = 1.0 \cdot 10^6 \text{ M}^{-1}\text{s}^{-1}$ ) and the fact that it is formed from the one electron oxidizing radical  $\text{Br}_2^{\bullet-}$  leads to the conclusion that this is the monoprotonated cation radical of primaquine ( $\text{PQH}^{2+\bullet}$ ). As  $\text{PQH}^{2+\bullet}$  is formed photochemically by a two-photon process, direct photoionisation by solar or room light exposure is not likely. However, the radical cation might be formed by photosensitized reactions of types I or II, as illustrated in Scheme 2. The relatively low reduction potential of primaquine at physiological pH ( $E^\circ$  ( $\text{PQH}^{2+\bullet} / \text{PQH}^+$ )  $< +1015 \text{ mV}$ ) combined with the long lifetime makes primaquine subject to oxidative degradation by formation of the monoprotonated cation radical. Reaction (7) is previously suggested as an important initial process in the photochemical degradation of primaquine [4]. Oxidation of the monoprotonated cation radical by molecular oxygen will make an important contribution to degradation under aerobic conditions. Previous studies also indicate that  $\text{OH}^\bullet$  is formed during photochemical degradation of primaquine (e.g. by degradation products). Formation of the  $\text{OH}^\bullet$  radical will increase the decomposition of primaquine in aqueous solution, as primaquine is oxidized by  $\text{OH}^\bullet$  at a high rate ( $k_q = 6.6 \cdot 10^9 \text{ M}^{-1}\text{s}^{-1}$ ).

Physicochemical properties and photoreactivity of primaquine is highly influenced by the protonation/deprotonation of the drug (Scheme 1). Hence, the formation of cation radicals of primaquine at various degree of protonation was studied by pulse radiolysis of primaquine (0.5 mM) in  $\text{N}_2\text{O}$ -saturated aqueous solutions containing 10 mM KBr at pH 2.3 ( $\text{PQH}_2^{2+}$ ), pH 5.6 ( $\text{PQH}^+$ ) and pH 10.1 (PQ). Fig. 7 shows the corrected (for ground state bleaching) transient absorption spectra that were observed immediately after the pulse (6  $\mu\text{s}$  at pH 2.3; 3  $\mu\text{s}$  at pH 5.6 and 8  $\mu\text{s}$  at pH 10.1, respectively). The transients at pH 2.3 and 5.6 have absorption maxima at 400-410 nm and 550-570 nm, i.e. identical with the spectrum detected at pH 7.4 (Fig. 4). The transient at pH 10.1 shows a different absorption spectrum, with a red shift of the first absorption maximum to 650 nm, a new (small) maximum at 470 nm and a blue shift to 385 nm (Fig. 7). The primaquine cation radical is most likely centred at the quinoline nitrogen. The proton that is attached to this functional group at acidic pH will thus be repulsed during the oxidation reaction with the  $\text{Br}_2^{\bullet-}$  radicals. The cation radicals formed at pH 2.3; 5.6 and pH 7.4 will then be identical, which is verified by equal second order rate constants for the transient decay ( $k_2 = 6.0\text{-}7.4 \cdot 10^8 \text{ M}^{-1}\text{s}^{-1}$ ) and illustrated in reactions (6) and (10). The transient absorption spectrum shows a different profile at alkaline pH, due to the shorter lived ( $k_2 = 1.0 \cdot 10^9 \text{ M}^{-1}\text{s}^{-1}$ ) non-protonated radical that is formed in accordance with reaction (11):



The reactions are illustrated in Scheme 1. The intramolecular repulsion between charged groups in  $\text{PQH}^{2+\bullet}$  will favour intermolecular interactions with the medium, while an intramolecular interaction between the quinoline radical centre and amine group in  $\text{PQ}^{\bullet+}$  is likely to shield part of the transient from the surroundings. This can explain the differences in

the transient absorption spectra as well as the longer lifetime of  $\text{PQH}^{2+\bullet}$  compared with that of  $\text{PQ}^{\bullet+}$ . It is interesting to compare these observations with absorption properties in the ground state, which are similar at alkaline and neutral pH (maximum at 350 nm, spectrum extended until 430 nm). Acidification and formation of the ground state primaquine dication leads to formation of two absorption maxima at 330 and 420 nm, respectively, and extension of the spectrum towards 500 nm. The first absorption band of primaquine in the ground state represents the transition of the lone-pair electrons ( $n\text{-}\pi^*$  transition) at the quinoline nitrogen. Intramolecular bonding is favoured in both alkaline and neutral media, leading to similar absorption properties. Intramolecular charge repulsion and extended interaction with the medium change the absorption properties at acidic pH [5].

The bimolecular quenching rate constants of  $\text{Br}_2^{\bullet-}$  by  $\text{PQH}_2^{2+}$  (pH 2.4) and PQ (pH 10.8) are  $k_q = 2.7 \pm 0.2 \cdot 10^9 \text{ M}^{-1} \text{ s}^{-1}$  and  $k_q = 4.2 \pm 0.1 \cdot 10^9 \text{ M}^{-1} \text{ s}^{-1}$ , respectively, as determined by the pseudo first order rate for the decay of transient absorption at 360 nm ( $\text{Br}_2^{\bullet-}$ ) detected in an  $\text{N}_2\text{O}$ -saturated aqueous medium containing 10 mM KBr ( $k_{\text{ref}} = 4.5 \pm 0.2 \cdot 10^3 \text{ s}^{-1}$  at pH 2.4,  $n = 3$ ;  $k_{\text{ref}} = 4.6 \pm 0.4 \cdot 10^3 \text{ s}^{-1}$  at pH 10.8,  $n = 2$ ) and in the presence of 10  $\mu\text{M}$  primaquine ( $k_{\text{obs}} = 3.1 \pm 0.1 \cdot 10^4 \text{ s}^{-1}$  at pH 2.4,  $n = 2$ ;  $k_{\text{obs}} = 4.7 \pm 0.05 \cdot 10^4 \text{ s}^{-1}$  at pH 10.8,  $n = 3$ ). The quenching rates are calculated according to Eq. 2. The quenching rate constant at pH 7.4 ( $\text{PQH}^+$ ) is previously determined to be  $k_q = 4.7 \cdot 10^9 \text{ M}^{-1} \text{ s}^{-1}$ . Primaquine as a monocation (neutral pH) and non-ion (alkaline conditions) thus undergo one electron oxidization by  $\text{Br}_2^{\bullet-}$  radical at diffusion controlled rates, while the dication (acidic medium) is oxidized at about half the rate. Protonation of the quinoline N and binding of the reactive n-electrons at low pH can explain the reduced reaction rate of the dication compared to the two other forms (Scheme 1).

The molar absorption coefficient ( $\epsilon$ ) of primaquine transients as a function of pH were calculated from the transient absorption maximum ( $A_{\text{max}}$ ) detected immediately after the laser pulse (3-12  $\mu\text{s}$ ), the optical path length (2.5 cm) and estimated concentration of transient assuming the yield of  $e_{\text{aq}}^-$  to be 2.7  $\mu\text{M kRad}^{-1}$  and 100% efficiency of the radical reactions:

$$\epsilon (\text{M}^{-1} \text{ cm}^{-1}) = A_{\text{max}} / [\text{PQ-transient}] \text{ M} \cdot 2.5 \text{ cm} \quad \text{Eq. 3}$$

The monoprotonated cation radical ( $\text{PQH}^{2+\bullet}$ ) has the same molar absorption coefficient in aqueous media at pH 5.6 and 7.4 ( $\epsilon_{400} = 7300 \text{ M}^{-1} \text{ cm}^{-1}$ ;  $\epsilon_{550} = 1700\text{-}1800 \text{ M}^{-1} \text{ cm}^{-1}$ ). In an acidified medium (pH 2.3) the calculated molar absorption coefficients are only slightly changed ( $\epsilon_{410} = 6500 \text{ M}^{-1} \text{ cm}^{-1}$ ;  $\epsilon_{570} = 2000 \text{ M}^{-1} \text{ cm}^{-1}$ ).

The  $\text{pK}_a$ -value of the cation radicals ( $\text{PQH}^{2+\bullet} / \text{PQ}^{\bullet+}$ ) was evaluated by pulse radiolysis of primaquine (10  $\mu\text{M}$ ) in  $\text{N}_2\text{O}$ -saturated aqueous solutions containing 10 mM KBr at pH 2.4; 7.1; 8.1; 9.5; 10.8 and 11.5. The plot of transient absorbance at 560 nm ( $\text{PQH}^{2+\bullet}$ ) as a function of pH is presented in Fig. 8. The steep decline in absorbance between pH 7 and 8 indicates that  $\text{PQH}^{2+\bullet} / \text{PQ}^{\bullet+}$  has a  $\text{pK}_a$ -value in this range, which is a significant change from the ground state ( $\text{pK}_a = 10.4$ ). The corresponding increase in transient absorption at 640 nm ( $\text{PQ}^{\bullet+}$ ) with an increase in pH did not demonstrate any characteristic behaviour that could be used in the calculations.

### 3.3. Studies of the scavenging reaction between primaquine and superoxide by pulse radiolysis

In order to study the reduction of primaquine by superoxide anions at physiological pH, the radical anion of primaquine (protonated / deprotonated) was identified by pulse radiolysis of


primaquine (0.2 mM) in N<sub>2</sub>-saturated 10 mM phosphate buffer pH 7.4 containing 1% t-butanol. The initially produced OH• and H• radicals react rapidly with t-butanol producing relatively unreactive t-butanol radicals [14]. The hydrated electron (e<sub>aq</sub><sup>-</sup>) is unaffected by the presence of t-butanol and is free to react with primaquine according to reaction (12):




According to Bisby [24] the unprotonated 8-aminoquinoline ring of primaquine appears to be the site for reduction by e<sub>aq</sub><sup>-</sup> at a diffusion controlled rate (k<sub>q</sub> = 2.47±0.1•10<sup>10</sup> M<sup>-1</sup>s<sup>-1</sup> at pH 7.3). The anion radical is supposed to be deprotonated (PQ•<sup>-</sup>) at neutral pH based on measured electron transfer reaction rates. The strongly reducing primaquine anion radical (E°<sub>PQH<sup>+</sup>/PQ•<sup>-</sup></sub> < -1022 mV) is efficiently quenched by molecular oxygen (k<sub>q</sub> = 2.31±0.25•10<sup>9</sup> M<sup>-1</sup>s<sup>-1</sup> at pH 6.9), likely by one-electron transfer by formation of superoxide anions (E°<sub>O<sub>2</sub>/O<sub>2</sub>•<sup>-</sup></sub> = -160 mV). According to these observations, the reverse reaction is not very likely.

The molar absorption coefficient of the anion radical was calculated by Eq. 3 at the absorption maximum at 350 nm (ε = 4700 M<sup>-1</sup> cm<sup>-1</sup>) and 490 nm (ε = 2100 M<sup>-1</sup> cm<sup>-1</sup>) (spectrum not shown). The hydrated electron has an absorption maximum at 720 nm and will not interfere with the spectrum [14]. The lifetime of PQH• / PQ•<sup>-</sup> in an anaerobic medium is in the milliseconds.

One-electron transfer from the superoxide radical anion (O<sub>2</sub>•<sup>-</sup>) to primaquine at physiological pH (PQH<sup>+</sup>) was subsequently studied by pulse radiolysis of primaquine (0.1 mM) in O<sub>2</sub>-saturated 10 mM phosphate buffer pH 7.4 containing 0.1 M sodium formate to increase the yield of O<sub>2</sub>•<sup>-</sup>. Pulse radiolysis of the sample produces superoxide radicals by reactions (13-15) [14]:



As expected, no reaction between primaquine and O<sub>2</sub>•<sup>-</sup> could be detected, as investigated at the absorption maximum of the primaquine anion radical (PQH•/ PQ•<sup>-</sup>) at 480 nm, illustrated by reaction (16):



The result is verified by the fact that no bleaching (detected as negative absorption) is observed by primaquine in the ground state at 360 nm (data not shown). These findings are in accordance with previous scavenging studies, where superoxide dismutase did not reduce photodegradation of primaquine although O<sub>2</sub>•<sup>-</sup> is formed in the reactions [3, 4]. Hence, primaquine will not be decomposed by self-sensitization by O<sub>2</sub>•<sup>-</sup> in the photochemical degradation process.

### 3.4. Photosensitized degradation of primaquine

The photochemical degradation of primaquine in the Suntest is significantly accelerated in deuterium oxide (D<sub>2</sub>O) compared to purified water, as 10(±3) % (n = 2) was decomposed

after only 5 min. irradiation in D<sub>2</sub>O while the drug had to be irradiated for 60 min. in water to obtain 13(±2) % (n = 2) decomposition. Reduction of the oxygen content by flushing with N<sub>2</sub>-gas prior to irradiation in D<sub>2</sub>O reduces the degradation from 86(±1) % (n = 2) to 45(±10) % (n = 3) after 15 min. exposure. The influence of oxygen concentration and the increased degradation in D<sub>2</sub>O indicate that <sup>1</sup>O<sub>2</sub> is involved in the reaction, which is likely since primaquine generates some <sup>1</sup>O<sub>2</sub> during laser flash photolysis (<sup>P</sup>Φ<sub>Δ</sub> = 0.025) and the drug is a quencher of <sup>1</sup>O<sub>2</sub> at moderate rate (k<sub>q</sub> = 2.6•10<sup>8</sup> M<sup>-1</sup> s<sup>-1</sup>) [27]. The first absorption band of ground state primaquine at 350 nm is decreased during photodegradation parallel to growth of a new maximum at 455 nm (i.e. outside the original absorption range). Irradiation at 455 nm was thus selected for investigation of a photosensitized degradation. Based on quenching studies we have previously suggested that initially formed photodegradation products or impurities of the raw material are sources of <sup>1</sup>O<sub>2</sub> in the photochemical reactions [4].

The photosensitized degradation of primaquine by monochromatic irradiation at 455 nm is illustrated in Fig. 9. The reactions can be expressed by apparent zero order degradation kinetics and the corresponding degradation rates (k<sub>obs</sub>) are given as relative values obtained under the selected experimental conditions. The results emphasize the possibility that one or more impurities in the raw material sensitize photodegradation of the drug (k<sub>obs</sub> = 3.8•10<sup>-9</sup> M s<sup>-1</sup>, R<sup>2</sup> = 0.947). Samples spiked with photochemical degradation products prior to exposure at 455 nm show increased degradation. Degradation products that are formed initially seem to be the most important, as the photosensitized degradation after irradiation in the Suntest for 5 min. (k<sub>obs</sub> = 1.2•10<sup>-8</sup> M s<sup>-1</sup>, R<sup>2</sup> = 0.999) and 15 min. (k<sub>obs</sub> = 1.5•10<sup>-8</sup> M s<sup>-1</sup>, R<sup>2</sup> = 0.985) are in the same range. The <sup>1</sup>O<sub>2</sub> quencher sodium azide (NaN<sub>3</sub>) prevents the reaction at the concentration added (10 mM). No significant degradation was observed in 5 min. irradiated samples during monochromatic irradiation at 455 nm for 15 min. after addition of the quencher, or in non-decomposed samples that were irradiated monochromatically for 30 or 60 min. in the presence of the quencher. These results support the hypothesis of <sup>1</sup>O<sub>2</sub> mediated photosensitized degradation of primaquine by impurities of the raw material and photodegradation product(s), although we were not able to detect <sup>1</sup>O<sub>2</sub> luminescence at 1270 nm by excitation at 455 nm. Low concentration(s) of the sensitizer(s) combined with low <sup>1</sup>O<sub>2</sub>-quantum yield(s) may be the reason.

The fluorescence quantum yield for the photodegradation product(s) of primaquine in D<sub>2</sub>O appears to be low (Φ<sub>F</sub> ~ 6•10<sup>-4</sup>), and the fluorescence lifetime is short (τ<sub>F</sub> < 0.5 ns). The fluorescence quantum yield of primaquine in water is low (Φ<sub>F</sub> ~ 9•10<sup>-4</sup>) and is only slightly increased in D<sub>2</sub>O (Φ<sub>F</sub> ~ 2•10<sup>-3</sup>). The fluorescence lifetime of primaquine is short (τ<sub>F</sub> < 0.5 ns) in water and in D<sub>2</sub>O.

Addition of NaN<sub>3</sub> partly retards photodegradation of primaquine in D<sub>2</sub>O when irradiated in the Monochromator at 350 nm, as percentage degradation is reduced from 10(±1) % (n = 2) to 5(±2) % (n = 4) after 30 min. irradiation in the presence of the quencher. Hence free radical reactions also contribute to photodegradation of primaquine, as postulated earlier by a series of scavenging experiments [3, 4, 7]. Formation and reactivity of the monoprotated cation radical is expected to play an essential part in the initial photochemical degradation of primaquine at physiological pH, as illustrated in Scheme 2. Formation of new photosensitizers (PQox) during drug photodegradation will accelerate the initially slow degradation rate with time, as previously observed [4].

## Acknowledgements

The authors are grateful to the STFC Daresbury Laboratory, Warrington, UK, for providing facilities and staff to perform the pulse radiolysis and laser flash photolysis experiments at the Free Radical Research Facility in the Synchrotron Radiation Department (Stations 0.1 and 0.2).

Many thanks to Jon Reierstad and Tove Larsen at the University of Oslo for help with drawing and scanning of figures.

## References

- [1] Editorial, WHO Drug Information Essential Drugs. *Malaria* 2, 1988, pp. 79-93.
- [2] S. Kristensen, A.-L. Grislingaas, J.V. Greenhill, T. Skjetne, J. Karlsen, H.H. Tønnesen, Photochemical stability of biologically active compounds: V. Photochemical degradation of primaquine in an aqueous medium, *Int. J. Pharm.* 100 (1993) 15-23.
- [3] S. Kristensen, L. Grinberg, H.H. Tønnesen, Photoreactivity of biologically active compounds. XI. Primaquine and metabolites as radical inducers, *Eur. J. Pharm. Sci.* 5 (1997) 139-146.
- [4] S. Kristensen, K. Nord, A.-L. Orsteen, H.H. Tønnesen, Photoreactivity of biologically active compounds. XIV. Influence of oxygen on light induced reactions of primaquine, *Pharmazie* 53 (1998) 98-103 .
- [5] S. Kristensen, Photoreactivity of biologically active compounds. XVII. Influence of solvent interactions on spectroscopic properties and photostability of primaquine, *Pharmazie* 60 (2005) 426-433.
- [6] S. Kristensen, J. Karlsen, H.H. Tønnesen, Photoreactivity of biologically active compounds. VI. Photohemolytical properties of antimalarials in vitro, *Pharm. Sci. Commun.* 4 (1994) 183-191.
- [7] S. Kristensen, R.-H. Wang, H.H. Tønnesen, J. Dillon, J.E. Roberts, Photoreactivity of biologically active compounds. VIII. Photosensitized polymerization of lens proteins by antimalarial drugs in vitro, *Photochem. Photobiol.* 61 (1995) 124-130.
- [8] M. Summerfield, G.R. Tudhope, Studies with primaquine in vitro: Superoxide radical formation and oxidation of haemoglobin, *Br. J. Clin. Pharmacol.* 6 (1978) 319-323.
- [9] P.J. Thornalley, A. Stern, J.V. Bannister, A mechanism for primaquine mediated oxidation of NADPH in red blood cells, *Biochem. Pharmacol.* 32 (1983) 3571-3575.
- [10] J.M. Megaw, L.A. Drake, *Photobiology: An overview*, in: E.M. Jackson (Ed.), *Photobiology of the Skin and Eye*, Marcel Dekker, New York, 1986, pp. 1-31.
- [11] R.V. Bensasson, J.C. Gramain, Benzophenone triplet properties in acetonitrile and water, reduction by lactams, *J. Chem. Soc. Faraday Trans. 1* (1980) 1801-1810.



- [12] R. Schmidt, C. Tanielian, R. Dunsbach, C. Wolf, Phenalenone, a universal referee compound for the determination of quantum yields of singlet oxygen  $O_2 (^1\Delta_g)$  sensitization, *J. Photochem. Photobiol. A* 79 (1994) 11-19.
- [13] S. Navaratnam, S.A. Jones, Primary process in the photochemistry of fenbufen in acetonitrile, *Photochem. Photobiol. A* 132 (2000) 175-180.
- [14] S. Navaratnam, Photochemical and photophysical methods used in the study of drug photoreactivity, in: H.H. Tønnesen (Ed.), *Photostability of Drugs and Drug Formulations*, second ed., CRC Press, Boca Raton, 2004, pp. 255-284.
- [15] S. Navaratnam, J. Claridge, Primary photophysical properties of ofloxacin, *Photochem. Photobiol* 72 (2000) 283-290.
- [16] M.A.J. Rodgers, P. Snowden, Lifetime of  $O_2 (^1\Delta_g)$  in water as determined by time resolved infrared luminescence measurements, *J. Am. Chem. Soc.* 104 (1982) 5541-5543.
- [17] S. Navaratnam, I. Hamblett, H.H. Tønnesen, Photoreactivity of biologically active compounds. XVI. Formation and reactivity of free radicals in mefloquine, *J. Photochem. Photobiol. B* 56 (2000) 25-38.
- [18] D.J. Holder, D. Allan, E.J. Land, S. Navaratnam, Establishment of pulse radiolysis facility on the SRS linac at Daresbury laboratory, in: *Proceedings of EPAC, 2002*, pp. 2804-2806.
- [19] J. Butler, B.W. Hodgson, B.M. Hoey, E.J. Land, J.S. Lea, E.J. Lindley, F.A.P. Rushton, A.J. Swallow, Experimental studies of some moderately fast processes initiated by radiation, *Radiat. Phys. Chem.* 34 (1989) 633-646.
- [20] G.E. Adams, J.W. Boag, B.D. Michael, J. Current, The pulse radiolysis of thiocyanate ion, in: M. Ebert, J.P. Keen, A.J. Swallow, J.H. Baxendale (Eds.), *Pulse Radiolysis*, Academic Press, London, 1965, pp. 117-129.
- [21] J.G. Calvert, J.N. Pitts Jr., *Photochemistry*, John Wiley and Sons, New York, 1966, p. 799.
- [22] C.D. Hufford, J.D. McChesney, Assignments of dissociation constants of primaquine by  $^{13}C$ -NMR spectroscopy, *J. Heterocycl. Chem.* 20 (1983) 273-275.
- [23] G. Viola, A. Salvador, L. Ceconet, G. Basso, D. Vadaldi, F. Dall'Acqua, G.G. Aloisi, M. Amelia, A. Barnarfina, L. Latterini, F. Ellsei, Photophysical properties and photobiological behaviour of amodiaquine, primaquine and chloroquine, *Photochem. Photobiol.* 83 (2007) 1415-1427.
- [24] R.H. Bisby, One-electron reduction of the antimalarial drug primaquine, studied by pulse radiolysis, *Free. Rad. Res. Comms.* 5 (1988) 117-124.
- [25] K.L. Stevenson, G.A. Papadantonakis, P.R. LeBreton, Nanosecond UV laser photoionization of aqueous tryptophan: temperature dependence of quantum yield,

mechanism, and kinetics of hydrated electron decay, *J. Photochem. Photobiol. A* 133 (2000)159-167.

[26] C.U. Valencia, E. Lemp, A.L. Zanocco, Quantum yields of singlet molecular oxygen O<sub>2</sub> (<sup>1</sup>Δ<sub>g</sub>), produced by antimalaric drugs in organic solvents, *J. Chil. Chem. Soc.* 48 (2003) 17-21.

[27] A.G. Motten, L.J. Martinez, N. Holt, R.H. Sik, K. Reszka, C.F. Chignell, Photophysical studies on antimalarial drugs, *J. Phys. Chem.* 98 (1999) 10352-10357.

[28] J.-C. Mialocq, Picosecond study of electron ejection in aqueous phenol and phenolate solutions, *J. Chem. Phys.* 72 (1980) 6338-6344.

[29] K.A. Connors, L.A. Gordon, V.J. Stella, Chemical stability of Pharmaceuticals. A Handbook for Pharmacists, second ed., John Wiley & Sons, New York, 1986, p. 93.

[30] P. Wardman, One electron reduction potentials of radicals in aqueous solutions, *J. Phys. Chem. Ref. Data.* 18 (1989) 1637-1755.

## Figure and table legends

### Figure 1

Transient difference absorption spectra from laser flash photolysis (355 nm) of primaquine in aqueous solution at pH 7.2 (laser energy  $\sim 50$  mJ/pulse).

A: Primaquine (0.1 mM) in  $N_2$ -saturated solution 0.5  $\mu s$  (■) and 5  $\mu s$  (□) after the pulse; Inset – transient decays at 410 nm (a) and 720 nm (b). B: Primaquine (0.3 mM) in air-saturated solution, 5  $\mu s$  after the pulse: Insets – power dependence of transient absorbance extrapolated to zero time at 700 nm (■) for  $N_2$  saturated 0.1 mM primaquine solution and 410 nm (□) for air saturated 0.3 mM primaquine solution as a) linear, and b) log-log plots.

### Figure 2

A plot of initial emission intensity (mV) at 1270 nm due to  $^1O_2$  as a function of laser energy (mJ) at 355 nm for  $O_2$ -saturated toluene solution of primaquine-base (●) and air-saturated toluene solution of perinaphthenone (□). Both solutions had the same absorbance of 0.25 at 355 nm.

### Figure 3

Transient difference absorption spectra observed at 7  $\mu s$  (□) and 191  $\mu s$  (●) after pulse radiolysis of primaquine (0.2 mM) in  $N_2O$ -saturated 10 mM phosphate buffer pH 7.4 (spectra normalised to a Dose of 8 Gy). Inset shows the changes in transient absorption at 400 nm ( $PQH^{2+}$ •).

### Figure 4

Transient difference absorption spectra observed at 5  $\mu s$  (□) and 51  $\mu s$  (●) after pulse radiolysis of primaquine (0.2 mM) in  $N_2O$ -saturated 10 mM phosphate buffer pH 7.4 containing 10 mM KBr (spectra normalised to a Dose of 8 Gy). Inset shows the changes in transient absorption at 410 nm ( $PQH^{2+}$ •).

### Figure 5

Lifetime of  $PQH^{2+}$ • (ms) as a function of sample oxygen concentration (% saturation) studied by pulse radiolysis of primaquine (0.2 mM) in  $O_2/N_2O$ -saturated 10 mM phosphate buffer pH 7.4 containing 10 mM KBr, detected as transient decay at 410 nm ( $R^2 = 0.980$ ). Inset shows the changes in transient absorption at 410 nm under 75 %  $O_2$ -saturation.

### Figure 6

Transient difference absorption spectra observed at 7  $\mu s$  (□), 27  $\mu s$  (●) and 69  $\mu s$  (Δ) after pulse radiolysis of primaquine (0.5 mM) in the presence of tryptophan (5 mM) in  $N_2O$ -saturated 10 mM phosphate buffer pH 7.4 containing 0.1 M KBr. Insets show the changes in transient absorption at 520 nm and 410 nm, respectively.

### Figure 7

Corrected (for ground state bleaching) transient absorption spectra at various degree of primaquine ionization, studied by pulse radiolysis of primaquine (0.5 mM) in  $N_2O$ -saturated aqueous solutions containing 10 mM KBr: pH 2.3 detected 6  $\mu s$  after the pulse (●), pH 5.6 detected 3  $\mu s$  after the pulse (□), pH 10.1 detected 8  $\mu s$  after the pulse (Δ).

### Figure 8

Transient absorbance extrapolated to the pulse at 560 nm as a function of pH, observed by pulse radiolysis of primaquine (10  $\mu$ M) in N<sub>2</sub>O-saturated aqueous solutions containing 10 mM KBr.

### Figure 9

Photosensitized degradation of primaquine in D<sub>2</sub>O by irradiation at 455 ( $\pm$  20) nm (72 mV). The samples contained primaquine diphosphate ( $\circ$ ); primaquine diphosphate ( $4 \cdot 10^{-5}$  M) pre-irradiated in the Suntest CPS (Atlas) for 5 min. ( $\blacksquare$ ) or primaquine diphosphate ( $4 \cdot 10^{-5}$  M) pre-irradiated in the Suntest CPS (Atlas) for 15 min. and supplied with an additional amount of drug prior to monochromatic irradiation ( $\blacktriangle$ ). Each point is the mean of 3 measurements ( $R^2 \geq 0.946$ ).

### Scheme 1

Postulated formation of primaquine (PQ) cation radicals at various degree of drug protonation.

### Scheme 2

Postulated mechanisms for the photosensitized degradation of primaquine (PQH<sup>+</sup>) in a neutral aqueous medium.

Sens = photosensitizing impurities in the raw material

PQox = oxidized photodegradation products of primaquine

**Table 1**  
**A summary of the photophysical properties and reaction rates of primaquine transients**

Transient / reaction	Solvent / pH	$\epsilon$ ( $M^{-1}cm^{-1}$ ) ( $\lambda_{max}$ , nm)	$k_2$ ( $M^{-1}s^{-1}$ ) ( $\lambda$ , nm)	${}^{PQ}\Phi_{\Delta}$	$k_q$ ( $M^{-1}s^{-1}$ )
PQ <sup>+</sup> •	Aqueous pH 10.1	5500 (385) 800 (470)			
PQH <sup>2+</sup> •	Aqueous pH 7.4	1600 (650) 7300 (400) 1700 (550)	$1.0 \pm 0.2 \cdot 10^9$ (650) $7.4 \pm 0.6 \cdot 10^8$ (410)		
	Aqueous pH 5.6	7300 (400) 1800 (550)	$7.1 \pm 0.5 \cdot 10^8$ (560)		
	Aqueous pH 2.3	6500 (410) 2000 (570)	$6.0 \pm 0.8 \cdot 10^8$ (560)		
	PQ•- / PQH•	Aqueous pH 7.4	4700 (350) 2100 (490)	$2.0 \pm 0.2 \cdot 10^9$ (490)	
<sup>1</sup> O <sub>2</sub> *	Toluene			0.025 ( $R^2 \geq 0.977$ )	
PQ + <sup>3</sup> FEN* →	Acetonitrile				$3.0 \pm 0.5 \cdot 10^9$
PQ + <sup>3</sup> BP* →	Toluene				$3.9 \pm 0.5 \cdot 10^9$
PQ + Br <sub>2</sub> •- →	Aqueous pH 10.8				$4.2 \pm 0.1 \cdot 10^9$
PQH <sub>2</sub> <sup>2+</sup> + Br <sub>2</sub> •- →	Aqueous pH 2.4				$2.7 \pm 0.2 \cdot 10^9$
PQH <sup>+</sup> + Br <sub>2</sub> •- →	Aqueous pH 7.4				$4.7 \cdot 10^9$ ( $R^2 = 0.998$ )
PQH <sup>+</sup> + OH• →	Aqueous pH 7.4				$6.6 \cdot 10^9$ ( $R^2 = 0.957$ )
PQH <sup>+</sup> + TRP• →	Aqueous pH 7.4				$2.5 \pm 0.3 \cdot 10^7$
PQH <sup>+</sup> + O <sub>2</sub> • <sup>-</sup> →	Aqueous pH 7.4				No reaction
PQH <sup>2+</sup> • + O <sub>2</sub> →	Aqueous pH 7.4				$1.0 \cdot 10^6$ ( $R^2 = 0.962$ )

Abbreviations: PQ = primaquine; FEN = fenbufen; BP = biphenyl; TRP = tryptophan;  $\epsilon$  = molar absorption coefficient;  $\lambda_{max}$  = absorption maximum;  $k_2$  = second order rate constant of transient decay, measured at wavelength  $\lambda$ ;  ${}^{PQ}\Phi_{\Delta}$  = quantum yield of <sup>1</sup>O<sub>2</sub>-formation sensitized by primaquine;  $k_q$  = bimolecular quenching rate constant.