



University of
Salford
MANCHESTER

Propagation and stability of two-colour spatial optical solitons

Bostock, C, Christian, JM and McDonald, GS

Title	Propagation and stability of two-colour spatial optical solitons
Authors	Bostock, C, Christian, JM and McDonald, GS
Type	Conference or Workshop Item
URL	This version is available at: http://usir.salford.ac.uk/18421/
Published Date	2011

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: usir@salford.ac.uk.

Propagation and stability of two-colour spatial optical solitons

C. Bostock, J. M. Christian, and G. S. McDonald

Materials & Physics Research Centre, University of Salford, U.K.

Two-colour spatial solitons comprise coupled nonlinear optical beams at two distinct temporal frequencies [1]. The components (which may be bright-like and/or dark-like) are localized in space and tend to overlap, thereby allowing the interplay between diffraction and nonlinear effects to result in stationary light structures. We will propose a more complete and realistic model for describing such phenomena. A key feature of our approach is that one may access multi-colour geometries involving beam propagation at *arbitrary angles and orientations* with respect to the reference direction – such considerations are central to multiplexing and interface scenarios, but lie far outside the reach of conventional theory. The modulational instability problem can be solved in a range of physically relevant regimes, and extensive computations have confirmed theoretical predictions (see figure 1). New families of exact analytical two-colour solitons are reported, each of which has *co-propagation* and *counter-propagation* classes that are related by geometrical transformation.




Figure 1. Modulational instability of the bright-dark soliton family – bright component in (a), dark component in (b) – in a focusing Kerr medium. Instability develops initially on the plane-wave background of the dark component, leading to filamentation. Nonlinearity provides a mechanism whereby this instability subsequently feeds through the system to destabilize the bright component.

References

- [1] R. De La Fuente and A. Barthelemy, *Opt. Commun.* **88**, 419–423(1992).
- [2] M. Shalaby and A. J. Barthelemy, *IEEE J. Quantum Electron.* **28**, 2736–2741 (1992).