



University of
Salford
MANCHESTER

Peristaltic wave propagation in compressible Maxwell fluids with boundary slip

Christian, JM, Tsiklauri, D and McDonald, GS

Title	Peristaltic wave propagation in compressible Maxwell fluids with boundary slip
Authors	Christian, JM, Tsiklauri, D and McDonald, GS
Type	Conference or Workshop Item
URL	This version is available at: http://usir.salford.ac.uk/id/eprint/18432/
Published Date	2006

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: usir@salford.ac.uk.

Peristaltic Wave Propagation in Compressible Maxwell Fluids with Boundary Slip

J. M. Christian, D. Tsiklauri and G. S. McDonald

Joule Physics Laboratory, Institute for Materials Research, School of Computing, Science and Engineering,
University of Salford, Greater Manchester, M5 4WT

Symposium: Theory and simulation of nanoscale and biological systems

Boundary slip has recently been observed by Craig et al. [Phys Rev Lett 87, 054504 (2001)]. Complementing this work, Tsiklauri [J Acoust Soc Am 112, 843 (2002)] incorporated non-zero boundary slip into a theoretical model of fluid-saturated porous media. This extended work by Tsiklauri and Beresnev [Phys Rev E 64, 036303 (2001)], which predicted new phenomena through a generalization of the classic description of peristaltic pumping: non-Newtonian effects led to unexpected changes in fluid behaviour within the deeply viscoelastic regime. Here, we present the most general description of peristaltic wave propagation that has been developed to date, capturing a wide variety of physical effects (including non-zero boundary slip, viscoelasticity, and compressibility). It thus provides an ideal analytical platform for investigating with greater rigour phenomena such as backflow, which has been suggested as a likely cause of vesico-ureteral reflux in urology. We have considered the peristaltic pumping of blood in a human artery, and have found that small changes in arterial radius can, in some parameter regimes, give rise to a spontaneous backflow effect. A distinct advantage with our model is that the generality lends itself to the description of flow processes in a range of other physical systems, including geophysical and industrial.