



University of  
**Salford**  
MANCHESTER

# Arbitrary-angle scattering of spatial solitons from dielectric optical interfaces

McCoy, EA, Christian, JM and McDonald, GS

<b>Title</b>	Arbitrary-angle scattering of spatial solitons from dielectric optical interfaces
<b>Authors</b>	McCoy, EA, Christian, JM and McDonald, GS
<b>Type</b>	Conference or Workshop Item
<b>URL</b>	This version is available at: <a href="http://usir.salford.ac.uk/id/eprint/18435/">http://usir.salford.ac.uk/id/eprint/18435/</a>
<b>Published Date</b>	2011

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.


For more information, including our policy and submission procedure, please contact the Repository Team at: [usir@salford.ac.uk](mailto:usir@salford.ac.uk).

# Arbitrary-angle scattering of spatial solitons from dielectric optical interfaces

E. A. McCoy, J. M. Christian, and G. S. McDonald

*Materials & Physics Research Centre, University of Salford, U.K.*

Wave–interface phenomena appear throughout Nature and are intrinsically angular in character. For instance, the behaviour of light waves at material boundaries underpins the entire field of optics. The seminal works on nonlinear beam refraction [1] considered scalar bright spatial solitons impinging on the interface between two Kerr-type media with different dielectric parameters, but where angles of incidence, reflection and refraction (relative to the interface) were constrained to be near-negligibly small. Our Group has recently been developing new mathematical and computational tools to describe *arbitrary-angle* refraction of similar beams [2]. We will report our latest research involving higher-order material effects (e.g., quintic-type nonlinearity). A generalized Snell's law describing beam refraction will be detailed and tested directly against fully-numerical calculations [see figure 1 (a)]. New effects are also uncovered, through simulations, in both small- and large-angle regimes [e.g., see figure 1 (b)].



**Figure 1.** (a) Comparison of theoretical Snell's-Law predictions (solid lines) against full numerical computations (points) for a range of interface parameters. (b) Simulation showing a non-trivial Goos-Hänchen shift for an incident Helmholtz spatial soliton that is close to the critical angle.

## References

- [1] A. B. Aceves, J. V. Moloney, and A. C. Newell, *Phys. Rev. A* **39**, 1809–1827 (1989); *Phys. Rev. A* **39**, 1828–1840(1989).
- [2] J. Sánchez-Curto, P. Chamorro-Posada, and G. S. McDonald, *J. Opt. A: Pure Appl. Opt.* **11** art. no. 054015 (2009); *Opt. Lett.* **32**, 1126–1128 (2007).