



University of  
**Salford**  
MANCHESTER

# The role of boundary conditions in kaleidoscope laser modes

McDonald, GS, Christian, JM and Huang, JG

<b>Title</b>	The role of boundary conditions in kaleidoscope laser modes
<b>Authors</b>	McDonald, GS, Christian, JM and Huang, JG
<b>Type</b>	Conference or Workshop Item
<b>URL</b>	This version is available at: <a href="http://usir.salford.ac.uk/18436/">http://usir.salford.ac.uk/18436/</a>
<b>Published Date</b>	2011

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: [usir@salford.ac.uk](mailto:usir@salford.ac.uk).


# The role of boundary conditions in kaleidoscope laser modes

G. S. McDonald<sup>1</sup>, J. M. Christian<sup>1</sup>, and J. G. Huang<sup>2</sup>

<sup>1</sup> *Materials & Physics Research Centre, University of Salford, U.K.*

<sup>2</sup> *School of Engineering and Mathematical Sciences, City University London, London, U.K.*

The complex character of transverse eigenmodes in one-dimensional (1D) unstable cavity lasers has been known for many years [1]. Early collaborations showed that the origin of such fractal (i.e., multiple spatial scale) structure lies in a subtle interplay between small-scale diffraction effects at the mirror edges and successive round-trip magnifications [2]. Kaleidoscope lasers are intuitive generalizations of the classic strip resonator to fully-2D geometries where the feedback mirror has a non-trivial transverse shape, such as a regular polygon [3]. The fundamental mechanism for fractal formation is preserved, but until recently these novel laser designs have remained largely unexplored. We will report on recent advances in our understanding of kaleidoscope lasers, made possible by new semi-analytical techniques. Aspects to be covered include mode patterns (see figure 1), eigenvalue spectra, convergence phenomena, and also the first calculations of fractal dimension for arbitrary cavity parameters.



**Figure 1.** Top row: Computations of the lowest-loss modes for a range of kaleidoscope lasers. Bottom row: magnification of the central portion of each of the corresponding patterns in the top row.

## References

- [1] G. P. Karman and J. P. Woerdman, *Opt. Lett.* **23**, 1909–1911 (1998).
- [2] G. H. C. New, M. A. Yates, J. P. Woerdman, and G. S. McDonald, *Opt. Commun.* **193**, 261–266 (2001).
- [3] G. S. McDonald *et al.*, *J. Opt. Soc. Am. B* **17**, 524–529 (2000); *Nature* **402**, 138 (1999).