



University of
Salford
MANCHESTER

Helmholtz solitons: a new angle in nonlinear science

Chamorro-Posada, P, McDonald, GS, Sanchez-Curto, J and Christian, JM

Title	Helmholtz solitons: a new angle in nonlinear science
Authors	Chamorro-Posada, P, McDonald, GS, Sanchez-Curto, J and Christian, JM
Type	Conference or Workshop Item
URL	This version is available at: http://usir.salford.ac.uk/id/eprint/18438/
Published Date	2007

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: usir@salford.ac.uk.

Helmholtz solitons: A new angle in nonlinear science

Pedro Chamorro-Posada¹
Julio Sánchez-Curto¹

Graham S. McDonald²
James M. Christian²

April 24, 2007

1. Dpto. de Teoría de la Señal y Comunicaciones e I. T., Universidad de Valladolid, ETSIT, Campus Miguel Delibes s/n, 47011 Valladolid, Spain.
2. Joule Physics Lab., School of Computing, Science and Engineering, University of Salford, Salford M5 4WT, UK.

The ubiquitous nonlinear Schrödinger (NLS) equation models soliton propagation in many different scenarios. In a nutshell, the NLS equation describes the slow modulation of the envelope of a linear wave when propagating in a weakly nonlinear medium. Within the context of spatial optical solitons, for instance, *slow* refers to the paraxial approximation which permits to derive the NLS equation from the more general nonlinear Helmholtz (NLH) equation [1] for the envelope of an optical beam which propagates along a definite axis, or infinitesimally close to it, and which is broad when compared to the wavelength, thus, preserving the weakly nonlinear nature of the propagating disturbance.

Restrictions on the propagation angle can be released using the NLH equation which restores the spatial symmetry required when angular considerations are fundamental such as in the collisions of solitons or when they impinge on a nonlinear interface [1]. A detailed account of recent progress in the field [1] plus new results will be presented, including exact soliton solutions and analytical and numerical investigations of their properties.

References

- [1] Chamorro-Posada P. et al. *Phys. Rev. E* **74** 036609 (2006), Sánchez-Curto J. et al. *Opt. Lett.* **32** 1126 (2007), Christian J.M. et al. *J. Phys.A: Math. Gen* **40** 1545 (2007).