



University of
Salford
MANCHESTER

Families of two-colour Helmholtz spatial solitons

Bostock, C, Christian, JM and McDonald, GS

Title	Families of two-colour Helmholtz spatial solitons
Authors	Bostock, C, Christian, JM and McDonald, GS
Type	Conference or Workshop Item
URL	This version is available at: http://usir.salford.ac.uk/id/eprint/18449/
Published Date	2011

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: usir@salford.ac.uk.

Families of Two-Colour Helmholtz Spatial Solitons

C. Bostock, J. M. Christian, and G. S. McDonald

Materials & Physics Research Centre, University of Salford, Greater Manchester, M5 4WT, U.K.

Multi-colour spatial solitons comprise localized optical components at distinct temporal frequencies [1]. The components (which may be bright-like and dark-like) tend to overlap in space, thereby allowing the interplay between linear spreading (diffraction) and nonlinear effects (self- and mutual-focusing) to result in an electromagnetic structure with a stationary intensity pattern. Two-colour spatial solitons for a Kerr-type medium were proposed by De La Fuenete and Barthelemy [2] within the context of an intuitive nonlinear Schrödinger model. Subsequent experiments, using continuous-wave (CW) laser light at red and green wavelengths, demonstrated that such mutually-trapped light beams could be generated in CS_2 waveguides [3]. This opened up the possibility of multi-colour photonic device applications and architectures [4].

Here, we introduce a novel Helmholtz model for two-colour CW optical fields whose temporal frequency separation is similarly large. A key advantage of our approach is that it allows one full access to multi-component geometries involving propagation at arbitrary angles and orientations with respect to the reference direction [5] – such considerations are central to off-axis configurations involving, for instance, beam multiplexing [6] and interface [7] scenarios. In contrast, classic paraxial models [2,3] capture angles (in the laboratory frame) that are negligibly, or near-negligibly, small [4]. The two-colour modulational instability problem can be solved in a range of physically relevant regimes. Bright-bright and bright-dark solitons are also reported, each of which having *co-propagation* and *counter-propagation* solution classes that are connected by geometrical transformation. Extensive computations [8] have confirmed the validity of analyses (see Fig. 1).




Fig. 1 Inherent instability of the exact analytical bright-dark Helmholtz soliton – bright component shown in (a), dark component in (b). Modulational instability develops initially on the finite-amplitude plane-wave background of the dark component in u_2 , leading to filamentation [the dominant spatial frequency in part (b) is predicted by linear analysis]. This instability then feeds through the system, via nonlinear coupling, to destabilize the bright component u_1 .

References

- [1] P. B. Lindquist, D. R. Andersen, and Y. S. Kivshar, “Multicolor solitons due to four-wave mixing,” *Phys. Rev. E* **57**, 3551 (1998).
- [2] R. De La Fuente and A. Barthelemy, “Spatial solitons pairing by cross phase modulation,” *Opt. Commun.* **88**, 419 (1992).
- [3] M. Shalaby and A. J. Barthelemy, “Observation of the self-guided propagation of a dark and bright spatial soliton pair in a focusing nonlinear medium,” *IEEE J. Quantum Electron.* **28**, 2736 (1992).
- [4] H. T. Tran, R. A. Sammut, and W. Samir, “Interaction of self-guided beams of different frequencies,” *Opt. Lett.* **19**, 945 (1994).
- [5] J. M. Christian, G. S. McDonald, and P. Chamorro-Posada, “Helmholtz-Manakov solitons,” *Phys. Rev. E* **74**, art. no. 066612 (2006).
- [6] P. Chamorro-Posada and G. S. McDonald, “Spatial Kerr soliton collisions at arbitrary angles,” *Phys. Rev. E* **74**, art. no. 036609 (2006).
- [7] J. Sánchez-Curto, P. Chamorro-Posada, and G. S. McDonald, “Helmholtz solitons at nonlinear interfaces,” *Opt. Lett.* **32**, 1126 (2007).
- [8] P. Chamorro-Posada, G. S. McDonald, and G. H. C. New, “Nonparaxial beam propagation methods,” *Opt. Commun.* **192**, 1 (2001).