



University of  
**Salford**  
MANCHESTER

# Propagation properties of Helmholtz power-law nonlinear surface waves

McCoy, EA, Christian, JM and McDonald, GS

<b>Title</b>	Propagation properties of Helmholtz power-law nonlinear surface waves
<b>Authors</b>	McCoy, EA, Christian, JM and McDonald, GS
<b>Type</b>	Conference or Workshop Item
<b>URL</b>	This version is available at: <a href="http://usir.salford.ac.uk/id/eprint/23001/">http://usir.salford.ac.uk/id/eprint/23001/</a>
<b>Published Date</b>	2012

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: [usir@salford.ac.uk](mailto:usir@salford.ac.uk).

## **Propagation properties of Helmholtz power-law nonlinear surface waves**

E. A. McCoy, J. M. Christian, and G. S. McDonald  
*Materials & Physics Research Centre,  
University of Salford, Greater Manchester M5 4WT, UK*

### **KEYWORDS:**

surface waves, spatial solitons, self-focusing, nonlinear materials, optical interfaces

### **THEME 1: NONLINEAR DYNAMICS**

#### **APPLICATION THEME 1: PHYSICAL SCIENCES**

Optical surface waves are a fundamental class of excitation in inhomogeneous nonlinear photonic systems. This type of laser light satisfies a Helmholtz-type governing equation, and is subject to certain boundary conditions. It also has an asymmetric cross-sectional shape due to abrupt changes in material properties that define the (e.g., planar) interface. The stability properties of surface-wave solution branches are notoriously difficult to predict. On the one hand, classic criteria (e.g., Vakhitov-Kolokolov) often fail; on the other hand, numerical computations have, historically, been performed only with approximated governing equations (e.g., of the Schrödinger type).

In recent research endeavours [J. M. Christian *et al.*, *J. At. Mol. Opt. Phys.*, *in press*], we have derived the surface waves for a Helmholtz-type interface model with refractive-index profile that depend on the local light amplitude to an arbitrary power  $0 < q < 4$ . This generic optical nonlinearity describes classes of semiconductors, doped filter glasses, and liquid crystals.

Here, we will present the first full investigation of Helmholtz nonlinear surface waves. Exact analytical solutions will be reported, and their properties explored. Extensive simulations with the full (i.e., un-approximated) governing equation have addressed some key issues surrounding the robustness of surface-wave solutions against spontaneous instabilities. New qualitative phenomena have also been predicted when considering interactions between surface waves and (obliquely-incident) spatial solitons (that is, self-collimated laser beams). The interaction angle between the surface-wave and incoming soliton is found to play a pivotal role in determining post-collision behaviour in the system, as is the nonlinearity exponent  $q$ .