



University of  
**Salford**  
MANCHESTER

# Helmholtz dark solitons: oblique propagation, saturable materials & bistability

Lundie, MJ, McDonald, GS and Christian, JM

<b>Title</b>	Helmholtz dark solitons: oblique propagation, saturable materials & bistability
<b>Authors</b>	Lundie, MJ, McDonald, GS and Christian, JM
<b>Type</b>	Conference or Workshop Item
<b>URL</b>	This version is available at: <a href="http://usir.salford.ac.uk/id/eprint/23015/">http://usir.salford.ac.uk/id/eprint/23015/</a>
<b>Published Date</b>	2012

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: [usir@salford.ac.uk](mailto:usir@salford.ac.uk).

# Helmholtz dark solitons: oblique propagation, saturable materials & bistability


M. J. Lundie, G. S. McDonald, and J. M. Christian

*Materials & Physics Research Centre, University of Salford, U.K.*

Email: [mark\\_lundie@gmx.com](mailto:mark_lundie@gmx.com)

Keywords: Helmholtz solitons, saturable refractive index, waveguide optics

Dark spatial optical solitons comprise a uniform background wave that is modulated by an obliquely-propagating 'dip' in the light intensity profile. Appearing throughout nonlinear science, these universal entities possess a phase topology that endows them with a remarkable degree of stability against perturbations to their shape. Hence, the photonics community is greatly interested in dark solitons for potential use as 'information bits' in future optical technologies. Our Group has been developing Helmholtz soliton theory for the past 14 years. This more sophisticated modelling approach completely eliminates the intrinsic angular limitations of classic paraxial models, where waves must travel along (or at near-negligibly-small angles with respect to) the laboratory longitudinal direction. The most recent system we have analyzed captures the generic effects of a saturable defocusing nonlinearity, where the locally-induced refractive-index change becomes bleached under high-intensity illumination. Families of exact bistable dark solitons have been derived by deploying a unique blend of mathematical methods (see figure 1), and linearization techniques have predicted the inherent stability of the background plane wave against small-amplitude modulations. Our new solutions have a raft of crucial asymptotic properties, complementing their bright [1] and paraxial [2] counterparts.



**Figure 1.** Intensity (left) and phase (right) quadratures of an exact Helmholtz dark soliton for a saturable defocusing nonlinearity as the net propagation angle in the laboratory frame (denoted by  $\theta - \theta_0$ ) is increased.

## References

- [1] J. M. Christian, G. S. McDonald, and P. Chamorro-Posada, *J. Opt. Soc. Am. B* **26**, 2323 (2009).
- [2] W. Krolikowski and B. Luther-Davies, *Opt. Lett.* **18**, 188 (1993).