



University of  
**Salford**  
MANCHESTER

# Dark & anti-dark spatiotemporal solitons: From cubic to cubic-quintic systems

Cowey, R, Christian, J M and McDonald, G S

<b>Title</b>	Dark & anti-dark spatiotemporal solitons: From cubic to cubic-quintic systems
<b>Authors</b>	Cowey, R, Christian, J M and McDonald, G S
<b>Type</b>	Conference or Workshop Item
<b>URL</b>	This version is available at: <a href="http://usir.salford.ac.uk/id/eprint/29296/">http://usir.salford.ac.uk/id/eprint/29296/</a>
<b>Published Date</b>	2013

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: [usir@salford.ac.uk](mailto:usir@salford.ac.uk).

# Dark & anti-dark spatiotemporal solitons: from cubic to cubic-quintic systems


R. Cowey, J. M. Christian, and G. S. McDonald

*Materials & Physics Research Centre, University of Salford, U.K.*

Email: [r.cowey@edu.salford.ac.uk](mailto:r.cowey@edu.salford.ac.uk)

Keywords: dark solitons, anti-dark solitons, spatiotemporal dispersion, *energy theme*

The origin of conventional models for optical pulse propagation lies in the universal slowly-varying envelope approximation (SVEA) accompanied by a Galilean boost to the local time frame. However, Biancalana and Creatore [1] have recently pointed out that the SVEA is not necessarily compatible with physical contexts involving spatial dispersion [2]. Their proposal has prompted us to reassess the way in which conventional pulse theory handles the linear part of the wave operator. In so-doing, we have constructed a framework for scalar optical pulses that is based upon transformation laws and frame-of-reference considerations [3]. Here, we present our most recent research analysing soliton propagation in systems with the universal cubic-quintic type of nonlinearity [4]. By deploying a range of analytical methods, we have been able to derive exact spatiotemporal dark and anti-dark solitons. These structures have a strongly geometrical flavour that is tightly connected to the symmetry properties of the governing equation. Multi-parameter asymptotic analyses demonstrate that classic conventional solitons are a subset of our new solutions. Extensive numerical calculations have also predicted a high degree of robustness of bright and dark pulses [see Figs. 1(a) and 1(b), respectively] against local perturbations to their temporal shape.



**Figure 1.** Simulations showing the stability of perturbed (a) bright (anomalous temporal dispersion) and (b) dark (normal temporal dispersion) spatiotemporal solitons. Blue (circle): unperturbed solution, where the pulse evolves with a stationary shape. Green (triangle), red (square) and black (diamond) curves: increasing perturbation, respectively.

## References

- [1] F. Biancalana and C. Creatore, *Opt. Express* **16**, 14882 (2008).
- [2] V. M Agranovich and V. L. Ginzburg, *Crystal optics with spatial dispersion, and excitons* (Springer, Berlin, 1984).
- [3] J. M. Christian, G. S. McDonald, T. F. Hodgkinson, and P. Chamorro-Posada, *Phys. Rev. Lett.* **108**, 033101 (2012).
- [4] K. I. Pushkarov, D. I. Pushkarov, and I. V. Tomov, *Opt. Quantum Electron.* **11**, 471 (1979).