



University of
Salford
MANCHESTER

Coupled spatiotemporal waves: New paradigms in vector soliton physics

Ashley, J T, Christian, JM and McDonald, GS

Title	Coupled spatiotemporal waves: New paradigms in vector soliton physics
Authors	Ashley, J T, Christian, JM and McDonald, GS
Type	Conference or Workshop Item
URL	This version is available at: http://usir.salford.ac.uk/id/eprint/29298/
Published Date	2013

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: usir@salford.ac.uk.

Coupled spatiotemporal waves: new paradigms in vector soliton physics

J. T. Ashley, J. M. Christian, and G. S. McDonald

Materials & Physics Research Centre, University of Salford, U.K.

Email: j.t.ashley@edu.salford.ac.uk

Keywords: vector solitons, spatiotemporal dispersion, *energy theme*

In this presentation, we propose a novel spatiotemporal generalization of Menyuk's classic equations [1] describing the propagation of two nonlinearly-coupled waves in a dispersive optical system (such as a fibre or planar waveguide). Our approach is similar in spirit to that recently introduced by Christian *et al.* [2] in a single-component context: both analyses are firmly rooted in special relativity-inspired frame-of-reference considerations, and are concerned with the universal properties of waves in fully second-order systems (where evolution is allowed to venture beyond the realms of the ubiquitous slowly-varying envelope approximation). Research highlights include the complete solution of the modulational instability (MI) problem for vectorized spatiotemporal continuous-wave solutions (the characteristic equation turns out to be an eighth-degree polynomial, rather than a quartic [1]). Predictions of the most-unstable frequency in the system, obtained from consultation of the MI spectrum, are in full agreement with extensive numerical calculations [see Figs. 1(a) & 1(b)]. We have also derived exact bright-bright solitons and investigated some of their key characteristics (e.g., robustness and the asymptotic recovery of Menyuk-type solutions). Future work will address spatiotemporal vector dark solitons [3,4] [see Fig. 1(c)].




Figure 1. (a) Typical long-wave MI spectrum when both waves experience anomalous temporal dispersion. (b) Extensive simulations have confirmed the prediction for the most-unstable frequency, denoted by Ω_{P0} [related to the dominant peak in part (a)]. (c) Instability of a conventional dark-dark type of initial condition used for the spatiotemporal governing equation.

References

- [1] C. R. Menyuk, IEEE J. Quantum Electron. **QE-23**, 174 (1987).
- [2] J. M. Christian, G. S. McDonald, T. F. Hodgkinson, and P. Chamorro-Posada, Phys. Rev. Lett. **108**, 034101 (2012).
- [3] V. V. Afanasyev, Y. S. Kivshar, V. V. Konotop, and V. N. Serkin, Opt. Lett. **14**, 805 (1989).
- [4] Y. S. Kivshar and S. K. Turitsyn, Opt. Lett. **18**, 337 (1993).