



University of
Salford
MANCHESTER

Vector spatial solitons beyond the slowly-varying envelope approximation

Bostock, C, Christian, JM and McDonald, GS

Title	Vector spatial solitons beyond the slowly-varying envelope approximation
Authors	Bostock, C, Christian, JM and McDonald, GS
Type	Conference or Workshop Item
URL	This version is available at: http://usir.salford.ac.uk/id/eprint/29314/
Published Date	2013

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: usir@salford.ac.uk.

Vector spatial solitons beyond the slowly-varying envelope approximation

C. Bostock, J. M. Christian, and G. S. McDonald

Materials & Physics Research Centre, University of Salford, Greater Manchester, M5 4WT

Key words: Helmholtz diffraction; nonlinear waves; modulational instability, spatial solitons; vector beams.

The simultaneous propagation of two spatially-overlapping and differently-coloured laser beams (e.g., at infra-red and green wavelengths) in photonic waveguides has attracted much attention since the early 1990s [Shalaby & Barthelemy, *IEEE J. Quantum Electron.* vol. 28, 2736 (1992)]. Historic analyses of the two-colour optical propagation problem have routinely adopted models of the nonlinear-Schrödinger class that are inherently bound by the slowly-varying envelope approximation (SVEA) [De La Fuente & Barthelemy, *Opt. Commun.* vol. 88, 419 (1992)]. While desirable in some respects (e.g., by considerable simplification of the governing equations), the SVEA can place some strong physical constraints on the validity of predictions made by Schrödinger-type systems in context of wave optics and the theory of nonlinear beams. Our approach avoids such limitations by dealing with the more general (i.e., un-approximated) Helmholtz-type governing equations.

In this presentation, we will deliver a concise overview of our investigations into a model of two-colour light beams beyond the SVEA. A plethora of new results will be provided, including the derivation of four exact analytical vector soliton families. A novel linearization technique, which generalizes conventional methods [Agrawal, *J. Opt. Soc. Am. B* vol. 6, 1072 (1990)] to capture the Helmholtz type of governing equation, is also developed to assess the resilience of the system against the emergence of spontaneous instabilities. Theoretical calculations are supported by fully-nonlinear computer simulations. We expect our results to play a key role in the design of future photonic devices that incorporate angular nonparaxial-beam configurations into their two-colour operational architecture and geometry.