



University of  
**Salford**  
MANCHESTER

# Giant Goos-Hänchen shifts and soliton refraction in systems with $\chi(5)$ susceptibility

McCoy, EA, Christian, JM and McDonald, GS

<b>Title</b>	Giant Goos-Hänchen shifts and soliton refraction in systems with $\chi(5)$ susceptibility
<b>Authors</b>	McCoy, EA, Christian, JM and McDonald, GS
<b>Type</b>	Conference or Workshop Item
<b>URL</b>	This version is available at: <a href="http://usir.salford.ac.uk/id/eprint/29315/">http://usir.salford.ac.uk/id/eprint/29315/</a>
<b>Published Date</b>	2013

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: [usir@salford.ac.uk](mailto:usir@salford.ac.uk).

# Giant Goos-Hänchen shifts and soliton refraction in systems with $\chi^{(5)}$ susceptibility

E. A. McCoy, J. M. Christian, and G. S. McDonald

*Materials & Physics Research Centre, University of Salford, Greater Manchester M5 4WT*

*Key words:* nonlinear materials; nonparaxial optics; optical interfaces;  
soliton refraction; spatial solitons;

A full understanding of how a spatial solitons (a unique type of nonlinear light beam) behave when the encounter the boundary between two dissimilar media is still some way off. On the one hand, light-interface interactions underpin essentially all optical technologies. On the other hand, the overwhelming majority of the literature is rooted in paraxial theory [c.f. Aceves *et al.*, Phys. Rev. A vol. 39, 1809 (1989)], whereby angles of incidence, reflection, and refraction are vanishingly small (in the *laboratory frame*).

We will present our latest research into a new class of nonparaxial model describing arbitrary-angle refraction. Novelty arises by supplementing the traditional cubic response – or “ $\chi^{(3)}$  susceptibility” – with an additional quintic contribution called the “ $\chi^{(5)}$  susceptibility” [Pushkarov *et al.*, Opt. Quantum Electron. vol. 11, 471 (1979)]. This term plays an important role, and can dominate the system, when the incident laser light is sufficiently strong. Non-trivial generalizations of results from earlier collaborations, which accommodated susceptibilities only of the cubic type [Sánchez-Curto *et al.*, Opt. Lett. vol. 35, 1347 (2010); vol. 32, 1127 (2007)], will be discussed. Extensive theoretical predictions (from a newly-derived Snell’s law) for beam refraction / critical angles in cubic-quintic material systems have generally been in good agreement with fully-numerical computations.

We have also made, to the best of our knowledge, the first predictions of giant Goos-Hänchen shifts [Goos and Hänchen, Ann. Phys. vol. 1, 333 (1947)] at cubic-quintic interfaces. A range of new qualitative phenomena, directly attributable to the effect of  $\chi^{(5)}$  susceptibility, will be reported.