



University of  
**Salford**  
MANCHESTER

# The spatiotemporal wave equation with a dual power-law nonlinearity

Cowey, R, Christian, JM and McDonald, GS

<b>Title</b>	The spatiotemporal wave equation with a dual power-law nonlinearity
<b>Authors</b>	Cowey, R, Christian, JM and McDonald, GS
<b>Type</b>	Conference or Workshop Item
<b>URL</b>	This version is available at: <a href="http://usir.salford.ac.uk/id/eprint/29318/">http://usir.salford.ac.uk/id/eprint/29318/</a>
<b>Published Date</b>	2013

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: [usir@salford.ac.uk](mailto:usir@salford.ac.uk).

# The spatiotemporal wave equation with a dual power-law nonlinearity

R. Cowey, J. M. Christian, and G. S. McDonald

*Materials & Physics Research Centre, University of Salford, Greater Manchester, M5 4WT*

*Key words:* dispersion, dual power-law nonlinearity, feedback loops, nonlinear wave equation, spatiotemporal solitons.

Solitons and soliton-like emergent phenomena appear ubiquitously throughout nonlinear science. These waves are truly fundamental entities, transcending the physical details of a system and resulting whenever linear dispersion (which tends to broaden a pulse in time) is opposed by nonlinearity (which tends to compress it). Such dynamic competition (feedback loops) may then facilitate the spontaneous formation of self-localizing and self-stabilizing wavepackets (solitons) whose profiles are often bell-shaped in time and stationary (invariant) throughout evolution in space. Here, we consider the classic dual power-law nonlinearity model [Micallef *et al.*, Phys. Rev. E vol. 54, 2936 (1994)] in the recently-proposed context of a spatiotemporal wave equation [Christian *et al.*, Phys. Rev. Lett. vol. 108, art. no. 034101 (2012)]. This type of governing equation has wide applicability in the field of photonics.

Our analysis begins with simple continuous waves (solutions that are temporally flat) and a perturbative investigation of their robustness against small disturbances (e.g., fluctuations due to random background noise). Our attention will then shift to hyperbolic solitons [that are quite tightly (i.e., exponentially) localized in time] and algebraic solitons (that have a much weaker Lorentzian-like localization). These two families are intimately related, with the latter appearing at the threshold for linear wave propagation (analytic continuation subsequently yields a class of temporally-delocalized periodic wave). We have also derived so-called boundary solitons, which connect (in a monotonic way) regions of finite-amplitude ‘flat’ solution to regions of zero amplitude. For each new class of wave, extensive simulations have been used to assess solution stability.