



University of
Salford
MANCHESTER

Alcohol industry sponsorship and hazardous drinking in UK university students who play sport

O'Brien, KS, Ferris, J, Greenlees, I, Jowett, S, Rhind, D, Cook, PA and Kypri, K

<http://dx.doi.org/10.1111/add.12604>

Title	Alcohol industry sponsorship and hazardous drinking in UK university students who play sport
Authors	O'Brien, KS, Ferris, J, Greenlees, I, Jowett, S, Rhind, D, Cook, PA and Kypri, K
Publication title	Addiction
Publisher	Wiley
Type	Article
USIR URL	This version is available at: http://usir.salford.ac.uk/id/eprint/31928/
Published Date	2014

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: library-research@salford.ac.uk.

Alcohol industry sponsorship and hazardous drinking in UK university students who play sport

Kerry S. O'Brien^{1,2*}, Jason Ferris³, Ian Greenlees⁴, Sophia Jowett⁵, Daniel Rhind⁶, Penny Cook⁷, Kypros Kypri⁸.

Behavioural Studies, Monash University, Australia¹; School of Psychological Sciences, University of Manchester, UK²; Institute for Social Science Research, University of Queensland, Australia³; Department of Sport and Exercise Sciences, University of Chichester, UK⁴; School of Sport, Exercise and Health Sciences, Loughborough University, UK⁵; Brunel Centre for Sport, Health and Wellbeing, Brunel University, UK⁶; ; School of Health Sciences, Salford University, UK⁷; School of Medicine & Public Health, University of Newcastle, Australia⁸

Declaration of Interest: The research was supported by grant funding from Alcohol Research UK.

KOB was also supported by funding from the Australian Research Council, VicHealth, and the Australian National Preventive Health Agency. There are no conflicts of interest to declare.

Running Head: Alcohol sponsorship of sport

Key Words: Sport, Athletes, Drinking, Hazardous, Alcohol Sponsorship.

* Author to whom correspondence should be addressed at: School of Political and Social Sciences, Monash University, VIC, Australia.

E-mail: kerrykez@gmail.com

Word Count: 3022

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/add.12604

This article is protected by copyright. All rights reserved.

Abstract

Aim: To examine whether receipt of alcohol industry sponsorship is associated with problematic drinking in UK university students who play sport.

Methods: University students (n=2450) participating in sports were invited to complete a pen-and-paper questionnaire by research staff approaching them at sporting facilities and in university settings. Respondents were asked whether they personally, their team, and/or their club were currently in receipt of sponsorship (e.g., money, free or subsidised travel, or sporting products), from an alcohol-related industry (e.g., bars, liquor stores, wholesalers), and whether they had solicited the sponsorship. Drinking was assessed using the Alcohol Use Disorders Identification Test (AUDIT).

Results: Questionnaires were completed by 2048 of those approached (response rate=83%).

Alcohol industry sponsorship was reported by 36% of the sample. After accounting for confounders (age, gender, disposable income, and location) in multivariable models, receipt of alcohol sponsorship by a team (adjusted $\beta_{adj}=.41$, $p=.013$), club ($\beta_{adj}=.73$, $p=.017$), team and club ($\beta_{adj}=.79$, $p=0.002$), and combinations of individual and team or club sponsorships ($\beta_{adj}=1.27$, $p<0.002$), were each associated with significantly higher AUDIT-Consumption substance scores. Receipt of sponsorship by team and club (aOR=2.04; 95% CI: 1.04-3.99) and combinations of individual and team or club sponsorships (aOR=4.12; 95% CI: 1.29-13.15) were each associated with increased odds of being classified a hazardous drinker (AUDIT score >8). Respondents who sought out sponsorship were not at greater risk than respondents who had, or whose teams or clubs had, been approached by the alcohol industry.

Conclusions: University students in the United Kingdom who play sport and who personally receive alcohol industry sponsorship or whose club or team receives alcohol industry sponsorship appear to have more problematic drinking behaviour than UK university students who play sport and receive no alcohol industry sponsorship. Policy to reduce or cease such sponsorship should be considered.

Introduction

Hazardous consumption of alcohol is a leading cause of mortality and morbidity in middle and high income countries [1], ranked above illicit drug use in terms of social, health, and financial costs [2].

In the UK alone, the cost of hazardous drinking is approximately £25 billion per year [3].

Hazardous drinking is particularly common in young people and university students [4,5], and even more so in university students who play sport [6-12]. Research from the USA, Australia, and New Zealand, suggests that sportspeople, and especially university students who play sport, drink more hazardously than their non-sporting peers and the general population [7-12]. Higher rates of drink-driving, anti-social behaviour, and unprotected sex were also found in university sports participants [6,7,11]. Only one published study has examined drinking in UK university sportspeople, finding that hazardous drinking was more prevalent among sportspeople than their non-sporting peers [12].

There has been no research examining whether alcohol industry sponsorship is associated with increased risk of hazardous drinking in the UK or elsewhere in Europe.

A review of the small literature on factors which might explain heavy drinking among sportspeople suggests that using alcohol for team cohesion and coping, and the drinking of peers probably play a role [7], along with specific practices such as drinking with teammates and opponents after matches [13]. There has been a strong interdependence between sport and alcohol industries for several hundred years in the UK. For example, pubs traditionally hosted and funded sport competitions for customer entertainment and gambling. During the industrial revolution many of the world's largest sporting clubs (e.g., Manchester United) were funded or owned by alcohol industries in the UK [14].

Sport continues to be a primary vehicle for the promotion of alcohol, with a large proportion of the alcohol industry's advertising and sponsorship budget spent on sport [15,16]. Alcohol industry marketing contributes to problem drinking [17-21]. Systematic reviews of longitudinal studies show that early exposure of young people to alcohol advertising and sponsorship is associated with stronger intentions to drink, and higher levels of alcohol consumption in later

adolescence and early adulthood [20,21]. A ban on alcohol advertising and sponsorship has been called for by peak medical bodies in the UK, Ireland, Australia, and South Africa [22-24], and in the UK House of Commons Health Select Committee Report on Alcohol [25]. The UK government's official response to calls for stronger regulation of alcohol advertising and sponsorship was that more evidence is required [26].

Alcohol industry sponsorship of sport refers not only to payments for event naming and product marketing rights (e.g., “Heineken & UEFA Champions League Football”), which are in effect advertising, but also to less conspicuous but potentially more harmful *direct to user* sponsorship [27-29]. This direct sponsorship occurs from grass-roots to elite level sport, encompassing the payment of club fees, provision of uniforms, payment of travel costs, and provision of alcohol at post match functions. In return, sponsored individuals, teams, and clubs are often required to wear the sponsor’s logo and to attend and drink at the sponsor’s premises [28]. For example, a bar provides a local football team with a cash payment to cover uniform costs. In return the team has that bar’s name printed on the jersey and agrees to drink there after games, bringing the opposition team and spectators, friends, and family. In this way the bar attracts new customers and creates a sense of obligation in club members.

A recent multinational EU longitudinal study found that children with indirect exposure to sports clubs receiving alcohol sponsorship reported more positive drinking expectancies and had higher odds of having consumed alcohol in the past month [30]. Studies from New Zealand and Australia found that sportspeople in receipt of alcohol industry sponsorship were more likely to be hazardous drinkers [28,29]. The previous work was unable account for possible confounders such as disposable income, nor whether participants sought out alcohol sponsorship, to examine the possibility that a predilection for drinking leads to sponsorship and not vice versa. There has been no UK research examining the association between alcohol industry sponsorship of sport and hazardous drinking among sportspeople despite both practices being common [31]. Our aim was to

examine whether receipt of alcohol industry sponsorship is positively associated with alcohol consumption and hazardous drinking in UK sportspeople.

Methods

Participants

To circumvent the possibility that some sporting organisations (e.g., clubs) would deny access to sportspeople thereby biasing findings, we approached participants directly at university playing fields, training facilities, and sport-related teaching venues and classes. The aim was to achieve heterogeneity in the exposure of interest, namely, alcohol sponsorship, rather than to estimate prevalence. Although the prevalence of alcohol industry sponsorship of sportspeople in the UK is unknown, alcohol industry practices and sporting cultures appear similar to those in New Zealand and Australia where 47% and 30%, respectively, of respondents approached via the same method were currently in receipt of alcohol sponsorship [28,29]. We therefore expected at least 30% of participants to be in receipt of alcohol industry sponsorship. The primary assumption underlying the inferences we seek to draw are that how participants came into the study would not be systematically related to the association of interest [32], i.e., whether receiving industry funding increases the likelihood of hazardous drinking. The risk of selection bias from this recruitment approach was judged likely to be lower than that arising from non-participation of some sporting bodies who may have considered involvement in the research to create a commercial or reputational risk.

It is also important to note that the participants in this study were not necessarily involved in university sport competitions, which is a common classification in US studies involving National Collegiate Athletic Associations. The university students surveyed here were identified as participants in sport not necessarily run by a university. Given that we cannot know how representative the participants are of the population of sportspeople or even of UK university students engaged in sport, prevalence rates are not estimated.

Sample size estimate

For multiple regression, the rule of thumb $N \geq (8/f^2) + (m-1)$ gives a sample size of 809 where m =number of predictors=10, f^2 =effect size=0.1 [33]. To examine relationships between sponsorship type and AUDIT score among those receiving sponsorship, we assumed that 30% of the sample would receive sponsorship. Accordingly, 450 respondents would be sufficient to detect an f^2 of 0.14, i.e. a small-moderate effect, equivalent to $R^2=12\%$.

Measures

Participants were presented with a short questionnaire assessing demographic details (age, gender, sports played, weekly disposable income, and geographical location), whether respondents had received alcohol industry sponsorship, and the Alcohol Use Disorders Identification Test (AUDIT) which consists of 10 questions with a total score range of 0-40 [34]. Its validity has been thoroughly established with a score of ≥ 8 indicative of hazardous drinking [35]. The AUDIT consumption subscale (AUDIT-C), consisting of the first three questions, is a measure of alcohol consumption, with a range 0-12 [34].

Receipt of alcohol industry sponsorship was assessed using an identical item and response format to those described in previous research [28,29]. In summary, participants were asked if they, their team, or their club currently received sponsorship (e.g., money, equipment, travel costs, discounted/free alcohol) from an alcohol industry body (e.g., a bar, hotel, liquor store, or producer). Participants who were uncertain about their team or club sponsorship arrangements were coded as 'no' for receipt of alcohol industry sponsorship. It should be noted that classification error arising from this assumption being incorrect can only bias associations toward the null.

Heavier drinkers may preferentially seek out alcohol industry sponsorship rather than heavier drinking resulting from receipt of alcohol sponsorship, an important issue in making inferences about the likely direction of any association identified [36]. Accordingly, we asked

participants who reported receiving alcohol industry sponsorship to report whether they, their team, or club, sought out alcohol industry sponsorship or whether they, their team, or club, had been approached by an alcohol industry body offering sponsorship.

Procedure

Data collection occurred between September 2010 and February 2012, encompassing in-season winter and summer sports (see Table 1). Venues for data collection were identified from university webpage listings and competition schedules posted in newsletters. Most venues were owned or leased by universities, and all were within a 2-mile radius of 10 universities situated in the North West, Midlands, London, and Southern region of England. None of the venues had bars associated with them. Non-team sport venues (e.g., tennis) with small numbers of participants present at any one time were visited up to five times.

Upon arrival at venues, researchers approached the nearest sportsperson and invited them to participate. Following acceptance or rejection of the invitation the data collector approached the next nearest sportsperson for participation, and so on until all potential participants at the venue had been approached. Researchers were on hand to assist with participant queries. Participants were offered a nominal incentive of £2 for participation, were informed that their participation would remain confidential and that identifying information was not being requested. The questionnaire took approximately 15 minutes to complete. Ethical approval was obtained from the human research ethics committees of the University of Manchester, Loughborough University, Brunel University, and the University of Chichester.

Questions about exposure of interest

Participants were asked “Do you personally, your team, or club receive sponsorship or support (e.g., financial payments, competition fees, clothing, club pourage rights, other goods) from an alcohol-related industry (e.g., pub, bar, winery, brewer, distillery, hotel, nightclub)? Responses options for

each of the following response categories were: Personally: *no* or *yes*; Team: *no* or *yes* or *uncertain*; and Club: *no* or *yes* or *uncertain*.

Statistical Analyses

Drinking outcomes of interest were alcohol consumption and hazardous drinking status. The AUDIT-C (henceforth referred to as alcohol consumption) was treated as a continuous variable. Participants with AUDIT scores ≥ 8 were classified as hazardous drinkers.

Multiple linear regression models adjusting for age, gender, location and disposable income, were used to examine associations between alcohol sponsorship variables and alcohol consumption scores. Multiple logistic regression models controlling for the same variables were used to estimate associations between sponsorship variables and hazardous drinker status. In all models clustering within sports was accounted for using the STATA *vce* command. Due to the large variation in the number of people within different sport types we also bootstrapped the models. We set bootstrapping to 1000 replications to ensure the variance estimators were sufficiently ranked. We used ANOVA to test for sponsorship status differences in age, disposable income, and AUDIT-C scores, and Pearson's Chi Squared tests for gender differences in the proportion of participants classified as hazardous drinkers.

Results

A sample of 2048 sportspeople (892 females, 44%) was recruited (response rate 83%). Table 2 presents the characteristics of respondents by sponsorship status. Eighty-one participants (4%) were abstainers. The proportion with hazardous drinking (AUDIT score ≥ 8) was 84%.

Sponsorship

Thirty-six per cent ($n = 575$) of participants were in receipt of alcohol industry sponsorship. It should be noted that 338 participants (19%) did not answer questions about disposable income,

reducing the number available for analysis to 1658. We found no association between whether participants indicated their disposable income and AUDIT-C scores (mean difference -0.04 points, 95% CI -0.34 to 0.26) or sponsorship seeking (difference -0.0003 in proportions, 95% CI -0.14 to 0.14). Participants who did not answer these questions were less likely to have received sponsorship (difference -0.24 in proportions 95% CI -0.28 to -0.19). The initial examination of the association between alcohol consumption and each of the modelled variables (Table 3) shows that all covariates were significantly associated with alcohol consumption patterns. After adjusting for these covariates, the final model (Table 3) indicates that sportspeople who received alcohol industry sponsorship had significantly higher alcohol consumption than sportspeople who did not ($\chi^2_{(1643)} = 431.17, P < .0001$). In particular, those sponsored at the level of *team, club, team and club*, or with *combinations of personal and team or club sponsorship* had higher alcohol consumption scores than sportspeople not sponsored by the alcohol industry.

Table 4 displays results for bivariate and multivariable logistic regression models examining the associations between explanatory variables and the hazardous drinker outcome. All covariates except for gender were associated with hazardous drinking. The overall association between sponsorship type and hazardous drinking was statistically significant ($\chi^2_{(4)} = 35.9, p < 0.001$). The risk of hazardous drinking was associated with *club* sponsorship and with *team and club* sponsorship. After adjusting for the other variables in Table 4 the association between receiving any sponsorship and being a hazardous drinker was statistically significant ($\chi^2_{(4)} = 11.93, P = 0.018$).

We also tested the hypothesis that heavier drinking sportspeople may preferentially seek out alcohol industry sponsorship. After adjusting for all other variables in regression models, those seeking out sponsorship did not have significantly greater alcohol consumption than those who were approached by an alcohol industry sponsor ($\beta_{Adj} 0.80, 95\% \text{ CI: } 0.52\text{-}1.22$). Similarly, seeking sponsorship was not significantly associated with being a hazardous drinker (aOR= 1.19, 95% CI: 0.62-2.27). Finally, we explored interaction effects between gender and type of sponsorship for each of the three multivariable models, finding that none was significant (p-values > 0.17).

Discussion

There had been no previous research examining the association between alcohol industry sponsorship and drinking among sports participants in Europe. After controlling for several confounders (age, gender, disposable income, location) receipt of alcohol industry sponsorship at any level (individual, team, club, or combinations of these) was associated with higher levels of alcohol consumption. Sportspeople in receipt of alcohol industry sponsorship had greater odds of being hazardous drinkers. The results are in line with findings from both Australia and New Zealand, which showed an association between receipt of alcohol sponsorship and hazardous drinking in university [29] and community [28] sporting samples. The results of this study build upon previous research by accounting for disposable income which is strongly associated with alcohol consumption in young people [37]; and by studying a greater number and wider distribution of regions than in previous studies.

We also tested whether the association might reflect heavier drinkers seeking out alcohol industry sponsorship, a possibility raised by alcohol industry bodies as an alternative explanation for findings of previous research [38,39]. After controlling for several known confounders, there was no significant difference in drinking outcomes for those who reported seeking alcohol sponsorship versus those being offered sponsorship by the alcohol industry. Although this does rule out the possibility of confounding by unidentified variables, it does suggest that the association between alcohol industry sponsorship and problematic drinking is unlikely to be due to the latter causing the former.

The primary limitation of the study is the cross-sectional design, which precludes causal attribution. The statistical control of confounders and the test of sponsorship seeking behaviour reduces, but does not eliminate, the possibility that other factors may account for the association between alcohol sponsorship and drinking outcomes. Furthermore, by treating geographical location of respondents as a fixed effect in the analysis the findings may not generalise to other UK locations.

Relatedly, because the sample was university students involved in sport, it is uncertain as to whether the associations would generalise to non-students. Notably, however, research in a New Zealand community sample had a similar pattern of results as found here [26].

The measures of exposure are simple statements rather than validated scales. The development of full scales with established reliability and validity may be warranted in future, however, the questions have high face validity and our pilot research preceding previous studies using these items shows that respondents understand the questions and can answer them reliably [28].

While consistent with findings from other countries [40-42], the high level of problematic drinking in UK university sportspeople is noteworthy regardless of sponsorship status. The AUDIT total score (mean=14.4) and proportion of hazardous drinkers (84%) in this population group were high compared with estimates of these parameters in a recent study in English university students (mean=9.9, and 61%, respectively) [42]. Similarly, total AUDIT scores were slightly higher than those seen in previous studies of sportspeople in New Zealand (mean=12.5) [8], however, it should be noted that the absence of a sampling frame and non-random selection makes comparison problematic. Interestingly, differences in drinking outcomes between men and women were small (see Table 2) which suggests that the sporting culture may be particularly detrimental to women's drinking.

There is vigorous debate in several countries over the need for bans on alcohol advertising and sponsorship, with calls for more evidence to inform these debates [22-26]. The present study provides some evidence from the UK, showing that alcohol industry sponsorship is possibly harmful. Taken together with recent longitudinal research showing that indirect exposure to alcohol sponsorship in sport is associated with the later development of drinking expectancies and behaviour [30], and the high levels of hazardous drinking in sport, health policy makers and sports administrators should consider whether the harms outweigh the financial benefits of alcohol sponsorship.

The tobacco industry has been prohibited from advertising during sports broadcasts and from sponsoring sport in many countries and there is no evidence to suggest that this has resulted in a decline in sport participation or performance. Similar action has been called for in regard to the alcohol industry [27], with emphasis on the application of a precautionary principle, in particular, on shifting the burden of proof to the proponents of the potentially harmful activity, i.e., the alcohol industry. Secondly, in absence of strong evidence as to whether the association is likely to be causal which may take many years to develop, public health authorities are compelled to take preventive action until such evidence is available [27].

An objection raised by some parties, including sporting organisations, is that industry funds are crucial for the survival of sporting activity. In the same way that hypothecated tobacco taxes have been used in some countries to fund health programs and elite and community-level sport, tax revenues from alcohol could be employed to fund sport. There would be value in further examining the association between alcohol industry sponsorship and later alcohol use in sports participants with prospective measurement of exposure, of investigating mechanisms by which this exposure increases the risk of hazardous drinking, and effects on women involved in sport.

Acknowledgements

The research was supported by a grant from Alcohol Research UK. KOB was also supported by funding from the Australian Research Council, VicHealth, Australian National Preventive Health Agency, and the Australian Drug Foundation. KK's contribution to the research was funded by a Senior Research Fellowship from the National Health & Medical Research Council.

References

1. Rehm J., Mathers C., Povova S., Thavorncharoensap M., Teerawattananon Y., Patra J. Global burden of disease and injury and economic cost attributable to alcohol use and alcohol-use disorders. *Lancet* 2009; **373**: 2223-33.
2. Nutt D. J., King L. A., Phillips L. D. Drug harms in the UK: a multicriteria decision analysis. *Lancet* 2010; **376**: 1558-1565.
3. Strategy Unit. Alcohol Harm Reduction Strategy for England. London: Prime Minister's Strategy Unit, UK, 2004.
4. Gill J. S. Reported levels of alcohol consumption and binge drinking within the UK undergraduate student population over the last 25 years. *Alcohol Alcohol* 2002; **37**: 109–120.
5. Kuo M., Adlaf E. M., Lee H. *et al.* More Canadian students drink but American students drink more: comparing college alcohol use in two countries. *Addiction* 2002; **97**: 1583-1592.
6. Nelson T.F., Wechsler H. Alcohol and college athletes. *Med Sci Sports Exerc* 2001; **33**: 43-47.
7. Martens M. P., Dams-O'Connor K., Beck N.C. A systematic review of college student-athlete drinking: Prevalence rates, sport-related explanations, and interventions. *J Substance Abuse Treatment* 2006; **31**: 305-316.
8. O'Brien K.S., Blackie J., Hunter J.A. Hazardous drinking in elite New Zealand sportspeople. *Alcohol Alcohol* 2005; **40**: 239-41.
9. O'Brien K.S., Kolt G., Webber A., Hunter J.A. Alcohol consumption in sport: The influence of sporting idols, friends and normative drinking practices. *Drug Alcohol Rev* 2010; **29**: 676-683.
10. Wechsler H., Davenport A.E., Dowdall G.W., Grossman S.J., Zanakos S.I. Binge drinking, tobacco, and illicit drug use and involvement in college athletics. *J Am Col Health* 1997; **45**: 195-200.
11. Leichliter J.S., Meilman P.W., Presley C.A., Cashin J.R. Alcohol use and related consequences among students with varying levels of involvement with college athletics. *J Am Col Health* 1998; **46**, 257-262.

12. Partington S., Partington E., Heather N., et al. The relationship between membership of a university sports group and drinking behaviour among students at English Universities. *Addiction Res Theory* 2013; **24**: 339–347.
13. O'Brien K.S., Kolt G., Webber A., Hunter J.A. Alcohol consumption in sport: The influence of sporting idols, friends and normative drinking practices. *Drug Alc Review* 2010; **29**: 676-683.
14. Collins T., Vamplew W. *Mud, Sweat and Beers: A Cultural History of Sport and Alcohol*. New York: Berg; 2002.
15. Center on Alcohol Marketing and Youth (2004). *Alcohol Advertising on Sports Television 2001 to 2003*. <http://camy.org/factsheets/pdf/AlcoholAdvertisingSportsTelevision2001-2003.pdf>
16. Lefton T. (31 August 2009). A-B shifts approach to sports ties. <http://www.sportsbusinessjournal.com/article/63394>
17. Babor T., Caetano R., Casswell S., Edwards G., Giesbrecht N., Graham K. et al. *Alcohol: No Ordinary Commodity*, 2nd ed. London: Oxford University Press; 2010.
18. Casswell S., Thamarangsi T. Reducing harm from alcohol: call to action. *Lancet* 2009; **373**: 2247–57.
19. Anderson P., Chisholm D., Fuhr D.C. Effectiveness and cost-effectiveness of policies and programmes to reduce the harm caused by alcohol. *Lancet* 2009; **373**: 2234–46.
20. Anderson P., De Bruijn A., Angus K., Gordon R., Hastings G. Impact of alcohol advertising and media exposure on adolescent alcohol use: a systematic review of longitudinal studies. *Alcohol Alcohol* 2009; **44**: 229–43.
21. Smith L.A., Foxcroft D.R. The effect of alcohol advertising, marketing and portrayal on drinking behaviour in young people: systematic review of prospective cohort studies. *BMC Pub Health* 2009; **9**: 51.
22. British Medical Association Board of Science. *Under the Influence: the Damaging Effect of Alcohol Marketing on Young People*. London: British Medical Association, 2009.

23. Australian Medical Association. Alcohol Marketing and Young People: Time for a new policy agenda. Kingston, ACT. Australian Medical Association, 2012.
24. Irish Medical Organisation. Submission to the Oireachtas Joint Committee on Health and Children on Alcohol Marketing. November 2011.
25. House of Commons Health Select Committee. Alcohol: First report of session 2009-10, Volume 1, London. The Stationary Office, 2010.
26. The Government Response to the Health Select Committee Report on Alcohol. London. The Stationary Office, 2010.
27. Kypri K., O'Brien K., Miller P. Time for precautionary action on alcohol industry funding of sporting bodies. *Addiction* 2009; **104**: 1949-1950.
28. O'Brien K.S., Kypri K. Alcohol industry sponsorship of sport and drinking levels in New Zealand sportspeople. *Addiction* 2008; **103**: 1961-1966.
29. O'Brien K.S., Miller PG, Kolt G, Martens M, Webber A. Alcohol industry and non-alcohol industry sponsorship of sportspeople and drinking. *Alcohol Alcohol* 2011; **46**: 210-213.
30. de Bruijn A., Tanghe J., Beccaria F., Buljaski M., Celata C., Gosselt, J. *et al.* Report on the impact of European alcohol marketing exposure on youth alcohol expectancies and youth drinking. Alcohol Measures for Public Health Research Alliance (AMPHORA), 2012.
31. Reilly J. (5 Sept 2010). Last orders on the cards for alcohol ads in sports: But most people oppose outright ban on drink firms' sponsorship. Sunday Independent.
<http://www.independent.ie/national-news/last-orders-on-the-cards-for-alcohol-ads-in-sports-2325328.html>
32. Ameratunga S.N., Norton R.N., MacMahon S.W., Smith G.S., Jackson R.T., Currie R., *et al.* A prospective cohort study of blood donors: methodological issues in the investigation of injuries and chronic diseases. *ANZJPH* 1998; **22**: 578-82.
33. Tabachnick, B., & Fidell, L. *Using multivariate statistics*. New York: Harper Jr Row; 2001.

34. Saunders J.B., Aasland O.G., Babor T.F., de la Fuente J.R., Grant M. Development of the Alcohol Use Disorders Identification Test (AUDIT): World Health Organisation Collaborative Project on Early Detection of Persons with Harmful Alcohol Consumption—II. *Addiction* 1993; **88**: 791-804.
35. Conigrave K.M., Saunders J.B., Reznik R.B. Predictive capacity of the AUDIT questionnaire for alcohol-related harm. *Addiction* 1995; **90**: 1479-1485.
36. Oakes J.M. The (mis)estimation of neighborhood effects: causal inference for a practicable social epidemiology. *Soc Sci Med* 2004; **58**: 1929-52.
37. Adams S., Blackburn M.L., Cotti C.D. Minimum wages and alcohol-related traffic fatalities among teens. *Rev Econ Statistics* 2012; **94**: 828-840.
38. Corcoran S (24th June 2013). European association against alcohol sponsorship ban in Ireland. <http://www.businessandleadership.com/marketing/item/41618-european-association/>
39. Glendinning M., (27 Nov 2008). Alcohol sponsorship all clear, says drinks group. Sport Business. <http://www.sportbusiness.com/britsport/168360/alcohol-sponsorship-all-clear-says-drinks-group>.
40. Martha C., Grélot L., Peretti-Watel P. Participants' sports characteristics related to heavy episodic drinking among French students. *Int J Drug Polic* 2009; **20**: 152–160.
41. Martens M.P., Dams-O'Connor K., Beck N.C. A systematic review of college student-athlete drinking: prevalence rates, sport-related explanations, and interventions. *J Substance Abuse Treatment* 2006; **31**: 305–16.
42. Heather N., Partington S., Partington E., et al. Alcohol use disorders and hazardous drinking among undergraduates at English universities. *Alcohol Alcohol* 2011; **46**: 270-277..

Table 1. Frequency and percentage of sports reported as being played by participants.

Sport	Frequency*	Percent
Football (Soccer)	452	18.49
Rugby (Union and League)	445	18.20
Hockey	207	8.47
Cricket	177	7.24
Netball	163	6.67
Basketball	124	5.07
Athletics	114	4.66
Lacrosse	95	3.89
Swimming	79	3.23
Tennis	52	2.13
Water polo	50	2.04
Badminton	49	2.00
Squash	45	1.84
American football	44	1.80
Volleyball	43	1.76
Martial arts	37	1.51
Rowing	35	1.43
Dance Sport	32	1.31
Body Building	32	1.31
Golf	28	1.15
Fencing	19	0.78
Gymnastics	17	0.70
Skiing	16	0.65
Weight Lifting	15	0.61
Trampoline	13	0.53
Equestrian	11	0.45
Cycling	11	0.45
Boxing	9	0.37
Cheer Leading	9	0.37
Table Tennis	7	0.29
Platform Diving	5	0.20
Sailing	4	0.16
Softball	3	0.12
Ice Skating	1	0.04
Lawn Bowls	1	0.04
Australian Football	1	0.04

* Note: the total frequency of sports played is 2445. This is due to some participants listing participation in multiple sports.

Table 2. Characteristics of the sample. Numbers other than mean and standard deviation scores are represented as raw counts and percentages (%). Significant sponsorship differences are indicated for age, disposable income, and all drinking behaviour scores.

Characteristic	No sponsorship n = 1473 (72)	Sponsorship n = 575 (28)	Total n = 2048
Sex **			
Female	614 (69)	278 (31)	892 (44)
Male	858 (74)	297 (26)	1155 (56)
Mean age **	20.05 (1.87)	19.77 (1.50)	19.97 (1.78)
Mean disposable income £	66.87 (83.42)	56.78 (52.90)	63.58 (75.00)
Mean alcohol consumption score ***	7.86 (2.81)	8.58 (2.39)	8.05 (2.72)
Mean AUDIT total score ***	13.97 (7.15)	15.74 (6.90)	14.47 (7.12)
Hazardous drinker (AUDIT 8+) ***	1212 (70)	512 (30)	1724 (84)
Alcohol dependence (AUDIT 16+) ***	568 (66)	287 (34)	855 (42)
Location ***			
North West	640 (63)	374 (37)	1014 (50)
Midlands	200 (74)	72 (26)	272 (13)
London	201 (85)	36 (15)	237 (11)
Southern England	432 (82)	93 (18)	525 (26)
Total	1473 (72)	575 (28)	2048 (100)
Level of sponsorship			
None	1473 (72)		
Team only		214 (10)	
Club only		195 (10)	
Team and Club only		144 (7)	
Personal combinations		22 (1)	

*Significant sponsorship difference at the $P < 0.05$ level, **Significant at the $P < 0.01$ level. ***Significant at the $P < 0.001$ level. Total (%) are with respect to the whole sample.

Table 3. Bivariate and multivariable regression analysis of alcohol consumption score after accounting for covariates.

Parameters ^a	Alcohol Consumption			
	Unadjusted β (95% CI)		Adjusted β^b (95% CI)	
Age	-0.16	(-0.22, -0.09)***	-0.15	(-0.22, -0.08)****
Male (Reference: Females)	0.39	(0.19, 0.96)**	0.38	(-0.13, 0.89)
Weekly disposable income (Reference: £0-20)				
£21-40	0.73	(0.09, 1.37)**	0.74	(0.23, 1.25)**
£41-60	1.09	(0.67, 1.53)***	1.09	(0.71, 1.46)****
£61-80	1.20	(0.76, 1.64)***	1.12	(0.80, 1.45)****
£81-100	1.11	(0.54, 1.68)**	1.14	(0.67, 1.60)****
>£100	0.60	(0.13, 1.08)*	0.63	(0.18, 1.09)**
Location (Reference: London)				
North West	1.52	(0.82, 2.22)***	1.24	(0.48, 2.00)**
Midlands	0.49	(-0.22, 1.20)	0.37	(-0.29, 1.02)
Southern England	1.22	(0.51, 1.94)***	1.12	(0.42, 1.83)**
Sponsorship (Reference: No alcohol sponsorship=0)				
Team	0.63	(0.27, 0.98)**	0.41	(0.07, 0.75)*
Club	1.00	(0.44, 1.56)***	0.73	(0.12, 1.34)*
Team and club	0.77	(0.40, 1.15)***	0.79	(0.28, 1.29)**
Personal combinations	1.28	(0.42, 2.14)*	1.27	(0.42, 2.11)**

*Significant at the $P < 0.05$ level, **Significant at the $P < 0.01$ level. ***Significant at the $P < 0.001$ level.

^aLevel of sponsorship categories are mutually exclusive, and are limited to those with sufficient numbers to permit valid statistical inference. AUDIT: Alcohol Use Disorders Identification Test; CI: confidence interval. ^b Linktest suggests that the overall model was correctly specified ($t=-0.06$; $p>0.45$); testing for multicollinearity was not possible due to bootstrapping the model.

Table 4. Bivariate and multivariable logistic regression analysis for hazardous drinking after accounting for covariates.

Parameters ^a	Hazardous drinking (AUDIT Total 8+)	
	OR (95% CI)	Adjusted OR ^{b,c} (95% CI)
Age	0.92 (0.87, 0.97)**	0.92 (0.88, 0.97)**
Male (Females as reference)	1.29 (0.77, 2.15)	1.26 (0.78, 2.01)
Weekly disposable income (Reference: £0-20)		
£21-40	1.92 (1.18, 3.14)**	2.03 (1.33, 3.10)**
£41-60	2.31 (1.57, 3.37)****	2.40 (1.66, 3.45)****
£61-80	3.03 (1.62, 5.68)***	2.96 (1.71, 5.11)***
£81-100	2.12 (1.30, 3.46)**	2.28 (1.38, 3.76)**
>£100	1.67 (1.14, 2.42)**	1.70 (1.13, 2.58)*
Location (Reference: London=0)		
North West	3.28 (1.93, 5.59)****	2.78 (1.73, 4.45)****
Midlands	1.52 (0.93, 2.50)	1.41 (0.91, 2.20)
Southern England	2.60 (1.66, 4.06)****	2.55 (1.65, 3.95)****
Sponsorship (Reference: No alcohol sponsorship=0)		
Team	1.50 (0.97, 2.30)	1.25 (0.82, 1.91)
Club	2.21 (1.14, 4.28)*	1.78 (0.87, 3.63)
Team and club	1.94 (1.18, 3.17)**	2.04 (1.04, 4.00)*
Personal combinations	4.30 (1.42, 13.01)**	4.12 (1.29, 13.15)*

*Significant at the $P < 0.05$ level. **Significant at the $P < 0.01$ level. ***Significant at the $P < 0.001$ level. ^aLevel of sponsorship categories are mutually exclusive, and are limited to those with sufficient numbers to permit valid statistical inference. AUDIT: Alcohol Use Disorders Identification Test; CI: confidence interval. ^bHosmer-Lemeshow tests also indicated a good fit for models (hazardous drinking, $\chi^2=5.29$, $p>.73$; and alcohol dependence, $\chi^2=15.22$ $p=.056$). ^cLinktest suggests that the overall model was correctly specified (hazardous drinking $t=-0.03$; $p>0.82$; alcohol dependence $t=0.08$; $p>0.55$).