



University of
Salford
MANCHESTER

Dissipative solitons of a spatiotemporal Ginzburg-Landau equation

Cooper, AR, Christian, JM, Bresnahan, DW, McDonald, GS and
Chamorro-Posada, P

Title	Dissipative solitons of a spatiotemporal Ginzburg-Landau equation
Authors	Cooper, AR, Christian, JM, Bresnahan, DW, McDonald, GS and Chamorro-Posada, P
Type	Conference or Workshop Item
URL	This version is available at: http://usir.salford.ac.uk/id/eprint/32869/
Published Date	2014

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: usir@salford.ac.uk.

Dissipative solitons of a spatiotemporal Ginzburg-Landau equation

A. R. Cooper¹, J. M. Christian¹, D. W. Bresnahan¹, G. S. McDonald¹,
and P. Chamorro-Posada²

¹ *Materials & Physics Research Centre, University of Salford,
Greater Manchester M5 4WT, U.K.*

² *ETSI Telecomunicación, Universidad de Valladolid,
Campus Miguel Delibes Paseo Belén 15, E-47011 Valladolid, Spain*

Complex Ginzburg-Landau (GL) equations play a fundamental role in our understanding of universal wave propagation in generic systems with dispersion, nonlinearity, gain, and loss. Over three decades ago, stationary bright-type solitons were derived by Pereira and Stenflo (PS solitons) for a cubically-nonlinear GL equation [Phys. Fluids, **20**, 1733 (1977)]. Dark-type solutions followed a short time later, introduced by Nozaki and Bekki (NB solitons) [J. Phys. Soc. Jpn. **53**, 5365 (1984)]. In photonics, such self-localizing optical wavepackets can typically arise when group-velocity dispersion is balanced by self-phase modulation, and nonlinear losses (e.g., from two-photon absorption) are compensated by linear gain (e.g., from doping in the host medium). In the corresponding GL model, linear gain tends to introduce instability (growth of infinitesimally-small perturbations) such that all its solutions are rendered unstable in the long term.

In this presentation, we introduce a spatiotemporal generalization of the classic GL equation. Our approach is based on a fully-second order scalar governing equation that omits the traditional prescription based on: (i) assumption of slowly-varying envelopes, and (ii) a Galilean boost to the local time frame (usually introduced for mathematical convenience). As a consequence, our model naturally describes wave phenomena from the laboratory reference frame. Moreover, the velocity combination rule that emerges is reminiscent of that which appears in Einstein's relativistic mechanics [Christian *et al.*, Phys. Rev. Lett. **108**, 034101 (2012)]. Exact bright and dark dissipative solitons have been derived, and their space-time geometry explored in detail. Crucially, the classic PS and NB solutions emerge asymptotically from their spatiotemporal generalizations in a simultaneous multiple limit. Of particular importance is that the inherent instability of the zero-amplitude solution, which plagues waves of the cubically-nonlinear GL equation, persists in the spatiotemporal case. However, we have taken the first steps toward stabilizing our model and preliminary results will be reported.