



University of  
**Salford**  
MANCHESTER

# Influence of wheel size on muscle activity and tri-axial accelerations during cross-country mountain biking

Hurst, HT, Atkins, SJ, Metcalfe, J, Rylands, L and Sinclair, JK

<http://dx.doi.org/10.1080/02640414.2016.1215497>

<b>Title</b>	Influence of wheel size on muscle activity and tri-axial accelerations during cross-country mountain biking
<b>Authors</b>	Hurst, HT, Atkins, SJ, Metcalfe, J, Rylands, L and Sinclair, JK
<b>Type</b>	Article
<b>URL</b>	This version is available at: <a href="http://usir.salford.ac.uk/id/eprint/40051/">http://usir.salford.ac.uk/id/eprint/40051/</a>
<b>Published Date</b>	2016

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: [usir@salford.ac.uk](mailto:usir@salford.ac.uk).



**Influence of wheel size on muscle activity and tri-axial accelerations during Cross-Country mountain biking**

Journal:	<i>Journal of Sports Sciences</i>
Manuscript ID	RJSP-2014-1186.R2
Manuscript Type:	Special Issue
Keywords:	Vibration, Cycling, Mountain biking, electromyography

SCHOLARONE™  
Manuscripts

Pre-Review Only

1  
2  
3 1 **Influence of wheel size on muscle activity and tri-axial accelerations during Cross-**  
4  
5 2 **Country mountain biking**

6  
7  
8 3

9  
10 4 **Running head:** Mountain biking and muscle activity

11  
12  
13 5 **Keywords:** vibrations, cycling, mountain biking, electromyography

14  
15  
16 6 **Word Count:** 3989

17  
18  
19 7

20  
21  
22 8

23 9 **Acknowledgments**

24  
25  
26 10 We would like to acknowledge Santa Cruz Bicycles in the USA and UK for the loan of the  
27  
28 11 test equipment and their support in this project. We would also like to confirm there were no  
29  
30 12 conflicts of interest and that Santa Cruz Bicycles' involvement was completely impartial. We  
31  
32 13 would also like to thank the participants for their time and effort in taking part in the study.  
33  
34

35  
36 14

37  
38 15

1  
2  
3 **1 Abstract**  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

2 This study aimed to investigate the influence of different mountain bike wheel diameters on  
3 muscle activity and **whether larger diameter wheels attenuate** muscle vibrations during cross-  
4 country riding. Nine male competitive mountain bikers (age  $34.7 \pm 10.7$  yrs; stature  $177.7 \pm$   
5  $5.6$  cm; body mass  $73.2 \pm 8.6$  kg) participated in the study. Riders performed one lap at race  
6 pace on 26", 27.5" and 29" wheeled mountain bikes. sEMG and acceleration (RMS) were  
7 recorded for the full lap and during ascent and descent phases at the gastrocnemius, vastus  
8 lateralis, biceps brachii and triceps brachii. No significant main effects were found by wheel  
9 size for each of the four muscle groups for sEMG or acceleration during the full lap and for  
10 ascent and descent ( $p > .05$ ). When data were analysed between muscle groups, significant  
11 differences were found between biceps brachii and triceps brachii ( $p < .05$ ) for all wheel sizes  
12 and all phases of the lap with the exception of for the 26" wheel during the descent. Findings  
13 suggest wheel diameter has no influence on muscle activity and vibration during mountain  
14 biking. However, more activity was observed in the biceps brachii during 26" wheel  
15 descending. This is possibly due to an increased need to manoeuvre the front wheel over  
16 obstacles.  
17  
18  
19  
20

## 1 Introduction

2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The physiological demands of cross-country mountain biking have been well reported over recent years (MacRae, Hise, & Allen, 2000; Lee, Martin, Anson, Grundy, & Hahn, 2002; Stapelfeldt, Schwirtz, Schumacher, & Hillebrecht, 2004; Impellizzeri, Rampinini, Sassi, Mognoni, & Marcora, 2005; Prins, Treblanche, & Myburgh, 2007). These studies focused primarily on aerobic and anaerobic contributions to the activity. However, few studies have investigated muscle activity during mountain biking. Hurst and Atkins (2006) proposed mountain bike riders utilised a high level of isometric muscular contractions during racing to aid postural control and control of the bicycle over obstacles.

12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Surface electromyography (sEMG) has been used to analyse muscle activity in road cyclists (Egaña, Ryan, Warmington, & Green, 2010; Matsuura, Arimitsu, Yuncki, & Yano, 2011; Blake, Champoux, & Wakeling, 2012). However, these studies focused on lower limb muscle activity and were laboratory based. Duc, Bertucci, Pernin, and Grappe (2008) investigated the influence of hand grip position on the handlebar drops compared to the tops, during uphill cycling on upper body muscle activity. However, the activity observed does not reflect those used in mountain biking, due to the more dynamic nature of upper body movements. Hurst et al. (2012) investigated the influence of course terrain on upper body sEMG during downhill riding in elite downhill and cross-country cyclist. They found significant differences in the muscles recruited between the groups, with downhill riders exhibiting greatest triceps brachii activity, whilst cross-country riders showed greater activity in the brachioradialis. Their study concluded that the differences in bicycle design between downhill and cross-country bicycles may in part have influenced the differences observed.

1  
2  
3 1  
4  
5 2 The standard mountain bike wheel diameter has long been 26". However, in the past 10 years  
6  
7 3 two more wheel size standards have been introduced, 29" and more recently 27.5". The  
8  
9 4 rationale for 29" wheels has been that the larger diameter improves roll over small trail  
10  
11 5 bumps and therefore improves contact with the ground and subsequent velocity. In addition,  
12  
13 6 it has been proposed by the cycling industry that the larger wheel will also reduce trail  
14  
15 7 vibrations observed in mountain biking (Levy & Smith, 2005; Faiss, Praz, Meichtry, Gobelet,  
16  
17 8 & Deriaz, 2007) being transmitted to the rider. Mester, Spitzenfeil, Schwarzer, & Seifriz  
18  
19 9 (1999) found that these vibrations lead to increased muscle activity to aid dampening of trail  
20  
21 10 shocks, resulting in reduced exercise efficiency. Justification for the newer 27.5" wheel  
22  
23 11 standard has been that it potentially provides a balance between the better handling  
24  
25 12 characteristics of the 26" wheel and the improved rolling properties of 29" wheels. However,  
26  
27 13 few academic studies exist determining the merits of these different wheel diameters on  
28  
29 14 performance. Macdermid, Fink, and Stannard (2014) did investigate differences in 3D  
30  
31 15 accelerations between 26" and 29" wheels and reported that 29" wheeled mountain bikes  
32  
33 16 resulted in significantly greater vibrations than the 26" wheel during a typical lap of a cross-  
34  
35 17 country course, possibly due to the increase in velocity observed. However, Macdermid and  
36  
37 18 Stannard (2014) also point out that a key limitation of their study was that they used the same  
38  
39 19 29" bicycle fitted with 26" wheels, thus potentially influencing their findings, as frame  
40  
41 20 geometry may not have been optimised for the smaller wheel size. In addition, they did not  
42  
43 21 investigate the influence of 27.5" wheels. Therefore, the aims of this study were to  
44  
45 22 investigate the influence of the 3 primary wheel standards on tri-axial accelerations and  
46  
47 23 muscle activity in upper and lower limbs during cross-country mountain biking. It was  
48  
49 24 hypothesised that the larger wheel diameters would significantly attenuate the muscle activity  
50  
51 25 and vibrations experienced by the riders more than the 26" wheel.  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 1  
4  
5  
6 2**Materials and methods**7 3 **Participants**

4 Nine male competitive mountain bikers (age  $34.7 \pm 10.7$  yrs; stature  $177.7 \pm 5.6$  cm; body  
5 mass  $73.2 \pm 8.6$  kg) participated in the study. All riders had a minimum of 5 years racing  
6 experience and competed at a National level in their respective age category. Ethical approval  
7 was granted by the University of Central Lancashire Ethics Committee and in accordance  
8 with the Declaration of Helsinki. Participants were informed both verbally and in writing of  
9 the test procedures and written informed consent was obtained.

10  
1111 **Course Profile**

12 All testing was completed on a purpose build cross-country mountain bike course at the  
13 British National Cycle Centre (Clayton Vale, Manchester). The course comprised of  
14 undulating technical sections and also featured one major climb and descent, for which data  
15 were analysed for these sections as well as the full lap. Figure 1 shows a global positioning  
16 system (GPS) trace of the course and profile. A Garmin Edge 810 GPS computer was used to  
17 record lap times and temperature. However, due to the level of tree cover and the  
18 inaccuracies of GPS under tree cover, actual distances were recorded with a calibrated trundle  
19 wheel. Distances of each section highlighted in figure 1 were; Start to 1 = 1.72 km; 1 to 2 =  
20 0.38 km (Climb); 2 to 3 = 0.47 km; 3 to 4 = 0.66 km (Descent); and 4 to Finish = 0.25 km;  
21 total lap distance = 3.48 km. Based on Ordnance Survey maps, the average gradient of the  
22 climb was  $5.8 \pm 0.3$  %, whilst the descent gradient was  $-6.1 \pm 0.4$  %. Testing took place over  
23 a four week period between the months of June and July. Mean temperature was  $18.5 \pm 1.5$   
24 °C, with all test sessions being performed in sunny dry conditions. Therefore, course  
25 conditions were consistent for all trials.

1

2 \*\*\*Figure 1 near here\*\*\*

3

4 **Surface EMG and Acceleration Analyses**

5 Surface electromyography (sEMG) data were recorded using a mobile wireless sensor system  
6 (Trigno Mobile, Delsys, USA) at 1926 Hz from the gastrocnemius, vastus lateralis, biceps  
7 brachii and triceps brachii. Parallel bar electrodes with a 1 cm spacing between electrodes  
8 were used. This method has been shown to significantly reduce the cross-talk between  
9 muscles when compared to the use of disc electrodes (De Luca, Kuznetsov, Gilmore, & Roy,  
10 2012). All recordings were taken from the left hand side of the body. Prior to sensor  
11 placement, the skin was prepared by shaving the area, lightly abrading and cleaning with  
12 alcohol wipes in order to minimise skin impedance. **Though impedance was not measured, it  
13 is accepted that this may have changed during testing due to sweating by the participants.  
14 However, in order to ensure the same muscles were assessed during each trial, the sEMG  
15 sensors were not removed between laps.** Sensors were positioned longitudinally in parallel to  
16 the muscle fibres on the medial aspect of each muscle. Placement of the sensors was in  
17 accordance with the Surface EMG for Non-Invasive Assessment of Muscles project  
18 recommendations. Once positioned on the muscles, sensors were secured in place using  
19 elasticated bandages. Though the use of elasticated bandages may have possibly reduced  
20 tissue movement artefacts, this would potentially improve indirect measurement of  
21 acceleration on the skeletal system. In addition, during pilot testing, sensors did not stay in  
22 situ using just hypoallergenic tape as a result of increased sweating. As such bandaging was  
23 deemed the most appropriate method for securing the sensors. Following data collection,  
24 sEMG data were full-wave rectified then filtered at 20 Hz using a second order low pass  
25 Butterworth filter.



1

2  
3  
4  
5  
6 According to Sinclair, Brooks, Edmundson, & Hobbs (2012), due to the nature of field-based  
7  
8 testing, normalisation of sEMG signals to a maximal voluntary isometric contraction are not  
9  
10 possible or appropriate, though their study looked at running. This was due to the dynamic  
11  
12 muscular activity involved. Therefore, they proposed that sEMG data should be normalised to  
13  
14 a dynamic peak task, that being the peak amplitude recorded during the field-based trials. As  
15  
16 cycling is also dynamic in nature, sEMG data in the present study are presented as a  
17  
18 percentage of the dynamic peak task for the full lap and the ascent and descent sections.  
19  
20  
21  
22

9

23  
24  
25 In addition to sEMG, the Delsys Trigno sensors also recorded wireless tri-axial acceleration  
26  
27 for measurement of muscle vibrations. The placement of accelerometers on soft tissue to  
28  
29 monitor vibrations has previously been validated (Lafortune et al. 1995; Coza et al 2010).  
30  
31 Nokes, Fairclough, Mintowt, Mackie, and Williams (1984) also reported that vibrations could  
32  
33 be reliably and validly recorded with muscle mounted accelerometers if they were preloaded  
34  
35 to overcome the influence of soft tissue properties. Given the use of elasticated bandages to  
36  
37 secure the sensors, an element of preload would have been applied to the accelerometers,  
38  
39 though the exact force could not be determined. In addition, Ziegert and Lewis (1979) found  
40  
41 that lower mass accelerometers had a minimal effect data acquisition. Sensors used in the  
42  
43 current study were 14 g in mass. Data were sampled at 148.1 Hz with a range of  $\pm 9$  g, and  
44  
45 analysed for total (XYZ) accelerations. Root mean squares (RMS) of the resultant XYZ data  
46  
47 were determined to analyse the vibrations experienced by the riders (ISO 1997). Similar  
48  
49 methods have been used in earlier studies of mountain biking vibrations (Macdermind et al.,  
50  
51 2014). Data were processed using EMGWorks Version 4.0 (Delsys Inc., Boston, USA) to  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 1 determine mean sEMG and RMS for the full lap and the ascent and descent phases prior to  
4  
5 2 later statistical analyses.  
6  
7

### 8 **3 Protocols**

9  
10 4 Following placement of the sensors participants were shown the course and allowed 1 hour to  
11  
12 5 familiarise themselves with the route. With the exception of one rider, all had previous  
13  
14 6 experience of racing or riding on the course used. Participants were required to complete one  
15  
16 7 lap of the course at self-determined race pace on each of the three wheel sizes. Thirty minutes  
17  
18 8 passive rest was given between trials to allow adequate recovery, this was then followed by a  
19  
20 9 10 minute re-warm, which consistent of low intensity cycling prior to the next trial. The test  
21  
22 10 bicycles were Santa Cruz Superlight full suspension mountain bikes (Santa Cruz Bicycles,  
23  
24 11 USA) with 120 mm and 100 mm of front and rear suspension respectively. All bicycles were  
25  
26 12 2014 models and were new and unused prior to the study. All three bicycles were fitted with  
27  
28 13 identical components, again new and unused, and differed component wise only in wheel  
29  
30 14 size, those being 26", 27.5" and 29" diameter. Tyre pressure was 35 psi for all trials.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

16 Frame size was a medium for all three bicycles. However, overall bicycle mass differed due  
17 to the differences in wheel diameter and wheel mass (13.69 kg, 13.93 kg and 14.15 kg for the  
18 26", 27.5" and 29" wheeled bicycles respectively). Therefore, to account for this and to  
19 eliminate any potential influence on results, bicycle mass was standardised to the heaviest  
20 bicycle (29" wheeled) by adding small weights to the lower downtube of the 26" and 27.5"  
21 wheeled bicycles. **Subsequently, any observed differences could be attributed more to rolling  
22 properties of the different wheel diameters than any differences in mass.** Top tube lengths  
23 were 590, 602 and 613 mm for the 26", 27.5" and 29" wheels respectively, whilst bottom  
24 bracket heights were 319, 326 and 337 mm, respectively. To accommodate differences in

1  
2  
3 1 rider stature a choice of two different stem lengths were offered (90 and 110 mm). Saddle  
4  
5 2 setback was also adjusted to ensure best fit. In addition, riders were allowed to use their own  
6  
7 3 pedals. Suspension systems were set up according to the manufacturers' recommendations for  
8  
9 4 each rider's individual weight and to allow for 10 percent sag in the travel. Both front and  
10  
11 5 rear suspension were run in fully open mode, whilst shock leverage ratios were optimised by  
12  
13 6 the manufacturer for each frame geometry.  
14  
15  
16  
17  
18  
19

20 8 Immediately prior to each trial participants lined up on the start line and were instructed to  
21  
22 9 perform a maximal vertical jump to provide a marker on the accelerometer data. This was  
23  
24 10 used to synchronise the accelerometer and sEMG data. Participants were then set off 10 s  
25  
26 11 post vertical jump. All sEMG and accelerometer data pre and 10 s post jump were discarded  
27  
28 12 from the analysis. The time the participants crossed the finish line was noted and all data after  
29  
30 13 this time was also discarded from the results. In the absence of a trigger system to help  
31  
32 14 identify where the climb and descents started and ended on the sEMG and accelerometer  
33  
34 15 traces, data were synchronised with the time date on the GPS unit attached to the handlebars.  
35  
36 16 The start and end of the main climb and descent were marked with signposts. Riders were  
37  
38 17 instructed to press the 'lap' button on the GPS at the start and end of these sections. This then  
39  
40 18 enabled the identification these sections on the sEMG and accelerometer traces by means of  
41  
42 19 the time stamps. When determining mean data in the EMGWorks software, these time blocks  
43  
44 20 were identified and subsequently analysed. Participants were randomly assigned to three  
45  
46 21 groups of three participants and trials were counterbalanced for each group.  
47  
48  
49  
50  
51  
52  
53  
54

## 55 23 **Statistical Analyses**

56  
57  
58  
59  
60

1 Normality of data were first confirmed by means of a Shapiro-Wilk test. Differences in  
2 muscle activity and accelerations between wheel sizes and between muscle groups within  
3 each wheel size were determined using within groups one-way repeated measures ANOVA's.  
4 Bonferroni corrections were used during post hoc analyses to control for type I errors. Where  
5 the homogeneity assumption was violated, the degrees of freedom were adjusted using the  
6 Greenhouse Geisser correction. Effect sizes were calculated using partial  $\eta^2$  ( $p^2$ ).  
7 Significance was accepted at the  $p \leq 0.05$  level (Sinclair, Taylor, & Hobbs, 2013) and data are  
8 presented as mean  $\pm$  standard deviation. All statistical procedures were conducted using  
9 SPSS 20.0 (SPSS Inc., Chicago, IL, USA).

## 11 Results

12 Though riders performed each lap at a self-determined race pace, no significant difference  
13 was found for mean velocity between wheel sizes ( $F_{(2,16)} = .70$ ;  $p = .45$ ;  $p^2 = .08$ ), with mean  
14 values of  $13.7 \pm .8$  (95% CI, 13.1, 14.3),  $13.6 \pm .8$  (95% CI, 13.0, 14.2) and  $13.9 \pm .8$  (95%  
15 CI, 13.3, 14.5)  $\text{km}\cdot\text{h}^{-1}$  being reported for the 26", 27.5" and 29" wheeled bicycles  
16 respectively. Additionally, no significant differences were found in lap times between wheel  
17 sizes ( $F_{(2,16)} = .69$ ;  $p = .45$ ;  $p^2 = .08$ ). Mean lap times were  $916.1 \pm 54.5$  (95% CI, 874.3,  
18  $957.6$ ),  $923.8 \pm 52.9$  (95% CI, 883.1, 964.5) and  $904.2 \pm 54.7$  (95% CI, 862.1, 946.3) s for  
19 the 26", 27.5" and 29" wheeled bicycles respectively.

## 21 Surface electromyography data

22 Analysis of sEMG data revealed no significant main effects for wheel size when analysed  
23 over the full lap for the gastrocnemius ( $F_{(2,16)} = .29$ ;  $p = .75$ ;  $p^2 = .04$ ), vastus lateralis ( $F_{(2,16)} =$

1 1.42;  $p = .27$ ;  $p^2 = .15$ ), biceps brachii ( $F_{(2,16)} = .83$ ;  $p = .40$ ;  $p^2 = .09$ ) or triceps brachii ( $F_{(2,16)}$   
2 = .12;  $p = .89$ ;  $p^2 = .02$ ). When data were analysed for the climb alone, no significant main  
3 effect was again reported for any muscle group by wheel size; gastrocnemius ( $F_{(2,16)} = .39$ ;  $p$   
4 = .69;  $p^2 = .05$ ), vastus lateralis ( $F_{(2,16)} = .95$ ;  $p = .41$ ;  $p^2 = .11$ ), biceps brachii ( $F_{(2,16)} = .44$ ;  $p$   
5 = .65;  $p^2 = .05$ ) and triceps brachii ( $F_{(2,16)} = .43$ ;  $p = .66$ ;  $p^2 = .05$ ) respectively. Analysis of  
6 descent sEMG also found no significant main effect for wheel size; gastrocnemius ( $F_{(2,16)} =$   
7 .23;  $p = .80$ ;  $p^2 = .03$ ), vastus lateralis ( $F_{(2,16)} = 1.64$ ;  $p = .23$ ;  $p^2 = .17$ ), biceps brachii ( $F_{(2,16)} =$   
8 .94;  $p = .37$ ;  $p^2 = .11$ ) or triceps brachii ( $F_{(2,16)} = .13$ ;  $p = .88$ ;  $p^2 = .02$ ). **Table 1** shows the  
9 mean values for sEMG as a percentage of dynamic peak task and 95% CI. When muscles  
10 were compared against each other within wheel sizes, significant differences ( $p < .05$ ) were  
11 found between the antagonistic muscles of the biceps brachii and triceps brachii within all  
12 wheel sizes when analysed over the full lap and the climb. However, during the descent  
13 section significant differences ( $p < .05$ ) were only reported between biceps brachii and triceps  
14 brachii within the 27.5" and 29" wheel sizes.

15  
16 \*\*\*Table 1 near here\*\*\*

### 18 Acceleration Data

19 Analysis of total XYZ acceleration amplitude (RMS) between muscles over the full lap  
20 revealed no significant main effect for wheel sizes in the gastrocnemius ( $F_{(2,16)} = 1.42$ ;  $p =$   
21 .27;  $p^2 = .15$ ), vastus lateralis ( $F_{(2,16)} = .48$ ;  $p = .63$ ;  $p^2 = .06$ ), biceps brachii ( $F_{(2,16)} = .85$ ;  $p =$   
22 .45;  $p^2 = .10$ ) or triceps brachii ( $F_{(2,16)} = .70$ ;  $p = .51$ ;  $p^2 = .08$ ). Data from the climbing section  
23 also showed no significant main effect for wheel size; gastrocnemius ( $F_{(2,16)} = .87$ ;  $p = .44$ ;  $p^2$

1 = .10), vastus lateralis ( $F_{(2,16)} = 2.73$ ;  $p = .10$ ;  $p^2 = .25$ ), biceps brachii ( $F_{(2,16)} = 1.50$ ;  $p = .25$ ;  
2  $p^2 = .16$ ) or triceps brachii ( $F_{(2,16)} = .90$ ;  $p = .43$ ;  $p^2 = .10$ ). Similarly, no significant main  
3 effects were found for wheel size during the main descent section of the lap; gastrocnemius  
4 ( $F_{(2,16)} = 1.94$ ;  $p = .18$ ;  $p^2 = .20$ ), vastus lateralis ( $F_{(2,16)} = .68$ ;  $p = .52$ ;  $p^2 = .08$ ), biceps brachii  
5 ( $F_{(2,16)} = .47$ ;  $p = .64$ ;  $p^2 = .06$ ) or triceps brachii ( $F_{(2,16)} = .51$ ;  $p = .61$ ;  $p^2 = .06$ ). **When**  
6 **muscles were compared against each other within each wheel size, significant differences**  
7 **existed ( $p < .001$ ) in total acceleration between the gastrocnemius and vastus lateralis as**  
8 **expected during the full lap, climb and descent sections. However, no significant differences**  
9 **in acceleration were found between the biceps brachii and triceps brachii for any section of**  
10 **the course or over the full lap. Table 2 shows the mean total acceleration amplitude (RMS)**  
11 **for each muscle and wheel size along with 95% CI.**

12  
13 **\*\*\*Table 2 near here\*\*\***

## 14 **Discussion**

15 The purpose of this study was to determine the influence of three different mountain bike  
16 wheel diameters on muscle activity and vibrations at different muscle sites during a typical  
17 cross-country lap at race pace. The key findings were that no significant differences were  
18 observed in muscle activity between the 26", 27.5" and 29" wheeled bicycles when analysed  
19 for the full lap and during the highlighted ascent and descent sections. Similarly, no  
20 significant differences were found in accelerations at the four different muscle sites between  
21 the three wheel sizes. Therefore, the hypothesis that larger wheels reduce muscle activity and  
22 vibrations should be rejected. However, other factors may also have influenced the results of  
23 the current study.

1  
2  
3 1  
4  
5  
6 2

The present study found significantly lower sEMG values than those reported by Hurst et al. (2012) in elite cross-country and downhill riders. Although the riders in the present study were not performing at an elite level, the differences between these studies are less likely the result of differences in athletic ability and more likely due to the limitation of normalising muscle activity during dynamic tasks to a maximal voluntary isometric contraction. Mountain biking involves numerous eccentric muscular contractions as a result of riding off drops to aid dampening and control of the bicycle. This most likely resulted in the greater than maximal voluntary isometric contraction values reported previously by Hurst et al. (2012). However, the present study normalised on course muscle activity to the peak dynamic task values elicited across all trials in accordance the methods previously reported (Sinclair et al., 2012). Results showed that irrespective of wheel size, mean muscle activity over the full lap, ascent and descent sections were not significantly different and represented a very low percentage of peak dynamic task, approximately 2-6 %. This low percentage may potentially be skewed somewhat due to high peak values occurring when riding off drops to help control the bicycle. The course used contained several drops of approximately 1 m in height. In addition, differences between the current study and that of Hurst et al. (2012) may also be attributed to the different sEMG methodologies used and the use of disc electrodes by Hurst et al. (2012), as opposed to the parallel bar electrodes used in the current study.

20

21 Though values did not reach a level of significance, the results show that biceps brachii  
22 activity was generally higher during all phases of the lap for the 26" wheel when compared to  
23 the 27.5" and 29" wheels. This may be the result of riders having to pull more on the handle  
24 bars increasing elbow flexion to lift the front wheel of the 26" wheel over obstacles and small

1  
2  
3 1 gaps. As a result of the larger wheel diameter, both the 27.5" and 29" wheels would  
4  
5 2 potentially roll over such obstacles more efficiently and would not drop into small gaps as  
6  
7 3 easily, thereby reducing muscle activity. This supposition would appear to be supported when  
8  
9  
10 4 data were analysed between muscle groups. Significant differences were found between the  
11  
12 5 antagonistic muscles of the biceps brachii and triceps brachii over the full lap and the main  
13  
14 6 climbing section for all wheel sizes. However, during the descent section, only significant  
15  
16 7 differences were found between these two muscle groups when riding the 27.5" and 29"  
17  
18 8 wheeled bicycles. The non-significant difference between biceps brachii and triceps brachii  
19  
20 9 during downhill on the 26" wheel bicycles again potentially indicates a greater demand on the  
21  
22  
23 10 biceps to manoeuvre the bicycle over and around obstacles. However, it is important to note  
24  
25 11 the relatively large standard deviations reported for the sEMG data. This may reflect  
26  
27 12 differences in riding styles of the participants, as riders may have adapted their riding styles  
28  
29 13 to suit the ride qualities of each bicycle. However, without video analysis of each ride, it is  
30  
31 14 difficult to quantify if or how much this differed between bicycles.  
32  
33  
34  
35 15

36  
37  
38 16 In addition, anthropometric differences may also have influenced the results and the large  
39  
40 17 standard deviations reported. Though not measured in the current study differences in limb  
41  
42 18 lengths between riders may have influenced these findings, as this would influence leverage  
43  
44 19 at the cranks and handlebars and thus affect how easily a rider could manoeuvre the bicycle  
45  
46 20 over or around trail obstacles. Given the lack of significant difference in mean velocity over  
47  
48 21 the course, difference between the three bicycles was less than  $0.4 \text{ km}\cdot\text{h}^{-1}$ , it is less likely that  
49  
50 22 would have influenced the muscle activity and vibrations encountered by the riders.  
51  
52  
53  
54 23



1  
2  
3 1 Lastly, upper limb sEMG may also have been influenced by braking activity. Hurst et al.  
4  
5 2 (2012) reported increased forearm activity in elite cross-country riders when compared to  
6  
7 3 downhill riders. This they proposed was possible the result of frame geometry differences and  
8  
9 4 smaller brake rotors. However, in the present study, equipment was the same on all bicycles  
10  
11 5 as was the head tube angle, therefore it is unlikely that these would have impacted on the  
12  
13 6 finding. Differences were more likely the result of riding style, as some riders may have  
14  
15 7 braked more into corners than other, resulting in the high standard deviations observed in  
16  
17 8 biceps brachii sEMG. Future research may seek to quantify the contribution of braking to  
18  
19 9 muscle sEMG activity.  
20  
21  
22  
23  
24 10

25  
26  
27 11 Previous research has tried to evaluate the external load imposed on riders during mountain  
28  
29 12 biking (Hurst, Swarén, Hébert-Losier, Ericsson, Sinclair, Atkins, & Holmberg, 2013). Their  
30  
31 13 study used a tri-axial accelerometer positioned between the scapulae. However, they reported  
32  
33 14 values as a global index of load/vibration on the whole body, which did not reflect the  
34  
35 15 vibrations exerted on specific muscle groups or joint. As such, the present study analysed  
36  
37 16 total acceleration (XYZ) (RMS) as an indicator of vibrations imposed on muscles in the  
38  
39 17 upper and lower limbs. Other than the expected differences in vibrations between the  
40  
41 18 gastrocnemius and vastus lateralis, due to the different locations from the body/bicycle  
42  
43 19 interface, no significant differences were found for muscles between wheel sizes or within  
44  
45 20 muscle groups. This is in contrast to the findings of Macdermid et al. (2014) who found the  
46  
47 21 29" wheel resulted in significantly greater vibrations than the 26" wheeled mountain bike. In  
48  
49 22 addition, data for vibrations presented in the current study are approximately double the  
50  
51 23 magnitude of those reported by Macdermid et al. (2014). This may be the result of  
52  
53 24 differences in sensor placement, as Macdermid et al. (2014) positioned their accelerometers  
54  
55 25 on the bony processes of the wrist and ankle, whilst sensors in the present study were  
56  
57  
58  
59  
60

1 positioned on soft tissue on the medial aspects of the muscles to be analysed. Chiementin,  
2 Rigaut, Crequy, Bolaers, and Bertucci (2012) also state that in order to assess vibrations in  
3 cycling the accelerometers should be located at the machine-hand interface. Though this is a  
4 valid point, within the present study positioning of the sensors in this region would have  
5 prevented the ability to determine muscle activity, due to the multi-sensor nature of the  
6 Delsys Trigno system. Therefore, these differences may account for much of the differences  
7 between Macdermid et al. (2014) and the present study.

8 Research by Lafortune, Hening, and Valiant (1995) and Coza, Nigg, and Fliri (2010)  
9 previously validated the use accelerometers placed on soft tissue to monitor vibrations on  
10 human muscle. Lafortune et al. (1995) did however reported values approximately double the  
11 amplitude of those recorded by accelerometers placed on the bone. Additionally, Kim et al.  
12 (1993) also found that skin mounted accelerometers increased vibrations by approximately 12  
13 % when compared to bone mounted accelerometers. Though it is generally believed that  
14 muscle would act as a damper and attenuate the accelerometer signals, during exercise the  
15 increased eccentric activity of the muscles upon landing forces, or those experienced during  
16 descending and jumping in mountain biking in our case, have been shown to increase  
17 intramuscular tension and therefore augment acceleration signals (Mizrahi, Verbitsky, &  
18 Isakov, 2000). This may in part account for some of the differences in results between the  
19 current study and those of Macdermid et al. (2014). Arpinar-Avsar, Birlik, Sezgin, and Soylu  
20 (2013) sought to quantify the vibrations imposed on the forearm of the rider when riding a  
21 mountain bike. They reported RMS values for acceleration again approximately half those  
22 reported in the present study. However, their study only involved riders riding over rough  
23 road for a distance of 250 m and not terrain representative of off-road cycling. This most  
24 likely contributed to the lower values reported.

25

1  
2  
3 1 Differences may also be attributed to the fact that full suspension bicycles were used in the  
4  
5 2 present study compared to the rigid hard tail mountain bike used by Macdermid et al. (2014).  
6  
7 3 As a consequence, this may have influenced riding dynamics. It is possible that the lower  
8  
9 4 accelerations observed by Macdermid et al. (2014) may have resulted from greater flexion of  
10  
11 5 the arms and legs to absorb trail shocks more effectively, whilst in the present study riders  
12  
13 6 may have relied more on the bicycle's suspension to absorb the shocks. However, this  
14  
15 7 potentially could have led to participants riding with straighter arms and legs and  
16  
17 8 unknowingly increasing the vibrations transmitted to the muscles, possibly during the shocks  
18  
19 9 rebound phase. This idea may be supported by Hurst et al. (2012) who found downhill riders  
20  
21 10 had a smaller degree of elbow flexion compared to cross-country riders as a result of the  
22  
23 11 greater level of suspension travel used on downhill specific bicycles.  
24  
25  
26  
27  
28  
29  
30

31 13 The non-significant differences in vibrations between the three wheel sizes in the present  
32  
33 14 study may again be due to the effects of the suspension systems effectively reducing trail  
34  
35 15 shock. In addition to the lack of rear suspension on the bicycle used in the Macdermid study,  
36  
37 16 they also fitted the 26" wheels to a frame designed for 29" wheels. This may have altered the  
38  
39 17 handling characteristics of the bicycle, as the bottom bracket and crank arms would have  
40  
41 18 been lower to the ground than when using frames designed specifically for the respective  
42  
43 19 wheel sizes.  
44  
45  
46  
47  
48  
49  
50

## 51 **Limitations**

52  
53 22 One potential limitation of the present study was the lack of gyroscopic and magnetometer  
54  
55 23 sensors within the sensor units used. The inclusion of such sensors could have aided in more  
56  
57  
58  
59  
60

1  
2  
3 1 accurate determination of the directionality of the acceleration forces experienced by the  
4  
5 2 riders. Such information could have helped identify differences in riding styles between  
6  
7 3 wheel sizes and whether riders were required to use greater lateral, vertical or horizontal  
8  
9 4 efforts to manoeuvre the bicycles through different sections of the course. In addition, the  
10  
11 5 inclusion of these sensors may have allowed the use of spectral analyses to determine the  
12  
13 6 frequency of vibrations in the vertical and horizontal planes, as with the current sensor  
14  
15 7 configuration and the movement of the riders it would have been practically impossible to  
16  
17 8 identify these vertical and horizontal frequencies.  
18  
19  
20  
21  
22  
23  
24

## 25 **Conclusions and practical implications**

26  
27  
28 11 In conclusion, this study revealed that larger wheel diameters did not significantly reduce  
29  
30 12 muscle activity or vibrations more so than the long standing traditional 26" wheel on the  
31  
32 13 particular course used. However, there appeared to be a trend for slightly greater biceps  
33  
34 14 brachii activity in the 26" wheel. This could have been the result of the need to lift the front  
35  
36 15 wheel over small obstacles more or differences in rider anthropometrics or a combination of  
37  
38 16 both. Additionally, differences in riding styles or adaptations to riding the different wheeled  
39  
40 17 bicycles may have played a part in muscle activity. Ultimately, the lack of statistical  
41  
42 18 differences between the measured parameters suggests that when ridden by trained mountain  
43  
44 19 bike riders all three wheel sizes are just as effective at controlling trail induced shocks and  
45  
46 20 that bicycle choice is largely a matter of personal preference. Within a field based setting,  
47  
48 21 there are likely to be a number of factors that influence muscle activity rather than wheel size  
49  
50  
51 22 alone.  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 **1 References**  
4

- 5  
6 2 Arpinar-Avsar, P., Birlik, G., Sezgin, Ö.C., & Soylu, A.R. (2013) The effects of surface-  
7 induced loads on forearm muscle activity during steering a bicycle. *Journal of Sports*  
8  
9 3 *Science and Medicine*, 12, 512-520.  
10  
11 4  
12  
13  
14 5 Blake, O.M., Champoux, Y., & Wakeling, J.M. (2012) Muscle Coordination Patterns for  
15  
16 6 Efficient Cycling. *Medicine and Science in Sports and Exercise*, 44, 926-938.  
17  
18  
19 7 Chimentin, X., Rigaut, M., Crequy, S., Bolaers, F., & Bertucci, W. (2012) Hand-arm  
20  
21 8 vibration in cycling. *Journal of Vibration Control*, 19, 2551-2560.  
22  
23  
24  
25 9 Coza, A., Nigg, B.M., & Fliri, L. (2010) Quantification of soft-tissue vibrations in  
26  
27 10 running: accelerometry versus high-speed motion capture. *Journal of Applied*  
28  
29 11 *Biomechanics*, 26, 367-373.  
30  
31  
32  
33 12 De Luca, C.J., Kuznetsoc, M., Gilmore, L.D., & Roy, S.H. (2012) Inter-electrode spacing  
34  
35 13 of surface EMG sensors: Reduction of crosstalk contamination during voluntary  
36  
37 14 contractions. *Journal of Biomechanics*, 45, 555-561.  
38  
39  
40  
41 15 Duc, S., Bertucci, W., Pernin, J.N., & Grappe, F. (2008) Muscular activity during uphill  
42  
43 16 cycling: Effect of slope, posture, hand grip position and constrained bicycle lateral sways.  
44  
45 17 *Journal of Electromyography and Kinesiology*, 18, 116-127.  
46  
47  
48  
49 18 Egaña, M., Ryan, K., Warmington, S.A., & Green, S. (2010) Effect of body tilt angle on  
50  
51 19 fatigue and EMG activities in lower limbs during cycling. *European Journal of Applied*  
52  
53 20 *Physiology*. 108, 649–656.  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1 Faiss, R., Praz, M., Meichtry, A., Gobelet, C., & Deriaz, O. (2007) The effect of  
4  
5 2 mountain bike suspensions on vibrations and off-road uphill performance. *Journal of*  
6  
7 3 *Sports Medicine and Physical Fitness*, 47, 151-158.  
8  
9  
10 4 Hurst, H.T., & Atkins, S. (2006) Power output of field-based downhill mountain biking.  
11  
12 5 *Journal of Sports Sciences*, 24(10), 1047-1053.  
13  
14  
15  
16 6 Hurst, H.T., Swarén, M., Hébert-Losier, K., Ericsson, F., Sinclair, J., Atkins, A., &  
17  
18 7 Holmberg, HC. (2012) Influence of course type on upper body muscle activity in elite  
19  
20 8 cross-country and downhill mountain bikers during off road downhill cycling. *Journal of*  
21  
22 9 *Science and Cycling*, 1(2), 2-9.  
23  
24  
25  
26 10 Hurst, H.T., Swarén, M., Hébert-Losier, K., Ericsson, F., Sinclair, J., Atkins, A., &  
27  
28 11 Holmberg, HC. (2013) GPS-based evaluation of activity profiles in elite downhill  
29  
30 12 mountain biking and the influence of course type. *Journal of Science and Cycling*, 2(1),  
31  
32 13 25-32.  
33  
34  
35  
36 14 Impellizzeri, F.M., Rampinini, E., Sassi, A., Mognoni, P., & Marcora, S. (2005)  
37  
38 15 Physiological correlates to off-road cycling performance. *Journal of Sports Sciences*, 23,  
39  
40 16 41-47.  
41  
42  
43  
44 17 Kim, W., Volshin, A.S., Johnson, S.H., & Simkin, A. (1993) Measurement of impulsive  
45  
46 18 bone motion by skim-mounted accelerometers. *Journal of Biomechanical Engineering*,  
47  
48 19 115, 47-52.  
49  
50  
51  
52 20 Lafortune, M.A., Hennig, E., & Valiant, G.A. (1995). Tibial shock measured with bone  
53  
54 21 and skin mounted transducers. *Journal of Biomechanics*, 28, 989-993.  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1 Lee, H., Martin, D.T., Anson, J.M., Grundy, D., & Hahn, A.G. (2002) Physiological  
4 characteristics of successful mountain bikers and professional road cyclists. *Journal of*  
5 2 *Sports Sciences*, 20, 1001-1008.  
6  
7 3  
8  
9  
10 4 Levy, M., & Smith, G. (2005) Effectiveness of vibration damping with bicycle  
11 suspension systems. *Sports Engineering*, 8, 99-106.  
12 5  
13  
14 6 Macdermid, P.W., Fink, P.W., & Stannard R. (2014) Transference of 3D accelerations  
15 during cross country mountain biking. *Journal of Biomechanics*, 47, 1829-1827.  
16 7  
17  
18 8 MacRae, H.S-H., Hise, K.J., & Allen, P.J. (2000) Effects of Front and Dual Suspension  
19 Mountain Bike Systems on Uphill Cycling Performance. *Medicine and Science in Sports*  
20 *and Exercise*, Vol. 32, 1276-1280.  
21 9  
22  
23 10 Matsuura, R., Arimitsu, T., Yuncki, T., & Yuno, T. (2011) Effects of resistive load on  
24 performance and surface EMG activity during repeated cycling sprints on a non-  
25 isokinetic cycle ergometer. *British Journal of Sports Medicine*. 45, 820-824.  
26 11  
27  
28 12 Mester, J., Spitzenfeil, P., Schwarzer, J., & Seifriz, F. (1999) Biological reaction to  
29 vibration – implications for sport. *Journal of Science and Medicine in Sport*, 2, 211-226.  
30 13  
31  
32 14 Mizrahi, J., Verbitsky, O., & Isakov, E. (2000) Shock accelerations and attenuation in  
33 downhill and level running. *Clinical Biomechanics*, 15, 15-20.  
34 15  
35  
36 16 Nokes, L., Fairclough, J.A., Minitowt, W.J., Mackie, I., & Williams, J. (1984) Vibration  
37 analysis of human tibia: The effects of soft tissue on the output from skin-mounted  
38 accelerometers. *Journal of Biomechanical Engineering*, 6, 223-226.  
39 17  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1 Prins, L., Terblanche, E., & Myburgh, K.H. (2007) Field and laboratory correlates of  
4 performance in competitive cross-country mountain bikers. *Journal of Sports Sciences*,  
5 25, 927 – 935.  
6  
7  
8  
9  
10 4 Saha, S., & Lakes, R.S. (1977) The effect of soft tissue on wave-propagation and  
11 vibration tests for determining the in vivo properties of bone. *Journal of Biomechanics*,  
12 10, 393-401.  
13  
14  
15  
16  
17 7 Sinclair, J., Brooks, D., Edmundson, C.J., & Hobbs S.J. (2012) The efficacy of EMG  
18 MVC normalisation techniques for running analyses. *Journal of Biomechanics*, 45(1),  
19 623.  
20  
21  
22  
23  
24  
25 10 Sinclair, J., Taylor, P.J., & Hobbs, S.J. (2013) Alpha level adjustments for multiple  
26 dependent variable analyses and their applicability – A review. *International Journal of*  
27 *Sport Science and Engineering*, 7, 17-20.  
28  
29  
30  
31  
32  
33 13 Stapelfeldt, B., Schwirtz, A., Schumacher, Y.O., & Hillebrecht, M. (2004) Workload  
34 demands in mountain bike racing. *International Journal of Sports Medicine*, 25, 294-300.  
35  
36  
37 15 Ziegert, J.C., & Lewis, J.L. (1979) The effects of soft tissue on measurements of  
38 vibrational bone motion and skin-mounted accelerometers. *Journal of Biomechanics*, 101,  
39 218.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3

**Figure 1. GPS profile of course. Numbers indicate the different sections of the course.**

For Peer Review Only

1

2 **Table 1. Mean  $\pm$  standard deviation and 95 % CI** for mean sEMG amplitude as a percentage  
3 of dynamic peak task for different wheel size during (A) full lap; (B) climb; and (C) descent.  
4 \*Significantly different to triceps brachii (26"); † significantly different to triceps brachii  
5 (27.5"); ‡ significantly different to triceps brachii (29").

6

For Peer Review Only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4

**Table 2.** Mean  $\pm$  standard deviation and 95% CI of mean acceleration amplitude (RMS) for different wheel size during (A) full lap; (B) climb; and (C) descent.

For Peer Review Only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



Figure 1  
164x101mm (96 x 96 DPI)

	26" Wheel	27.5" Wheel	29" Wheel
	Mean ± sd (95% CI)	Mean ± sd (95% CI)	Mean ± sd (95% CI)
<b>Full Lap</b>			
Gastrocnemius	4.71 ± 3.23 (2.22, 7.19)	4.86 ± 3.28 (2.33, 7.38)	4.83 ± 3.12 (2.43, 7.22)
Vastus Lateralis	4.56 ± 2.54 (2.61, 6.52)	4.94 ± 2.99 (2.64, 7.23)	4.62 ± 2.31 (2.84, 6.40)
Biceps Brachii	1.78 ± 1.66 (0.51, 3.06)*	1.17 ± 0.74 (0.61, 1.74) <sup>†</sup>	1.32 ± 0.67 (0.80, 1.85) <sup>‡</sup>
Triceps Brachii	3.61 ± 1.28 (2.62, 4.59)	3.58 ± 1.34 (2.55, 4.61)	3.66 ± 1.70 (2.84, 4.48)
<b>Climb</b>			
Gastrocnemius	4.91 ± 3.33 (2.35, 7.47)	4.68 ± 3.10 (2.30, 7.05)	4.81 ± 3.07 (2.46, 7.17)
Vastus Lateralis	4.62 ± 2.51 (2.70, 6.55)	4.95 ± 3.04 (2.61, 7.28)	4.60 ± 2.41 (2.74, 6.45)
Biceps Brachii	1.43 ± 1.04 (0.63, 2.22)*	1.18 ± 0.87 (0.51, 1.85) <sup>†</sup>	1.09 ± 0.78 (0.49, 1.69) <sup>‡</sup>
Triceps Brachii	3.26 ± 1.20 (2.33, 4.18)	3.39 ± 1.17 (2.50, 4.29)	3.43 ± 1.02 (2.64, 4.21)
<b>Descent</b>			
Gastrocnemius	4.70 ± 3.43 (2.07, 7.34)	4.56 ± 3.24 (2.07, 7.05)	4.63 ± 3.12 (2.23, 7.03)
Vastus Lateralis	4.64 ± 2.94 (2.37, 6.90)	4.78 ± 2.87 (2.58, 6.99)	4.30 ± 2.14 (2.65, 5.95)
Biceps Brachii	2.29 ± 3.05 (-0.06, 4.63)	1.19 ± 0.74 (0.62, 1.76) <sup>†</sup>	1.44 ± 0.75 (0.86, 2.02) <sup>‡</sup>
Triceps Brachii	3.91 ± 1.35 (2.88, 4.95)	3.85 ± 1.60 (2.62, 5.08)	3.79 ± 1.12 (2.93, 4.65)

	26" Wheel	27.5" Wheel	29" Wheel
	Mean ± sd (95% CI)	Mean ± sd (95% CI)	Mean ± sd (95% CI)
<b>Full Lap</b>			
Gastrocnemius	68.50 ± 2.78 (66.37, 70.64)	66.28 ± 4.00 (63.21, 69.36)	67.35 ± 2.23 (65.64, 69.07)
Vastus Lateralis	57.66 ± 2.06 (56.08, 59.24)	57.26 ± 1.02 (56.47, 58.04)	57.26 ± 1.71 (55.95, 58.57)
Biceps Brachii	58.88 ± 5.39 (54.74, 63.03)	56.02 ± 7.12 (50.55, 61.49)	56.58 ± 5.70 (52.20, 60.96)
Triceps Brachii	62.48 ± 5.71 (58.09, 66.88)	61.44 ± 2.24 (59.71, 63.16)	61.10 ± 2.97 (58.82, 63.38)
<b>Climb</b>			
Gastrocnemius	66.73 ± 4.27 (63.45, 70.01)	66.07 ± 3.98 (63.02, 69.13)	64.91 ± 2.43 (63.04, 66.78)
Vastus Lateralis	56.82 ± 2.87 (54.61, 59.03)	57.18 ± 1.89 (55.73, 58.64)	55.90 ± 1.74 (54.57, 57.24)
Biceps Brachii	57.08 ± 7.12 (51.61, 62.56)	54.02 ± 8.21 (47.11, 60.33)	52.75 ± 5.96 (48.17, 57.33)
Triceps Brachii	60.31 ± 6.84 (55.05, 65.56)	59.63 ± 3.90 (56.63, 62.63)	58.39 ± 4.04 (55.29, 61.50)
<b>Descent</b>			
Gastrocnemius	69.68 ± 3.31 (67.13, 72.23)	66.91 ± 4.29 (63.61, 70.21)	68.28 ± 2.56 (66.31, 70.25)
Vastus Lateralis	58.05 ± 2.20 (56.36, 59.74)	57.34 ± 1.50 (56.19, 58.49)	57.64 ± 2.09 (56.03, 59.25)
Biceps Brachii	60.48 ± 6.21 (55.71, 65.25)	58.29 ± 7.99 (52.15, 64.44)	58.93 ± 6.04 (54.30, 63.57)
Triceps Brachii	64.45 ± 5.87 (59.93, 68.96)	63.38 ± 2.60 (61.38, 65.38)	63.31 ± 3.40 (60.70, 65.93)