



University of
Salford
MANCHESTER

Modulation of vascular reactivity by perivascular adipose tissue (PVAT)

Agabiti Rosei, C, Painsi, A, De Ciuceis, C, Withers, SB, Greenstein, A, Heagerty, AM and Rizzoni, D

<http://dx.doi.org/10.1007/s11906-018-0835-5>

Title	Modulation of vascular reactivity by perivascular adipose tissue (PVAT)
Authors	Agabiti Rosei, C, Painsi, A, De Ciuceis, C, Withers, SB, Greenstein, A, Heagerty, AM and Rizzoni, D
Type	Article
URL	This version is available at: http://usir.salford.ac.uk/id/eprint/46513/
Published Date	2018

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: usir@salford.ac.uk.

**MODULATION OF VASCULAR REACTIVITY BY PERIVASCULAR
ADIPOSE TISSUE (PVAT).**

Claudia Agabiti Rosei*, Anna Paini*, Carolina De Ciuceis*, Sarah Withers, Adam Greenstein, Anthony M. Heagerty, Damiano Rizzoni*.

Department of Medicine, Manchester University, United Kingdom, and * Clinica Medica, Department of Medical and Surgical Sciences, University of Brescia, Italy.

Running title: Effects of inflammation PVAT-dependent and oxidative stress on modulation of vascular tone

Keywords: perivascular adipose tissue, small arteries, hypoxia, oxidative stress, melatonin.

Wordcount: 7933

Address:

Clinica Medica, Department of Medical and Surgical Sciences,
University of Brescia.

c/o 2^a Medicina Spedali Civili di Brescia

Piazza Spedali Civili 1

25100 Brescia

Tel. 030 3905251

Fax 030 3384348

e-mail: claudiaagabitirosei@libero.it

Abstract

Purpose of Review: In this review we discuss the role of perivascular adipose tissue (PVAT) in the modulation of vascular contractility and arterial pressure, focusing on the role of the renin-angiotensin-aldosterone system and oxidative stress/inflammation.

Recent Findings: PVAT possesses an relevant endocrine-paracrine activity, which may be altered in several pathophysiological and clinical conditions. During the last two decades it has been shown PVAT may modulate vascular reactivity. It has also been previously demonstrated that inflammation in adipose tissue may be implicated in vascular dysfunction. In particular, adipocytes secrete a number of adipokines with various functions, as well as several vasoactive factors, together with components of the renin-angiotensin system which may act at local or at systemic level. It has been shown that the anticontractile effect of PVAT is lost in obesity, probably as a consequence of the development of adipocyte hypertrophy, inflammation, and oxidative stress.

Summary: Adipose tissue dysfunction is interrelated with inflammation and oxidative stress, thus contributing to endothelial dysfunction observed in several pathological and clinical conditions such as obesity and hypertension. Decreased local adiponectin level, macrophage recruitment and infiltration, and activation of renin-angiotensin-aldosterone system could play an important role in this regards.

1. Potential role of PVAT in the modulation of arterial pressure

The majority of blood vessels are surrounded by variable amounts of PVAT. For many years PVAT was routinely removed from arteries during *in vitro* studies, in order to simplify the experiments on isolated blood vessels. However, due to the increasing evidence of the role of the adipose tissue as an endocrine organ (1) and of the close proximity of PVAT to a vast network of blood vessels, investigators have begun to study the effects of PVAT (Table 1).

In 1991 Soltis and Cassis were the first to test the hypothesis that PVAT might be important for vascular regulation (2). They demonstrated the presence of a diminished response to noradrenaline in intact arteries compared with vessels having PVAT removed in isolated aortic ring preparations from rats. They attributed this effect to the dense sympathetic innervation in PVAT (2). Ten years later the interest in PVAT was renewed by Lohn et al (3). They demonstrated that the vasocontractile response with angiotensin II (Ang II), serotonin and phenylephrine was 95%, 80% and 30% lower respectively after removal of PVAT (3). Furthermore this effect was independent of nitric oxide (NO) formation and appeared to be mediated by a substance released from intact aortic rings surrounded with fat into the organ bath since transferral of bath solution from vessels with fat or cultured rat adipocytes to vessels without fat resulted in rapid relaxation of pre-contracted arteries. The factor involved was considered to be more likely a protein than a lipid, as it was not adsorbed by essentially fatty acid-free serum albumin and it was inactivated by heating. Therefore, the Authors concluded that “adventitium derived relaxing factor” (ADRF) was released by PVAT (3).

ADRF release was shown to depend on extracellular calcium and can be modulated by changes in calcium concentration in the plasma *in vivo* (4). Furthermore, ADRF effect is regulated by intracellular tyrosine kinase, protein kinase A and voltage dependent smooth muscle potassium channel but not by NO, cytochrome P450 pathways or prostaglandins (4). We now know that the relaxing factor is released from the adipose tissue and not from the tunica adventitia (5) and therefore we can use the term PVAT-derived relaxing factor (PDRF). Subsequent studies confirmed the role of PVAT as an inhibitor of arterial contraction in internal thoracic arteries (6) and in human internal mammary arteries (7). The mechanism of the vasodilatory capacity of PVAT varies depending on anatomical location and animal species. In rat aorta, PDRF appears to cause vasorelaxation by opening ATP-dependent K⁺ (K_{ATP}) channels in vascular smooth muscle cell (4), whilst in rat mesenteric arteries, voltage dependent K⁺ channels (K_v) seem to be involved (8). In a recent study Gao et al identified two different pathways for explaining the PVAT relaxation on rat aorta. They described an endothelium-dependent and endothelium independent mechanism, working through release of nitric oxide with calcium sensing K⁺ channels activation (K_{Ca}) (5) and hydrogen peroxide with soluble guanylate cyclase activation (5), respectively. Several studies both *in vitro* and *in vivo* have confirmed the anticontractile property of PVAT but despite several candidates being proposed, the real identity of PDRF is partly unknown.

Candidates for PDRF :

Leptin: Leptin has been proposed as PDRF because of its direct vasodilator effect by release of nitric oxide with endothelium-dependent mechanism (9,10). However, leptin possesses also contractile effects by activation of the sympathetic nervous

system and several other effects such as increase in vasoconstrictor endothelin-1 (11) and in toxic reactive oxygen species (ROS) (12). Therefore, leptin may contribute to arterial pressure homeostasis and to the anticontractile effect of PVAT but its role as PDRF seems to be unlikely.

Adiponectin: The PVAT has been shown to secrete other vasodilator products like hydrogen peroxide (13) and adiponectin (14,15). In particular adiponectin is a protein secreted exclusively by adipose tissue (16) and it has been reported to possess several protective properties against hypertension (17,18,19), atherosclerosis (20) and diabetes (21). Several studies have demonstrated that adiponectin acts as a vasodilator by activation of endothelial NO synthase (eNOS) and consequent increase of NO production (21). It has the ability to inhibit macrophage activation, to reduce proliferation of vascular smooth muscle cells, to improve insulin signalling pathways and to reduce ROS levels (22,23). Hypoadiponectinemia has been correlated to endothelial dysfunction (24), obesity (25), hypertension (18), atherosclerosis (19), type 2 diabetes (20) and myocardial infarction. In particular, it has been shown that in patients with coronary artery disease epicardial tissue expresses lower adiponectin levels (26). Recent studies have demonstrated adiponectin-mediated anticontractile properties of PVAT in human small arteries, which represent the main regulators of peripheral resistance and hence of blood pressure (27). In particular, in that study it has been demonstrated that the anticontractile function of PVAT disappears in healthy human arteries after blockade of adiponectin receptor 1 (27). Considering its beneficial and protective effects, adiponectin could be an important therapeutic target. Therefore adiponectin might represent a reliable candidate for the role of PDRF. Aside from vasodilator products, PVAT has been also reported to secrete vasoconstrictor peptides like endothelin and Ang II (28) (Figure 1).

Inflammatory cytokines: The adipokines group also includes proinflammatory cytokines (such as interleukin 6 [IL-6], tumor necrosis factor [TNF- α]) chemokines (like interleukin 8 [IL-8] and monocyte chemoattractant protein-1 [MCP-1]) growth factors (transforming growth factor- β [TGF- β]) and pro-thrombotic factors (eg, plasminogen activator inhibitor type 1 [PAI-1]).

There is a compelling evidence for an increased level of adipokines in inflammatory state, such as in obesity, and of a decreased cardiometabolic protective role of adiponectin. These obesity-related changes in adipose tissue are probably linked to the development of insulin resistance, type 2 diabetes mellitus, hypertension and, thus, metabolic syndrome (29-32).

In addition to this emergent role of PVAT as modulator of arterial tone, in a recent study it has been suggested that PVAT may act also as a regulator of vein function. In particular PVAT is able to reduce contractile responses of the inferior vena cava to various agonists mediated by angiotensin (1-7) through activation of K_v channels (33). Therefore the PVAT may play a role as dual modulator of vascular tone.

It is clear that adipose tissue and in particular PVAT is a secretory and endocrine organ which is involved in the regulation of vascular tone. PVAT might play an important role in the modulation of blood pressure, in the hemodynamic homeostasis and in other conditions associated to vascular dysfunction such as atherosclerosis, obesity and type 2 diabetes mellitus. This suggests that a balance between adipose tissue-derived vasodilator and vasoconstrictor mediators might be extremely important for the maintenance of an appropriate vascular tone.

Table 1: Potential mechanisms involved in vascular-relaxation by PVAT

Study	District	Species	Agonist	Effect of PVAT on vascular tone	Potential mechanism proposed
Soltis et al (1991) ²	Aorta	Rat	Noradrenaline	Vasodilatation	Increased uptake of Noradrenaline with PVAT
Lohn et al (2002) ³	Aorta	Rat	Phenylephrine Angiotensin II Serotonin	Vasodilatation	1. Endothelium independent action 2. Activation of ATP dependent K ⁺ channels (K _{ATP})
Dubrovskaja et al (2003) ⁴	Aorta	Rat	Phenylephrine and Serotonin	Vasodilatation	1. Ca ²⁺ dependent 2. Tyrosine & protein kinase A dependent intracellular signalling pathways 3. Independent of perivascular nerves
Archer SL et al (2003) ⁷	Internal mammary arteries	Human	Bradykinin, Acetylcholine & 11,12-epoxyeicosatrienoic acid	Vasodilatation	Activation of Smooth Muscle Cells large-conductance Ca ²⁺ -activated K ⁺ channels (BK _{Ca}).
Verlohren et al (2004) ⁸	Mesenteric arteries	Rat	Phenylephrine Adrenaline Serotonin	Vasodilatation	1. Endothelium independent action 2. Activation of Voltage dependent K ⁺ channels (K _v)
Gao et al (2005) ⁶	Internal thoracic artery	Human	Phenylephrine U46619	Vasodilatation	Activation of Calcium sensing K ⁺ channels (K _{Ca})
Galvez et al (2006) ⁹	Mesenteric arteries	Rat	Serotonin U46619	Vasodilatation	Activation of Voltage dependent K ⁺ channels (K _v)
Gao et al (2007) ⁵	Aorta	Rat	Phenylephrine Serotonin	Vasodilatation	1. Endothelium-dependent relaxation by Nitric oxide release & K _{Ca} activation 2. Endothelium independent mechanism involving H ₂ O ₂ & activation of soluble Guanylate Cyclase
Malinowski et al (2007) ³⁴	Internal thoracic artery	Human	Angiotensin II Serotonin	Vasodilatation	non-Nitric Oxide, non-prostacyclin dependant mechanism
Greenstein et al (2009) ²⁷	Subcutaneous gluteal biopsy small arteries & mesenteric arteries	Human and rat	Norepinephrine	Vasodilatation	Endothelium-dependent relaxation by adiponectin through an increase Nitric oxide bioavailability
Lee RM et al (2009) ^{35,36}	Mesenteric resistance arteries, Aorta	Rat	Phenylephrine, serotonin, Angiotensin II	Vasodilatation	Endothelium-dependent relaxation by angiotensin 1-7 through Nitric Oxide, via activation K _{Ca} channels
Lu C et al (2011) ³³	Inferior vena cava	Rat	Phenylephrine, U46619 & 5-hydroxytryptamine	Vasodilatation	Endothelium-dependent relaxation by angiotensin 1-7 via the activation K _v channels




Figure 1: Hormones and other substances secreted by adipose tissue. Abbreviations: ASP, acylation stimulating protein; IL-6, interleukin 6; PAI-1, plasminogen activator inhibitor type 1; TF, tissue factor; TGF- β , transforming growth factor β ; TNF α , tumor necrosis factor α . Diagram modified from Ahima RS, Flier JS. Adipose tissue as an endocrine organ. *Trends in Endocrinology & Metabolism* 2000; 11(8): 327-32.

2. The adipose tissue and the renin-angiotensin axis

The renin-angiotensin system (RAS) is involved in systemic blood pressure regulation and in renal electrolyte homeostasis. Several studies have demonstrated the presence of local RAS activity in both white and brown perivascular adipose tissue (37).

A number of animal studies have demonstrated the expression of most of the RAS system components in adipose tissue (38), however data from human tissues are relatively few (39) (Figure 2).




Figure 2: Perivascular adipose tissue and local Renin-Angiotensin System. From Cassis LA, Police SB, Yiannikouris F, Thatcher SE. Local adipose tissue renin-angiotensin system. *Curr Hypertens Rep* 2008; 10(2): 93-98.⁽⁴⁰⁾

Angiotensinogen (AGT) expression in adipose tissue was first identified in periaortic brown adipose tissue in 1987 (41). Subsequently, AGT secretion and AGT-mRNA were detected in rat and human adipose tissue depots, in adipocytes isolated from rat arterial vessels walls and mesentery (42), in primary cultured human adipocytes (35) and in differentiating human pre-adipocytes (43). Thus the AGT expression and secretion is considered a late marker of adipocyte differentiation (44). Various studies have demonstrated that AGT expression is higher in visceral compared to subcutaneous adipose tissue in rats (45) and humans (46). In addition, white adipose tissue is considered to be one of the major sources of angiotensinogen outside the liver (47). The AGT activation by free fatty acids and overfeeding provides a possible link between adipose-tissue RAS, obesity/insulin resistance and obesity/hypertension.

Although renin activity has been detected in brown adipose tissue (48), Gálvez-Prieto et al showed the absence of renin expression in mesenteric and aortic PVAT from male Wister Kyoto rats (38). Nevertheless, they confirmed the presence of the (pro)renin receptor which may act with a circulating enzyme to increase local synthesis of angiotensin I from AGT. The peroxisome proliferator activated receptors (PPARs), of which two isoforms (α and γ) have been identified in PVAT, have both been shown to positively regulate renin transcription (49). This receptor has been detected in subcutaneous and visceral human adipose tissue (50).

Ang II is the major component of RAS with a variety of physiological actions including vasoconstriction, stimulation of aldosterone release from the adrenal gland, renal sodium and water reabsorption, increase of blood pressure, cell growth, inflammation process, vascular and cardiac remodelling, activation of the sympathetic

nervous system and ROS production (51). In the adipose tissue Ang II may play a role in adipocyte growth and differentiation, stimulating lipogenesis, pre-adipocytes recruitment and their differentiation in mature adipocytes (40).

The exact role of Ang II in the differentiation and proliferation of human adipocytes remains controversial; it has been shown that preadipocytes from human adipose tissue express a functional renin angiotensin system (RAS) (52). Sharma hypothesized that Ang II may be able to inhibit the adipogenic differentiation of primary cultured preadipocytes in humans (53). Ang II levels in mesenteric adipose tissue are higher than in periaortic adipose tissue (38), demonstrating a regional different expression of the RAS.

The effects of Ang II are mediated by two main membrane receptors: the angiotensin type 1 receptor (AT1R) and the angiotensin type 2 receptor (AT2R). Matsushita et al. demonstrated that Ang II stimulated differentiation of adipocytes through AT1R and inhibited it through AT2R (54). The AT1R is responsible for most biological effects of Ang II, such as pressure, trophic and pro-inflammatory effects, whereas the AT2R antagonizes several AT1R-mediated effects. It has been demonstrated that Ang II through AT1R stimulates transcription factor expression, which may promote adipocytes differentiation and increase of triglycerides content in human adipocytes culture and in 3T3-L1 cells, thus leading to adipocyte hypertrophy (55). The crucial role of AT1R is confirmed by the demonstration that, in essential hypertensives patients, in adipocytes in culture and in a rat model of type 2 diabetes, AT1R antagonist is able to increase adiponectin concentrations and improve insulin sensitivity (56,57). In addition, it has been demonstrated that telmisartan, a selective AT1R-blocker, increases adiponectin level and reduces myocardial damage in infarcted Zucker diabetic fatty rats models (58). It has been already shown that plasma

adiponectin levels were significantly increased in patients with hypertension after treatment with telmisartan (59). A previous study had also shown that treatment with the AT1R antagonist, olmesartan, in Ang II-infused rats resulted in an increase of adiponectin level (60)

Conversely, the binding between Ang II and its AT2R may determine tissue regeneration (25) and up-regulation of adiponectin production in neonatal rat ventricular myocytes (61)

Although the role of Ang II on differentiation and proliferation of adipocytes is still controversial, Ang II -related effects on adipose tissue, with its trophic and pro-inflammatory effects, could play an important role for vascular function and structure and could be implicated in hypertension-related obesity.

3. Changes of the RAS-axis in obesity and hypertension: contribution of the tissue RAS of PVAT

Abdominal obesity is an important risk factor for the development of arterial hypertension, insulin resistance and type 2 diabetes. Abdominal obesity is also associated with metabolic and morphological abnormalities including sodium retention, endothelial dysfunction, left ventricular hypertrophy, dyslipidemia, hyperinsulinemia, microalbuminuria, elevated markers of inflammation and oxidative stress (62).

Increasing evidence suggests a link between renin-angiotensin-aldosterone system (RAAS) and other components of the metabolic syndrome, in particular abdominal obesity. A Japanese study has shown that oxidative stress, an important mechanism of metabolic dysregulation in obesity, is able to reduce mRNA-AGT levels in adipose tissue of obese and in hypertrophic murine adipocytes (63). Oxidative stress is closely

correlated to RAAS; Ang II is a potent inducer of reactive oxygen substances (ROS) and during obesity systemic and related adipose tissue Ang II levels are increased.

In addition to Ang II, aldosterone is also an important mediator of RAAS effects. In obesity-related hypertension aldosterone levels are higher and they are associated with endothelial dysfunction (64) and inflammation (65). In a study of 2009, it has been shown that the blockade of mineralcorticoid-receptors improves adipocyte dysfunction and insulin resistance in a murine model of obesity (66). A recent review discusses the known interaction between adiponectin and aldosterone, demonstrating an inverse relationship (67). Several effects of aldosterone are mediated through pathways similar to those induced by Ang II. Furthermore, recent data demonstrate the presence of cross-talk between aldosterone and Ang II which modulates and stimulates the signalling transduction such as the Ang II effects on PVAT (68).

The tissue RAS of PVAT may operate in combination or independently of circulating RAS. The precise function of RAS in perivascular adipose tissue is still controversial but may contribute to vascular tone, structural and functional alterations, especially in pathological conditions such as hypertension and obesity, amplifying the effect of systemic RAS. It has been hypothesized cross-talk between smooth muscle cells and endothelial cells which could explain at least in part the implication of tissue RAS in the perivascular adipose tissue on structural alterations of small resistance arteries.

In view of the relevant role of the RAS system and PVAT on vascular function, it is interesting to understand the possible relationship between them.

In 2006 Gálvez and colleagues observed several changes in PVAT function and mass in spontaneously hypertensive rats (SHR) which could contribute to increased vascular resistance (9). In the SHR model, PVAT has a reduced anticontractile effect,

being the mass of mesenteric PVAT less than in control mice with smaller adipocytes and lower leptin content (9). Subsequently, Lee demonstrated that the changes on PVAT structure and function were observed also in Ang II-induced hypertension in adult male Wistar normotensives rats (35). In addition, it has been also shown that Ang II is able to mediate PVAT-associated increase of contractile response to perivascular neuronal excitation by EFS, possibly through superoxide production (69). Recently the same author confirmed the reduced anticontractile function of PVAT in SHR (70) and this functional impairment could involve angiotensin (1-7) (70). In fact, angiotensin (1-7) blockade is able to inhibit the anticontractile effect of PVAT (36). Ang 1-7 has been also implicated on anticontractile function showed by PVAT on inferior vena cava (33).

In a study of 2011, in a mouse model, we demonstrated that PVAT in small mesenteric arteries loses its anticontractile effect, after stimulation with two physiological stimuli of inflammation: aldosterone and hypoxia (71).

In that study we demonstrated that the presence and activation of macrophages in adipose tissue was a key modulator of the increased in contractility in arteries with PVAT following induction of inflammation. Among multiple factors that may be involved in determining the vascular consequences of obesity and hypertension, we have shown a favorable effect of eplerenone in inducing a reduction of the inflammatory effects of both aldosterone and hypoxia (71). This property of mineralocorticoid-receptors blockers may be of potential therapeutic interest. In a recent study we demonstrated that also an ACE-inhibitor and an AT1R blocker have similar effects on anticontractile properties of the perivascular adipose tissue following hypoxia (72). In particular, we have shown that hypoxia induced a loss of

this anticontractile effect which could be completely prevented with captopril or telmisartan. Therefore our *in-vitro* induction of a hypoxic environment can simulate the loss of anticontractile perivascular adipose tissue function seen *in vivo* in obese patients, and this can be prevented using inhibitors of the renin–angiotensin cascade; RAS blockade may protect the vasculature and may reverse the effect of perivascular and systemic RAS.

4 The role of oxidative stress and inflammation in PVAT function

It is well known that vascular dysfunction is mediated by adipose tissue inflammation which could be contrasted by anti-inflammatory agents such as adiponectin. Recently a review from Huang Cao et al described the important role of PVAT on regulation of vascular tone underlining the importance of the functional integrity of PVAT, rather than the amount of PVAT itself, because it's essential for the control of blood pressure and it's protective against the development of hypertension and cardiovascular disease (73). In particular in lean conditions the PVAT is able to secrete several adipokines which act on endothelial cells through a nitric-oxide dependent mechanism and directly on small muscle cells to induce vasorelaxation of the vessel wall. Instead during obesity the secretory profile of PVAT change with a reduction of expression of vasorelaxing factors and an increase in vasoconstricting factors with cell infiltration leading to inflammatory state (73). The cause of the onset of inflammation remains still unknown although a recent hypothesis is related to hypoxia; in obesity, adipocytes become hypertrophic leading to inadequate perfusion and consequent local hypoxia. Hypoxia-inducible factor (HIF-1 α) alpha is a key mediator of hypoxia. Its levels are higher in adipose tissue of obese subjects and they

decrease after weight loss (74). HIF-1 α is a main hypoxia-inducible transcription factor responsible for increased inflammatory mediator production, like TNF- α and IL-6, and for reduced adiponectin level (75). Greenstein *et al* have shown that the application of TNF- α and IL-6 to PVAT around healthy blood vessels decreased the dilator effect of PVAT. Similarly, it has been shown that the induction of experimental hypoxia for 2.5 hours causes inflammation and loss of anticontractile function of PVAT. Both conditions could be reversed and the anticontractile property rescued by catalase and superoxide dismutase or cytokine antagonists (27). From these observations it is clear that PVAT loses its dilating effect after the increase of oxidative stress.

In addition to hypoxia, there are also different HIF activators such as growth factors, cytokines and hormones. In particular, it has been demonstrated that in vascular smooth muscle cells, Ang II is a potent activator of HIF-1 α by mechanisms even different from hypoxia, involving also mitochondrial reactive oxygen species (mtROS) (76,77). As previously mentioned, it has been shown that hypoxia and aldosterone can reproduce the effect of an inflammatory phenotype in adipose tissue, with loss of anticontractile property (71). The macrophage activation is responsible of both conditions because free radical scavengers are able to rescue the anticontractile function of adipose tissue (71). In conclusion, inflammation and oxidative stress are interrelated and associated with adipose tissue dysfunction and they could contribute to arterial stiffness and hypertension (78,79), maybe through endothelial dysfunction (Figure 3). Macrophages in adipose tissue represent a key modulator of oxidative stress and systemic inflammation through the production of IL-6 and ROS (80). The important role of IL-6 and MCP-1 in the recruitment of inflammatory cells, in particular monocytes and macrophage, and in the pathology of vascular disease

during obesity is described in a recent review (73). Decreased adiponectin level is involved in the process of macrophage infiltration (81) and is a consequence of inflammation and oxidative stress (80) (Figure 3).




Figure 3: Hypothetical link between adipose tissue dysfunction, inflammation, oxidative stress and vascular disease.

Therefore it's clear that adiponectin is one of the key mediator of PVAT function which is altered in obesity. It was demonstrated that the effect of adiponectin on vascular tone is mediated by activation of potassium channel Calcium-dependent (BKCa channels) on vascular smooth muscle cells and adipocytes and by endothelial

mechanisms (82): adiponectin hyperpolarizes wild-type arteries, but not vessels from BKCa- knockout mice. Furthermore it seems that stimulated release of a hyperpolarizing factor (maybe adiponectin) from mesenteric artery PVAT induces electrical changes that involve myocyte BKCa channels (83). Thereafter the intracellular downstream mechanisms which involve cGMP-dependent protein kinase (PKG) as a regulators of normal PVAT function was described. (84). Recently the involvement of BKCa channels in anticontractile function of perivascular adipose tissue was confirmed (85). In particular, it has been shown that the anticontractile effect of PVAT completely disappeared after pre-incubation with iberiotoxin, a scorpion toxin that inhibits BKCa channels, in ob/ob mice, an animal model of obesity (B6. V-Lepob/OlaHsd) (85). In that study also the effects of melatonin, an endogenous hormone with antioxidant and vasculoprotective properties, on anticontractile properties of PVAT were addressed.

5 The role of melatonin in vascular dysfunction

Melatonin is an endogenous hormone that exerts antioxidant, antiinflammatory, antihyperlipidemic, and antihypertensive actions and it also modulates insulin secretion and action (86).

It has been demonstrated that melatonin is able to improve mesenteric small resistance artery structure and endothelial function in spontaneously hypertensive rats (87). It has been shown that melatonin therapy is able to improve blood pressure, lipid profile, and parameters of oxidative stress in patients with metabolic syndrome (88). Furthermore, chronic administration of melatonin in rats fed a normal or high-fat diet, resulted in significant reduction in body weight, in circulating insulin, glucose and triglyceride mean levels and was able to modulate the normal circadian pattern of

plasma adiponectin (89). It has also been proposed that melatonin-induced hypertrophy and functional activation of brown adipose tissue may provide a potential target for treatment of obesity in humans (90). Therefore, in obesity melatonin might be able to reduce pro-inflammatory cytokines and to increase adiponectin level with consequent modulation of PVAT function. Recently, it has been demonstrated that prolonged administration of melatonin is able to reduce hyperglycemia associated with obesity in hyperphagic mouse and to ameliorate the inflammation in the perivascular environment (85). In particular ob/ob mice treated with melatonin showed a marked reduction in the expression of endothelin-1 (ET-1), interleukin-6 (IL-6) , and metalloproteases 2 and 9; in addition we observed that the increased expression of both TNF- α and CD68 in visceral fat sections from ob/ob mice was significantly reduced after melatonin treatment. Anticontractile function of PVAT, partially lost in a model of animal obesity, may be restored by melatonin treatment but only in presence of an intact PVAT, indicating the importance of PVAT oxidative stress in vascular dysfunction observed in obese animals (85).

Aging represents another vascular dysfunction condition which is associated with a progressively decrease of the nighttime peak of melatonin concentrations. During aging structural and functional changes have been observed. In particular, the effect of aging on vascular endothelium and small muscle cells (SMC) has been widely investigated, while less is known about the changes of PVAT. A senescence-accelerated prone mouse (SAMP8), a model of age-related vascular dysfunction with associated increase in blood pressure and cognitive decline has been recently investigated (91, 92). It has been demonstrated that in SAMP8 mice there was an overexpression of endothelin 1 (ET-1), inducible nitric oxide synthase (iNOS) and cyclooxygenase 2 (COX-2), all markers of oxidative stress together with a reduced

level of endothelial nitric oxide synthase (eNOS) and cyclooxygenase 1 (COX-1) (93). In addition, in SAMP8 mice PVAT had lost its protective anti contractile effect. Also long term treatment with melatonin in SAMP8 was able to increase some vasculoprotective markers, to decrease oxidative stress and inflammation and to restore the anticontractile effect of perivascular adipose tissue (93). Decreased expression of adiponectin and adiponectin receptor 1 was also observed in visceral fat of untreated aging mice, whereas a significant increase was observed after melatonin treatment (93). The increased production/activity of adiponectin might contribute to explain the improvement of the anticontractile action of perivascular fat observed in the mesenteric small resistance arteries of SAMP8 mice after chronic treatment with melatonin (93).

6 Conclusions

In conclusion, several evidences suggest a key role of PVAT as regulator of vascular tone through different mechanisms. It is clear that the RAS activity in adipose tissue plays an important role on PVAT function and the correction of the imbalance between the different RAS components is crucial for the maintenance of vascular tone. Besides the known antihypertensive effects of ACE-inhibitors and AT1R-blockers and their effect on reduction of the incidence of new-onset diabetes (94-96), they are able to increase circulating levels of adiponectin. In particular, it has been demonstrated that the levels of myocardial adiponectin and its type 1 receptor are decreased in type 2 diabetic rats and that telmisartan treatment may correct these alterations (97). In addition melatonin represents a potential therapeutic biological target in clinical conditions such as obesity and for delaying physiological and pathological alteration related to early vascular aging.

Further studies are needed for fully understanding the potential implications of perivascular adipose tissue properties on vascular function in humans.

REFERENCES

References Papers of particular interest, published recently, have been highlighted as:

- Of importance
- Of major importance

(1) Galic S, Oakhill JS, Steinberg GR. Adipose tissue as an endocrine organ. *Mol Cell Endocrinol.* 2010 Mar 25;316(2):129-39.

(2) •• Soltis EE., Cassis LA. Influence of adipose tissue on rat aortic smooth muscle responsiveness. *Clinical & Experimental Hypertension – Part A . Theory & Practice* 1991; 13(2): 277-96. *first demonstration that PVAT might be important for vascular regulation*

(3) Lohn M, Dubrovskaja G, Lauterbach B, Luft FC, Gollasch M. Sharma AM. Periadventitial fat releases a vascular relaxing factor. *FASEB Journal* 2002; 16(9):1057-63.

(4) Dubrovskaja G, Verlohren S, Luft FC, Gollasch M. Mechanisms of ADRF release from rat aortic adventitial adipose tissue. *American Journal of Physiology – Heart & Circulatory Physiology* 2003; 286(3): H1107-13.

(5) Gao YJ, Lu C, Su LY, Sharma AM, Lee RMKW. Modulation of vascular function by perivascular adipose tissue: the role of endothelium and hydrogen peroxide. [comment]. *British Journal of Pharmacology* 2007; 151(3): 323-31.

(6) Gao Y-J, Zeng Z-H, Teoh K, Sharma AM, Abouzahr L, Cybulsky I, et al. Perivascular adipose tissue modulates vascular function in the human internal thoracic artery. *Journal of Thoracic & Cardiovascular Surgery* 2005; 130(4): 1130-6

(7) Archer SL, Gragasin FS, Wu X, Wang S, McMurtry S, Kim DH, et al. Endothelium-derived hyperpolarizing factor in human internal mammary artery is 11,12- epoxyeicosatrienoic acid and causes relaxation by activating smooth muscle BK (Ca) channels. *Circulation* 2003; 107(5): 769-76.

(8) Verlohren S, Dubrovskaja G, Tsang S-Y, EEssin K, Luft FC, Huang Y, et al. Visceral periadventitial adipose tissue regulates arterial tone of mesenteric arteries [comment]. *Hypertension* 2004; 44(3): 271-6.

(9) Gálvez B, de Castro J, Herold D, Dubrovskaja G, Arribas S, González MC, Arangué I, Luft FC, Ramos MP, Gollasch M, Fernández Alfonso MS. Perivascular adipose tissue and mesenteric vascular function in spontaneously hypertensive rats. *Arterioscler Thromb Vasc Biol.* 2006 Jun;26(6):1297-302.

(10) Baker AR, Silva NF, Quinn DW, Harte AL, Pagano D, Bonser RS, Kumar S, McTernan PG. Human epicardial adipose tissue expresses a pathogenic profile of

adipocytokines in patients with cardiovascular disease. *Cardiovasc Diabetol*. 2006 Jan 13;5:1.

(11) Quehenberger P, Exner M, Sunder-Plassmann R, Ruzicka K, Bieglmayer C, Endler G, et al. Leptin induces endothelin-1 in endothelial cells in vitro. *Circ Res*. 2002;90(6):711–8.

(12) Korda M, Kubant R, Patton S, Malinski T. Leptin-induced endothelial dysfunction in obesity. *Am J Physiol Heart Circ Physiol*. 2008;295(4):H1514–21.

(13) Ketonen J, Shi J, Martonen E, Mervaala E. Periadventitial adipose tissue promotes endothelial dysfunction via oxidative stress in diet-induced obese C57Bl/6 mice. *Circ J*. 2010;74(7):1479–87.

(14) Yamamoto K, Kiyohara T, Murayama Y, Kihara S, Okamoto Y, Funahashi T, Ito T, Nezu R, Tsutsui S, Miyagawa JI, Tamura S, Matsuzawa Y, Shimomura I, Shinomura Y. Production of adiponectin, an anti-inflammatory protein, in mesenteric adipose tissue in Crohn's disease. *Gut*. 2005 Jun;54(6):789-96.

(15) Trujillo ME, Scherer PE. Adiponectin—journey from an adipocyte secretory protein to biomarker of the metabolic syndrome. *Journal of Internal Medicine* 2005;257(2):167-75.

(16) Chow W-S, Cheung BMY, Tso AWK, Xu A, Wat NMS, Fong CHY, et al. Hypoadiponectinemia as a predictor for the development of hypertension: a 5-year prospective study. [comment]. *Hypertension* 2007; 49(6): 1455-61.

(17) Li H-Y, Chiu Y-F, Hwu C-M, Sheu WH-H, Hung Y-J, Fujimoto W, et al. The negative correlation between plasma adiponectin and blood pressure depends on obesity: a family-based association study in SAPPHERE. [comment]. *American Journal of Hypertension* 2008; 21(4): 471-6.

(18) • Iwashima Y, Katsuya T, Ishikawa K, Ouchi N, Ohishi M, Sugimoto K, et al. Hypoadiponectinemia is an independent risk factor for hypertension. *Hypertension* 2004; 43(6): 1318-23. *This is one of the first study that describes the several protective properties of adiponectin against hypertension*

(19) Yamauchi T, Kamon J, Waki H, Imai Y, Shimozawa N, Hioki K, et al. Globular adiponectin protected ob/ob mice from diabetes and ApoE-deficient mice from atherosclerosis. *Journal of Biological Chemistry* 2003; 278(4): 2461-8.

(20) Yamauchi T, Kamon J, Waki H, Terauchi Y, Kubota N, Hara K, et al. The fat-derived hormone adiponectin reverses insulin resistance associated with both lipodystrophy and obesity. [comment]. *Nature Medicine* 2001; 7(8): 941-6.

(21) Deng G, Long Y, Yu YR, Li MR (2010). Adiponectin directly improves endothelial dysfunction in obese rats through the AMPK-eNOS Pathway. *Int J Obes (Lond)* 2010, 34: 165–171

- (22) Luo N, Liu J, Chung BH, Yang Q, Klein RL, Garvey WT, et al. Macrophage adiponectin expression improves insulin sensitivity and protects against inflammation and atherosclerosis. *Diabetes*. 2010;59(4):791–9.
- (23) Li FY, Cheng KK, Lam KS, Vanhoutte PM, Xu A. Cross-talk between adipose tissue and vasculature: role of adiponectin. *Acta Physiol (Oxf)*. 2011;203(1):167–80
- (24) Cao Y, Tao L, Yuan YX, Jiao XY, Lau WB, Wang YJ, et al. Endothelial dysfunction in adiponectin deficiency and its mechanisms involved. *J Mol Cell Cardiol*. 2009;46(3):413–9.
- (25) Sowers JR. Endocrine functions of adipose tissue: focus on adiponectin. *Clin Cornerstone* 2008, 9: 32–38, discussion 39–40
- (26) Iacobellis G, Pistilli D, Gucciardo M, Leonetti F, Miraldi F, Brancaccio G, et al. Adiponectin expression in human epicardial adipose tissue in vivo is lower in patients with coronary artery disease. *Cytokine*. 2005;29(6):251–5.
- (27) •• Greenstein AS, Khavandi K, Withers SB, Sonoyama K, Clancy O, Jeziorska M, Laing I, Yates AP, Pemberton PW, Malik RA, Heagerty AM. Local inflammation and hypoxia abolish the protective anticontractile properties of perivascular fat in obese patients. *Circulation* 2009; 119(12): 1661-70. *This is the first demonstration of anticontractile effect of perivascular adipose tissue in obese patients*

- (28) An SJ, Boyd R, Wang Y, Qiu X, Wang HD. Endothelin-1 expression in vascular adventitial fibroblasts. *Am J Physiol Heart Circ Physiol* 2006;290: H700-H708
- (29) Berg AH, Scherer PE. Adipose tissue, inflammation, and cardiovascular disease. *Circ Res.* 2005; 96: 939–949.
- (30) Wellen KE, Hotamisligil GS. Inflammation, stress, and diabetes. *J Clin Invest.* 2005; 115: 1111–1119.
- (31) Despres JP, Lemieux I. Abdominal obesity and metabolic syndrome. *Nature.* 2006; 444: 881–887.
- (32) Lazar MA. How obesity causes diabetes: not a tall tale. *Science.* 2005; 307: 373–375.
- (33) Lu C, Zhao AX, Gao YJ, Lee RM. Modulation of vein function by perivascular adipose tissue. *Eur J Pharmacol.* 2011 Apr 25;657(1-3):111-6.
- (34) Malinowski M, Deja MA, Gołba KS, Roleder T, Biernat J, Woś S. Perivascular tissue of internal thoracic artery releases potent nitric oxide and prostacyclin-independent anticontractile factor. *Eur J Cardiothorac Surg.* 2008 Feb;33(2):225-31.
- (35) Lee RM, Ding L, Lu C, Su LY, Gao YJ. Alteration of perivascular adipose tissue function in angiotensin II-induced hypertension. *Can J Physiol Pharmacol.* 2009 Nov;87(11):944-53

- (36) Lee RMKW, Lu C, Su LY, Gao YJ. Endothelium-dependent relaxation factor released by perivascular adipose tissue. *J Hypertens*. 2009;27(4):782–90.
- (37) • Engeli S, Gorzelniak K, Kreutz R, Runkel N, Distler A, Sharma AM. Co-expression of renin-angiotensin system genes in human adipose tissue. *Journal of Hypertension* 1999; 17: 555-560. *The study describes the role of the renin-angiotensin system in human adipose tissue*
- (38) Gálvez-Prieto B, Bolbrinker J, Stucchi P, de Las Heras AI, Merino B, Arribas S, Ruiz-Gayo M, Huber M, Wehland M, Kreutz R, Fernandez-Alfonso MS. Comparative expression analysis of the renin-angiotensin system components between white and brown perivascular adipose tissue. *Journal of Endocrinology* 2008; 197: 55-64.
- (39) Goossens GH, Jocken JW, Blaak EE, Schiffrers PM, Saris WH, van Baak MA. Endocrine role of the renin-angiotensin system in human adipose tissue and muscle: effect of beta-adrenergic stimulation. *Hypertension* 2007 Mar;49(3):542-7.
- (40) • Perivascular adipose tissue and local Renin-Angiotensin System. From Cassis LA, Police SB, Yiannikouris F, Thatcher SE. Local adipose tissue renin-angiotensin system. *Curr Hypertens Rep* 2008; 10(2): 93-98. *This study is a review of the important role of adipose tissue renin-angiotensin system*

41) Campbell DJ, Habener JF. Cellular localization of *AGT* gene expression in brown adipose tissue and mesentery: quantification of messenger ribonucleic acid abundance using hybridization in situ. *Endocrinology* 1987; 121: 1616-1626

(42) Cassis LA, Lynch KR, Peach MJ. Localization of angiotensinogen messenger RNA in rat aorta. *Circulation Research* 1988a; 62: 1259-1262

(43) Shiling P, Mallow H, Trindl A, Loffler G. Evidence for a local renin angiotensin system in primary cultured human preadipocytes. *Internal Journal Obes Relat Metab Disord.* 1999; 23: 336-341.

(44) Saye J, Linch KR, Peach MJ. Changes in angiotensinogen messenger RNA in differentiating 3T3- F442A adipocytes. *Hypertension* 1990; 15: 867-871

(45) Hainault I, Nebout G, Ardouin BB, Quignard-Boulangé A. Developmental changes in *AGT* expression and its secretion in the Zucker rat: adipose tissue-specific effect of *FA* genotype. *Int J Obes Relat Metab Disord.* 1998; 22(suppl 3): S103 Abstract.

(46) Giacchetti G, Faloia E, Sardu C, Mariniello B, Garrapa GGM, Gatti C, Camilloni MA, Mantero F. Different gene expression of the RAS in human subcutaneous and visceral adipose tissue. *Int J Obes Relat Metab Disord.* 1999;23(suppl 5): S71. Abstract.

- (47) Phillips MI, Speakman EA, Kimura B. Levels of angiotensin and molecular biology of the tissue renin angiotensin systems. *Regulatory Peptides* 1993; 43: 1-20.
- (48) Shenoy U, Cassis L. Characterization of renin activity in brown adipose tissue. *Am J Physiol.* 1997; 272: C989- C999.
- (49) Kupiers,I., van der Harst, P., Navis, G., et al. Nuclear hormone receptors as regulators of the renin-angiotensin-aldosterone system. *Hypertension* 2008; 51: 1442-1448.
- (50) Archard V, Boullu-Ciocca S, Desbriere R, Nguyen G, Grino M. Renin receptor expression in human adipose tissue. *American Journal of Physiology, Regulatory, Integrative and Comparative Physiology* 2007; 292: 274-282.
- (51) Touyz RM, Schiffrin EL. Signal transduction mechanisms mediating the physiological and pathophysiological actions of angiotensin II in vascular smooth muscle cells. *Pharmacol Rev* 2000; 52: 639-672
- (52) Schling P, Mallow H, Trindl A, Löffler G. Evidence for a local renin angiotensin system in primary cultured human preadipocytes.. *Int J Obes Relat Metab Disord.* 1999 Apr;23(4):336-41.
- (53) Sharma AM, Janke J, Gorzelniak K, Engeli S, Luft FC. Angiotensin blockade prevents type 2 diabetes by formation of fat cells. *Hypertension* 2002; 40: 609-611.

(54) Matsushita K, Wu Y, Okamoto Y, Pratt RE, Dzau VJ. Local renin-angiotensin expression regulates Human mesenchymal stem cell differentiation to adipocytes. *Hypertension* 2006; 48: 1095-1102.

(55) Jones BH, Standridge MK, Moustaid N. Angiotensin II increases lipogenesis in 3T3-L1 and human adipose cells. *Endocrinology* 1997; 138(4): 1512-1519.

(56) Furuhashi M, Ura N, Higashiura K, et al. Blockade of the renin-angiotensin system increases adiponectin concentrations in patients with essential hypertension. *Hypertension*. 2003;42:76-81

(57) Lee MH, Song HK, Ko GJ, et al. Angiotensin receptor blockers improve insulin resistance in type 2 diabetic rats by modulating adipose tissue. *Kidney Int*. 2008;74:890-900

(58) Rinaldi B, Di Filippo C, Capuano A, Donniacuo M, Sodano L, Ferraraccio F, Rossi F, D'Amico M. Adiponectin elevation by telmisartan ameliorates ischaemic myocardium in Zucker diabetic fatty rats with metabolic syndrome. *Diabetes Obes Metab*. 2012 Apr;14(4):320-8.

(59) Nakamura T, Kawachi K, Saito Y, Saito T, Morishita K, Hoshino J, Hosoi T, Iwasaki T, Ohyama Y, Kurabayashi M. Effects of ARB or ACE-inhibitor administration on plasma levels of aldosterone and adiponectin in hypertension. *Int Heart J*. 2009;50:501-512.

- (60) Ran J, Hirano T, Fukui T, Saito K, Kageyama H, Okada K, Adachi M. Angiotensin II infusion decreases plasma adiponectin level via its type 1 receptor in rats: an implication for hypertension-related insulin resistance. *Metabolism*. 2006;55:478–488.
- (61) Guo B, Li Y, Han R, Zhou H, Wang M. Angiotensin II upregulation of cardiomyocyte adiponectin production is nitric oxide/cyclic GMP dependent. *Am J Med Sci*. 2011 May;341(5):350-5.
- (62) Onat A. Metabolic syndrome: nature, therapeutic solutions and options. *Expert Opin Pharmacother*. 2011 Aug;12(12):1887-1900.
- (63) Okada S, Kozuka C, Masuzaki H, Yasue S, Ishii-Yonemoto T, Tanaka T, Yamamoto Y, Noguchi M, Kusakabe T, Tomita T, Fujikura J, Ebihara K, Hosoda K, Sakaue H, Kobori H, Ham M, Lee YS, Kim JB, Saito Y, Nakao K. Adipose tissue-specific dysregulation of angiotensinogen by oxidative stress in obesity. *Metabolism Journal* 2010; 59(9): 1241-1251.
- (64) Farquharson CA, Struthers AD. Aldosterone induces acute endothelial dysfunction in vivo in humans: evidence for an aldosterone-induced vasculopathy. *Clin Sci (Lond)* 2002; 103(4): 425-431.
- (65) Sun Y, Zhang J, Lu L, Chen SS, Quinn MT, Weber KT. Aldosterone-induced inflammation in the rat heart: role of oxidative stress. *The American Journal of Pathology* 2002; 161(5): 1773-1781.

- (66) Hirata A, Maeda N, Hiuge A, Hibuse T, Fujita K, Okada T, Kihara S, Funahashi T, Shimomura I. Blockade of mineralcorticoid receptor reverses adipocyte dysfunction and insulin resistance in obese mice. *Cardiovascular Research* 2009.
- (67) Flynn C, Bakris GL. Interaction between Adiponectin and Aldosterone. *Cardiorenal Med.* 2011;1(2):96-101.
- (68) Lemarié CA, Paradis P, Schiffrin EL. New insights on signaling cascades induced by cross-talk between angiotensin II and aldosterone. *J Mol Med* 2008; 86(6): 673-678.
- (69) Lu C, Su LY, Lee RM, Gao YJ. Mechanisms for perivascular adipose tissue-mediated potentiation of vascular contraction to perivascular neuronal stimulation: the role of adipocyte-derived angiotensin II. *Eur J Pharmacol.* 2010 May 25;634(1-3):107-12.
- (70) Lu C, Su LY, Lee RM, Gao YJ. Alterations in perivascular adipose tissue structure and function in hypertension. *Eur J Pharmacol.* 2011 Apr 10;656(1-3):68-73.
- (71) •• Withers SB, Agabiti Rosei C, Livingstone DM, Little MC, Aslam R, Malik RA, Heagerty AM. Macrophage activation is responsible for loss of anticontractile function in inflamed perivascular fat. *Artheroscler Thromb, Vascular, Biol* 2011

Apr;31(4):908-13. *This study demonstrates the involvement of macrophage activation in loss of anticontractile function of perivascular adipose tissue*

(72) • Rosei CA, Withers SB, Belcaid L, De Ciuceis C, Rizzoni D, Heagerty AM. Blockade of the renin-angiotensin system in small arteries and anticontractile function of perivascular adipose tissue. *J Hypertens*. 2015 May;33(5):1039-45. doi: 10.1097/HJH. *This is the first demonstration of restore anticontractile effect of perivascular fat with blockade of the renin-angiotensin system in small arteries*

(73) • Huang Cao ZF1, Stoffel E, Cohen P. Role of perivascular adipose tissue in vascular physiology and pathology. *Hypertension* 2017; 69:770-777. *This is a recent overview of the role of perivascular adipose tissue in vascular structure and reactivity*

(74) Canello R, Henegar C, Viguier N, Taleb S, Poitou C, Rouault C, et al. Reduction of macrophage infiltration and chemoattractant gene expression changes in white adipose tissue of morbidly obese subjects after surgery-induced weight loss. *Diabetes* 2005; 54(8): 2277-86.

(75) Chen B, Lam KSL, Wang Y, Wu D, Larn MC, Shen J, et al. Hypoxia dysregulates the production of adiponectin and plasminogen activator inhibitor-1 independent of reactive oxygenspecies in adipocytes. *Biochemical e Biophysical Research Communications* 2006; 341 (2): 549-56.

(76) Patten DA, Lafleur VN, Robitaille GA, Chan DA, Giaccia AJ, Richard DE. Hypoxia-inducible factor-1 activation in nonhypoxic conditions: the essential role of mitochondrial-derived reactive oxygen species. *Mol Biol Cell* 2010; 21(18): 3247-57.

(77) Elisabeth L. Pagé, Geneviève A Robitaille, Pouysségur J, Richard DE. Induction of hypoxia-inducible factor-1 α by transcriptional and translational mechanisms. *J Biol Chem* 2002; 277(50): 48403-9.

(78) • Schnabel R, Larson MG, Dupuis J, Lunetta KL, Lipinska I, Meigs JB, Yin X, Rong J, Vita JA, Newton-Cheh C, Levy D, Keaney JF Jr, Vasani RS, Mitchell GF, Benjamin EJ. Relations of inflammatory biomarkers and common genetic variants with arterial stiffness and wave reflection. *Hypertension*. 2008 Jun;51(6):1651-7.

Source National Heart Lung and Blood Institute Framingham Study, Framingham, Mass., USA. *This study shows that inflammation and oxidative stress are interrelated and associated with adipose tissue dysfunction and they could contribute to arterial stiffness and hypertension*

(79) Chae CU, Lee RT, Rifai N, Ridker PM. Blood pressure and inflammation in apparently healthy men. *Hypertension*. 2001 Sep;38(3):399-403.

(80) Furukawa S, Fujita T, Shimabukuro M, Iwaki M, Yamada Y, Nakajima Y, Nakayama O, Makishima M, Matsuda M, Shimomura I. Increased oxidative stress in obesity and its impact on metabolic syndrome. *J Clin Invest*. 2004 Dec;114(12):1752-61.

(81) Kawanami D, Maemura K, Takeda N, Harada T, Nojiri T, Imai Y, Manabe I, Utsunomiya K, Nagai R. Direct reciprocal effects of resistin and adiponectin on vascular endothelial cells: a new insight into adipocytokine-endothelial cell interactions. *Biochem Biophys Res Commun*. 2004 Feb 6;314(2):415-9.

(82) • Lynch FM1, Withers SB, Yao Z, Werner ME, Edwards G, Weston AH, Heagerty AM. Perivascular adipose tissue-derived adiponectin activates BK(Ca) channels to induce anticontractile responses. *Am J Physiol Heart Circ Physiol*. 2013 Mar 15;304(6):H786-95. *This study is the first demonstration that the mechanism underlies the anticontractile effect of perivascular adipose tissue involves the activation of BK(Ca) channels*

(83) Weston AH, Egner I, Dong Y, Porter EL, Heagerty AM, Edwards G. Stimulated release of a hyperpolarizing factor (ADHF) from mesenteric artery perivascular adipose tissue: involvement of myocyte BKCa channels and adiponectin. *Br J Pharmacol*. 2013 Aug;169(7):1500-9

(84) • Withers SB, Simpson L, Fattah S, Werner ME, Heagerty AM. cGMP-dependent protein kinase (PKG) mediates the anticontractile capacity of perivascular adipose tissue. *Cardiovasc Res*. 2014 Jan 1;101(1):130-7. *This study shows the role of cGMP-dependent protein kinase (PKG) as a principal mediators of perivascular adipose tissue activity*

(85) Agabiti-Rosei C, De Ciuceis C, Rossini C, Porteri E, Rodella LF, Withers SB, Heagerty AM, Favero G, Agabiti-Rosei E, Rizzoni D, Rezzani R. Anticontractile

activity of perivascular fat in obese mice and the effect of long-term treatment with melatonin. *J. Hypertens.* 2014 Jun;32(6):1264-74

(86) Bonnefont-Rousselot D1, Collin F, Jore D, Gardès-Albert M. Reaction mechanism of melatonin oxidation by reactive oxygen species in vitro. *J Pineal Res.* 2011 Apr;50(3):328-35.

(87) Rezzani R, Porteri E, De Ciuceis C, Bonomini F, Rodella LF, Paiardi S, Boari GE, Platto C, Pilu A, Avanzi D, Rizzoni D, Agabiti Rosei E. Effects of melatonin and Pycnogenol on small artery structure and function in spontaneously hypertensive rats. *Hypertension.* 2010 Jun;55(6):1373-80.

(88) Koziróg M1, Poliwczak AR, Duchnowicz P, Koter-Michalak M, Sikora J, Broncel M. Melatonin treatment improves blood pressure, lipid profile, and parameters of oxidative stress in patients with metabolic syndrome. *J Pineal Res.* 2011 Apr;50(3):261-6

(89) Ríos-Lugo MJ1, Cano P, Jiménez-Ortega V, Fernández-Mateos MP, Scacchi PA, Cardinali DP, Esquifino AI. Melatonin effect on plasma adiponectin, leptin, insulin, glucose, triglycerides and cholesterol in normal and high fat-fed rats. *J Pineal Res.* 2010 Nov;49(4):342-8.

(90) • Tan DX, Manchester LC, Fuentes-Broto L, Paredes SD, Reiter RJ. Significance and application of melatonin in the regulation of brown adipose tissue metabolism: relation to human obesity. *Obes Rev.* 2011 Mar;12(3):167-88. *This study describes*

melatonin function in brown adipose tissue metabolism suggesting a potential target for treatment of obesity in humans

(91) Takeda T. Senescence-accelerated mouse (SAM): a biogerontological resource in aging research. *Neurobiol Aging*. 1999 Mar-Apr;20(2):105-10.

(92) Butterfield DA, Poon HF. The senescence-accelerated prone mouse (SAMP8): a model of age-related cognitive decline with relevance to alterations of the gene expression and protein abnormalities in Alzheimer's disease. *Exp Gerontol*. 2005 Oct;40(10):774-83.

(93) Agabiti-Rosei C, Favero G, De Ciuceis C, Rossini C, Porteri E, Rodella LF, Franceschetti L, Maria Sarkar A, Agabiti-Rosei E, Rizzoni D, Rezzani R. Effect of long-term treatment with melatonin on vascular markers of oxidative stress/inflammation and on the anticontractile activity of perivascular fat in aging mice. *Hypertens Res*. 2017 Jan;40(1):41-50.

(94) Narkiewicz K. Diagnosis and management of hypertension in obesity. *Obes Rev*. 2006 May;7(2):155-62.

(95) Kotchen TA. Obesity-related hypertension: epidemiology, pathophysiology, and clinical management. *Am J Hypertens*. 2010 Nov;23(11):1170-8.

(96) Mancia G, De Backer G, Dominiczak A, Cifkova R, Fagard R, Germano G, Grassi G, Heagerty AM, Kjeldsen SE, Laurent S, Narkiewicz K, et al. 2007

Guidelines for the management of arterial hypertension: The Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *Eur Heart J.* 2007 Jun;28(12):1462-536.

(97) Guo Z, Zheng C, Qin Z, Wei P. Effect of telmisartan on the expression of cardiac adiponectin and its receptor 1 in type 2 diabetic rats. *J Pharm Pharmacol.* 2011 Jan;63(1):87-94.