



University of  
**Salford**  
MANCHESTER

# Standardization and methodological considerations for the isometric mid-thigh pull

Comfort, P, Dos'Santos, T, Beckham, G, Stone, M, Guppy, S and Haff, GG

<http://dx.doi.org/10.1519/SSC.0000000000000433>

<b>Title</b>	Standardization and methodological considerations for the isometric mid-thigh pull
<b>Authors</b>	Comfort, P, Dos'Santos, T, Beckham, G, Stone, M, Guppy, S and Haff, GG
<b>Publication title</b>	Strength and Conditioning Journal
<b>Publisher</b>	Lippincott, Williams & Wilkins (Wolters Kluwer Health)
<b>Type</b>	Article
<b>USIR URL</b>	This version is available at: <a href="http://usir.salford.ac.uk/id/eprint/48647/">http://usir.salford.ac.uk/id/eprint/48647/</a>
<b>Published Date</b>	2019

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: [library-research@salford.ac.uk](mailto:library-research@salford.ac.uk).

1     **Standardization and methodological considerations for the Isometric Mid-Thigh Pull**

2     Paul Comfort<sup>1#</sup>, Thomas Dos'Santos<sup>1</sup>, George K. Beckham<sup>2</sup>, Michael H. Stone<sup>3</sup>, Stuart. N.  
3     Guppy<sup>4</sup>, G. Gregory Haff<sup>4</sup>.

4     <sup>1</sup> Directorate of Sport, Exercise and Physiotherapy, University of Salford, Salford, Greater  
5     Manchester, UK

6     <sup>2</sup> Kinesiology Department, California State University, Monterey Bay, Seaside, CA.

7     <sup>3</sup> Center of Excellence for Sport Science and Coach Education, Department of Exercise and  
8     Sport Science, East Tennessee State University, Johnson City, TN.

9     <sup>4</sup> Centre for Exercise and Sports Science Research, Edith Cowan University, Joondalup,  
10    Australia

11    #Corresponding Author – [p.comfort@salford.ac.uk](mailto:p.comfort@salford.ac.uk)

12

13

14

15

16

17

18

19

20

21

22

23

## 24 **Standardization and methodological considerations for the Isometric Mid-Thigh Pull**

25

### 26 **Abstract**

27 The isometric mid-thigh pull (IMTP) is commonly used to assess an athlete's force generation  
28 ability. This test is highly reliable and is simple and relatively quick to perform. The data that  
29 can be determined from the force-time curves generated by the test have been shown to be  
30 closely related to performance capacities in a variety of dynamic athletic tasks. However,  
31 within the scientific literature there are inconsistencies in the data collection procedures and  
32 methods used for data analysis that may impact the resultant output and the ability to compare  
33 and generalize results. Therefore, the primary aim of this review is to identify the differences  
34 in IMTP testing procedures and data analysis techniques, while identifying the potential impact  
35 this may have on the data collected. The secondary aim is to provide recommendations for  
36 the standardization of testing procedures to ensure that future IMTP data is of maximal benefit  
37 to practitioners and researchers.

38

### 39 **Introduction**

40 Maximal strength underpins performance in many athletic tasks (15, 55, 63) and as such,  
41 monitoring strength, usually via repetition maximum (RM) testing, is commonly performed by  
42 practitioners and researchers. While RM testing is reliable (12, 24, 28), it can be perceived as  
43 fatiguing, posing an increased potential for injury risk, and only providing information related  
44 to the maximal load lifted. In contrast, isometric testing, such as the isometric mid-thigh pull  
45 (IMTP), is potentially safer (18), less fatiguing, and allows for the quantification of peak force  
46 (PF), force at a variety of epochs, and can provide several measures of the rate of force  
47 development (RFD) (11, 21, 26, 30, 32, 33). The diagnostic ability of these measures may be  
48 of importance when considering time constrained tasks within sports, such as jumping,

49 sprinting and change of direction. Importantly, the IMTP has been shown to be highly reliable  
50 both within and between sessions, with low variability and low measurement error (8, 11, 18,  
51 24, 26, 27, 32).

52 Performance in the IMTP has been associated with performance in numerous athletic tasks  
53 (7, 18, 30, 33, 40, 41, 45, 46, 49, 59, 64, 66, 67, 69, 72, 73). Specifically, absolute PF has  
54 been associated with weightlifting performance (7, 30), 1RM squat and power clean (45-47,  
55 49, 59, 69, 73), 1RM deadlift (18), vertical jump performance (39-41, 53, 60, 64, 67), short  
56 sprint and change of direction times (59, 64), sprint cycling performance (60), and throwing  
57 performance (72) (Table 1). In contrast, West et al. (71) reported no meaningful relationships  
58 between absolute PF and short sprint times or jump height, although they did observe large  
59 correlations between relative PF (PF/body weight) and these variables in rugby league  
60 players. Similarly, Nuzzo et al. (49) reported only a small relationship between absolute PF  
61 and jump height but a large relationship between relative PF and jump height (Table 1). The  
62 range of associations between PF and performance in other tasks is summarised in Figure 1.  
63 Researchers have also reported relationships between allometrically scaled PF and  
64 performance in athletic tasks (60, 72), demonstrating similar correlations to those observed  
65 when ratio scaling is used (60).

66

67

68

69

70


71

72

**Table 1: Relationships between peak force and performance in other activities**

Author(s)	Subjects	1RM	Sprint	Jump	Change of Direction	Other
Haff et al. (39)	8 trained (>2 years) men 1RM PC = 1.21 kg.kg <sup>-1</sup>			SJ PF: r = 0.76		Force during dynamic MTP 90% 1RM: r = 0.77 100% 1RM: r = 0.80
Stone et al. 60	30 competitive sprint cyclists			CMJ height: r = 0.59 CMJ PP: r = 0.79 SJ height: r = 0.51 SJ PP: r = 0.78		Absolute PF & Sprint cycling performances: r = 0.49-0.55 Relative PF & Sprint cycling performances: r = 0.45-0.60 AS PF & Sprint cycling performances: r = 0.45-0.58
Haff et al. (30)	6 elite women weightlifters	Snatch: r = 0.93		CMJ PP: r = 0.88 SJ PP: r = 0.92		
Kawamori et al. (39)	8 male collegiate weightlifters 1RM PC = 1.39 kg.kg <sup>-1</sup>			CMJ PF: r = 0.87 CMJ PRFD: r = 0.85 CMJ PP: r = 0.95 CMJ height: r = 0.82 SJ height: r = 0.87		Force during dynamic MTP 90% 1RM: r = 0.82
McGuigan et al. (47)	8 division III collegiate wrestlers	PC: r = 0.97 Squat: r = 0.96 BP: r = 0.73				
McGuigan & Winchester (45)	22 college football players 1RM PC = 1.11 kg.kg <sup>-1</sup> 1RM Squat = 1.75 kg.kg <sup>-1</sup>	PC, Squat, BP: r = 0.61-0.72*				
Nuzzo et al. (49)	12 division I collegiate athletes 1RM PC = 1.28 kg.kg <sup>-1</sup> 1RM Squat = 1.91 kg.kg <sup>-1</sup>	PC: r = 0.74		CMJ PP: r = 0.75 Relative PF & CMJ height: r = 0.59		
Kraska et al. (41)	41 female and 22 male collegiate athletes			SJ: r = 0.40 SJ20: r = 0.55 CMJ: r = 0.36 CMJ20: r = 0.55 AS PF: SJ: r = 0.47		

				SJ20: r = 0.52 CMJ: r = 0.41 CMJ20: r = 0.52		
Whittington et al. (72)	7 NCAA Division I track and field athletes					Ball throw distance PF: r = 0.89 AS PF: r = 0.91
McGuigan et al. (46)	26 recreationally trained men 1RM Squat = 1.30 kg•kg <sup>-1</sup>	Squat: r = 0.97 BP: r = 0.99			CMJ height: r = 0.72	
Khamoui et al. (40)	19 recreationally trained men			Relative PF & CMJ height: r = 0.61		Relative PF & high pull PV: r = -0.60
West et al. (71)	39 professional rugby league players		Relative PF & 10 m sprint time: r = 0.37	Relative PF & CMJ height: r = 0.45		
Spiteri et al. (59)	12 competitive female basketball players	IMTP relative PF, back squat: r = 0.81			T-Test: r = -0.85 505 COD = -0.79	
Winchester et al. (73)	26 recreationally trained men 1RM Squat = 1.30 kg•kg <sup>-1</sup>	Squat: r = 0.97 BP: r = 0.99			CMJ height: r = 0.72	
Secomb et al. (53)	15 elite surfers				CMJ height: r = 0.65 SJ height: r = 0.58	
Beckham et al. (7)	12 collegiate-national level weightlifters	Snatch: r = 0.83 Clean & Jerk: r = 0.84 Total: r = 0.84				
Thomas et al. (64)	14 collegiate team sport athletes		5 m: r = -0.57 20 m: r = -0.69			505mod: r = -0.57
Thomas et al. (67)	22 collegiate team sport athletes				CMJ PF: r = 0.45	
Wang et al. (69)	15 collegiate rugby players	Squat: r = 0.866				
PC = Power Clean; BP = Bench Press; SJ = Squat Jump; CMJ = Countermovement Jump; 505mod = Modified 505 change of direction PF = Peak Force; PP = Peak Power; PV = Peak Velocity; PRFD = Peak Rate of Force Development; AS = Allometrically Scaled *Individual correlations not reported						



76

77 Figure 1: Relationships between isometric mid-thigh pull peak force and performance in other tasks (References  
78 in Table 1)

79

80 Another way to examine the isometric force-time curve is to measure force at specific time  
81 epochs (e.g. 50-250 ms). It has been reported that these time specific forces are associated  
82 with squat jump (SJ) and countermovement jump (CMJ) height (force at 50-, 90, 250 ms) (41),  
83 weightlifting performance (force at 100-, 150-, 200-, 250 ms) (7) and 1RM back squat (90-250  
84 ms) (69). Additionally, allometrically scaled force at 150 ms was reported to be related to mean  
85 and maximum club head speed during a golf swing (42), with allometrically scaled force at 50-  
86 , 90- and 250 ms also related to jump performance (41) (Table 2). In contrast, however, force  
87 at 30-250 ms was not related to 1RM deadlift performance (18).

88

89

90

91

92 **Table 2: Relationships between time specific force and performance in other activities**

93

Author(s)	Subjects	1RM	Sprint	Jump	Other
Kraska et al. (41)	41 female and 22 male collegiate athletes			PF50 SJ: r = 0.33 SJ20: r = 0.52 CMJ: r = 0.27 CMJ20: r = 0.50 AS PF50: SJ: r = 0.33 SJ20: r = 0.48 CMJ20: r = 0.45 PF90 SJ20: r = 0.37 CMJ20: r = 0.33 AS PF90: CMJ20: r = 0.48 PF250 SJ: r = 0.39 SJ20: r = 0.56 CMJ: r = 0.34 CMJ20: r = 0.54 AS PF250 SJ: r = 0.42 SJ20: r = 0.51 CMJ: r = 0.34 CMJ20: r = 0.48	
Beckham et al. (7)	12 collegiate-national level weightlifters	F100 Snatch: r = 0.65 Clean & Jerk: r = 0.64 Combined Total: r = 0.65 F150 Snatch: r = 0.64 Clean & Jerk: r = 0.61 Combined Total: r = 0.62 F200 Snatch: r = 0.73 Clean & Jerk: r = 0.71			



		Combined Total: $r = 0.72$ F250 Snatch: $r = 0.80$ Clean & Jerk: $r = 0.80$ Combined Total: $r = 0.80$			
West et al. (71)	39 professional rugby league players		F100 & 10 m: $r = -0.66$ Relative F100 & 10 m: $r = -0.68$	F100 & CMJ PP: $r = 0.55$ Relative F100 & CMJ PP: $r = 0.38$ Relative F100 & CMJ height: $r = 0.43$	
Wang et al. (69)	15 collegiate rugby players	Squat F90: $r = 0.76$ F100: $r = 0.78$ F150: $r = 0.78$ F200: $r = 0.77$ F250: $r = 0.82$			
Leary et al. (42)	12 recreational golfers				<i>Golf Club Head Speed</i> ASF150 & Mean Club Head Speed: $r = 0.46$ ASF150 & Max' Club Head Speed: $r = 0.47$
F90 = Force at 90 ms; F100 = Force at 100 ms; F150 = Force at 150 ms; F200 = Force at 200 ms; F250 = Force at 250 ms AS = Allometrically Scaled; SJ20 = Squat Jump with 20 kg; CMJ20 = Countermovement Jump with 20 kg					

94  
95

96

97 Equivocal results regarding the relationships between measures of RFD and performance in dynamic athletic tasks have been reported in the  
98 scientific literature. When examining how the RFD is quantified two main methods exist within the literature (32). The first method is to quantify  
99 the peak RFD (PRFD) that occurs during the IMTP with a predefined moving window, most typically lasting between 2-40 ms (32) (Table 3).  
100 When this method is utilized for analyzing the force-time curve conflicting results exist within the scientific literature with some authors reporting

101 significant relationships between the RFD and dynamic performance activities (30, 33, 39, 41),  
102 while others report no meaningful relationship with 1RM performance (7, 45-47), or SJ and  
103 CMJ performances (40, 49, 67). These difference may be attributable to the moving window,  
104 with Maffiuletti et al. (43) cautioning against the use of short windows (e.g. 2 ms) as they may  
105 be too sensitive to unsystematic variability and therefore less reliable. The second method for  
106 evaluating the RFD is to examine time dependant epochs (32). The use of time dependent  
107 epoch has been shown to be an effective method for examining the RFD during the IMTP and  
108 relating it to various sports performance tasks. For example, Spiteri et al. (58) report that  
109 athletes who produce higher RFD to 90 ms and 100 ms are able to demonstrate faster agility  
110 times during a 45 ° cutting task. One possible explanation why some RFD measures relate to  
111 dynamic performance activities and others do not is the method of calculation and reliability of  
112 the method. For example, Haff et al. (32) have shown that the only PRFD measure that is  
113 reliable is when a 20 ms moving window is used, supporting previous suggestions by  
114 Maffiuletti et al. (43). Conversely, using time dependent epochs such as 0-90 ms, 0-150 ms,  
115 0-200 ms and 0-250 ms to calculate the mean RFD across the specific duration produces  
116 much more reliable results and generally have better relationships to dynamic performance  
117 measures. Therefore, it is generally recommended that using time specific RFD epochs is  
118 warranted when using the IMTP as a performance diagnostic tool (32).

119

120

121

122

123

124

125

**Table 3: Relationships between RFD and performance in other activities**

Author(s)	Subjects	1RM	Sprint	Jump	Change of Direction	Other
Haff et al. (33)	8 trained (>2 years) men 1RM PC = 1.21 kg•kg <sup>-1</sup>			PRFD SJ Power: r = 0.76 SJ Height: r = 0.82		RFD during dynamic MTP 80% 1RM: r = 0.84 90% 1RM: r = 0.88 100% 1RM: r = 0.84
Haff et al. (30)	6 elite women weightlifters	PRFD Snatch: r = 0.79 Combined Total: r = 0.80		PRFD CMJ PP: r = 0.81 SJ PP: r = 0.84		
McGuigan et al. (47)	8 division III collegiate wrestlers					PRFD & Coaching Ranking: r = 0.62
Kawamori et al. (39)	8 male collegiate weightlifters 1RM PC = 1.39 kg•kg <sup>-1</sup>					Force during dynamic MTP 90% 1RM: r = 0.69 120% 1RM: r = 0.74
Nuzzo et al. (49)	12 division I collegiate athletes 1RM PC = 1.28 kg•kg <sup>-1</sup> 1RM Squat = 1.91 kg•kg <sup>-1</sup>			PRFD CMJ PP: r = 0.65		
Kraska et al. (41)	41 female and 22 male collegiate athletes			PRFD SJ: r = 0.48 SJ20: r = 0.66 CMJ: r = 0.43 CMJ20: r = 0.62		
Whittington et al. (72)	7 NCAA Division I track and field athletes					Ball throw distance: r = 0.78
Khamoui et al. (40)	19 recreationally trained men					RFD50 & high pull PV: r = 0.56 RFD100 & high pull PV: r = 0.56

West et al. (71)	39 professional rugby league players		PRFD 10 m: r = -0.66	PRFD CMJ height: r = 0.39		
Beckham et al. (7)	12 collegiate-national level weightlifters	RFD200 Snatch: r = 0.65 Combined Total: r = 0.60 RFD250 Snatch: r = 0.78 Clean & Jerk: r = 0.72 Combined Total: r = 0.75				
Thomas et al. (64)	14 collegiate team sport athletes		PRFD 5 m: r = -0.58 20 m: r = 0.71		PRFD 505mod: r = -0.57	
Wang et al. (69)	15 collegiate rugby players		5 m: PRFD: r = -0.54 RFD30: r = 0.57 RFD50: r = 0.53		Pro agility: PRFD: r = -0.52 RFD30: r = 0.52 RFD50: r = 0.53 RFD90: r = 0.53 RFD100: r = 0.52	
PRFD = Peak RFD; RFD30 = Mean RFD between 0-30 ms; RFD50 = Mean RFD between 0-50 ms; RFD90 = Mean RFD between 0-90 ms RFD100 = Mean RFD between 0-100 ms; RFD200 = Mean RFD between 0-200 ms; RFD250 = Mean RFD between 0-250 ms; PV = Peak Velocity						

128

129

130

131

132

133

Another method for analysing the force-time curve derived from an IMTP is to examine the isometric impulse (67, 68). For example, impulse values across different epochs (0-100, 0-200 and 0-300 ms) have been associated with 5- and 20 m sprint times as well as 505 change of direction times (64), peak force and power during the SJ and CMJ (68) (Table 4). While determining the isometric impulse of various epochs within the force-time curve achieved during the IMTP yields useful information much more research is needed to understand how best to utilise this measurement in a sports performance monitoring program.

134  
135

**Table 4: Relationships between time specific impulse and performance in other activities**

Author(s)	Subjects	Sprint	Jump	Change of Direction
Thomas et al. (64)	14 collegiate team sport athletes	Imp100 5 m: r = -0.71 20 m: r = 0.75 Imp300 5 m: r = -0.74 20 m: r = 0.78		Imp100, 505mod: r = -0.58 Imp300, 505mod: r = -0.62
Thomas et al. (67)	22 collegiate team sport athletes		Imp100 SJ PF: r = 0.57 SJ PP: r = 0.60 CMJ PF: r = 0.64 CMJ PP: r = 0.51 Imp200 SJ PF: r = 0.56 SJ PP: r = 0.59 CMJ PF: r = 0.63 CMJ PP: r = 0.50 Imp300 SJ PF: r = 0.58 SJ PP: r = 0.60 CMJ PF: r = 0.63 CMJ PP: r = 0.49	
Imp100 = Impulse over 100 ms; Imp200 = Impulse over 200 ms; Imp300 = Impulse over 300 ms SJ = Squat Jump; CMJ = Countermovement Jump; PF = Peak Force; PP = Peak Power				

136  
137

138

139 The PF achieved during the IMTP has also been used to monitor adaptations to training (5,  
140 36, 50, 51, 57, 70, 74), with some authors also including RFD (36, 51, 52, 74). PF and peak  
141 RFD have also been used in an attempt to identify levels of fatigue or recovery (4, 29, 35, 44).  
142 More recently researchers have started to investigate the potential of the IMTP to investigate  
143 between-limb asymmetries, using dual force platforms (1-3) and a unilateral stance IMTP (25,  
144 65). Additionally, the PF during the IMTP has been divided by the PF during a SJ or CMJ, to  
145 calculate the dynamic strength index (DSI; ratio of PF during the CMJ or SJ and IMTP PF), in  
146 attempt to identify if an athlete needs to focus more on maximal force production or rapid  
147 dynamic force production (14, 52, 54, 56, 66).

148

#### 149 **Variation in Testing and Data Analysis Procedures**

150 Unfortunately, there is substantial variation across testing protocols reported within the  
151 scientific literature, including differences in knee and hip joint angles (120-150° and 124-175°,  
152 respectively), sampling frequency (500-2000 Hz), pull onset identification thresholds including  
153 absolute (20-75 N) and relative (2.5-10% body weight) threshold values, and smoothing and  
154 filtering approaches, with some authors not stating hip angles, thresholds or filtering  
155 procedures (Table 5). In addition, if practitioners or researchers are intending to use published  
156 values for comparison they should be mindful that some data is presented as net force (gross  
157 force – body weight) while others report gross measures, along with ratio and allometric  
158 scaling used in some studies. These two latter approaches may impact the results less, as  
159 allometric scaling uses an exponent related to body mass (13) although allometric scaling will  
160 reduce the resultant values compared to ratio scaling, with greater variation introduced  
161 depending on the exponent used (Table 5).

162

163

**Table 5: Reported Testing and Data Analysis Procedures**

Author(s)	Knee Angle	Hip Angle	Sampling Frequency	Onset Threshold	Scaling	Smoothing & Filtering	RFD Calculation
Haff et al. (33)	144 ± 5°	145 ± 3°	500 Hz	---	Net Force	---	PRFD (2 ms window)
Stone et al. (60)	140-145°	---	600 Hz	---	Net Absolute, Relative and AS	---	PRFD (1.7 ms window)
Haff et al. (30)	127-145° *	---	600 Hz	---	Net Force	---	PRFD (1.7 ms window)
McGuigan et al. (47)	130°	---	500 Hz	---	Absolute	---	PRFD (2 ms window)
Kawamori et al. (39)	141±10°	124±11°	500 Hz	---	---	---	PRFD (2 ms window)
Haff et al. (31)	127-145° *	---	600 Hz	---	Net Force	---	PRFD (1.7 ms window)
Nuzzo et al. (49)	140°	---	1000 Hz	---	Ratio	---	Mean RFD
Winchester et al. (74)	130°	---	---	---	Net	---	---
Winchester et al. (73) #	---	---	---	---	---	---	---
McGuigan & Winchester (45)	130°	---	960 Hz	---	---	---	--- Assumed peak due to the values
Kraska et al. (41)	120-135°	170-175° ¥ In line with Haff et al (1997)	1000 Hz	---	Absolute & AS	---	--- Assumed peak due to the values
Whittington et al. (72)	120-135° 'Self-selected'	170-175° 'Self-selected'	1000 Hz	---	---	---	PRFD (1 ms window)
McGuigan et al. (46)	130°	---	960 Hz	---	---	---	---

					Assumed Net due to the values		Assumed mean due to the values
West et al. (71)	120-130° ¥ In line with Haff et al (2005), Stone et al (2004)	---	1000 Hz	5SD of mean force after trigger	Net	Dual pass Butterworth filter (low pass, 20 Hz cut-off)	PRFD (1 ms window)
Crewther et al. (16)	120-130° ¥ In line with Haff et al (2005), Stone et al (2004)	---	1000 Hz	---	Net	Dual pass Butterworth filter (low pass, 20 Hz cut-off)	PRFD (1 ms window)
Beckham et al. (6)	¥ In line with Haff et al. (1997) and Kraska et al. (2009)	¥ In line with Haff et al. (1997) and Kraska et al. (2009)	1000 Hz	---	Absolute & AS	4 <sup>th</sup> Order Butterworth low pass filter 100 Hz	Not included
Beckham et al. (7)	120-135°	175°	1000 Hz	---	Absolute, Ratio & AS	4 <sup>th</sup> Order Butterworth low pass filter 100 Hz	Mean & PRFD (1 ms window)
Sheppard et al. (56)	130°	155-165°	600 Hz	---	Net	---	Not included
Comfort et al. (11)	120°, 130°, 140°, 150° & Self-selected (133 ± 3°)	125°, 145° & Self-selected (138 ± 4°)	600 Hz	40 N	Absolute	---	PRFD (1.7 ms window)
Thomas et al. (64)	Self-selected	Self-selected	600 Hz	---	Absolute	4 <sup>th</sup> Order Butterworth low pass filter 16 Hz	PRFD (1.7 ms window)
Thomas et al. (67)	Self-selected	Self-selected	600 Hz	---	Absolute & Relative	4 <sup>th</sup> Order Butterworth	PRFD (1.7 ms window)



						low pass filter 16 Hz	
Thomas et al. (66)	Self-selected	Self-selected	600 Hz	---	Absolute	4 <sup>th</sup> Order Butterworth low pass filter 16 Hz	Not included
Haff et al. (32)	140.0 ± 6.6°	137.6 ± 12.9°	1000 Hz	---	Net	Rectangular smoothing with a moving half-width of 12	PRFD (20 ms window) RFD <sub>30, 50, 90, 100,</sub> 150, 200, 250
Secomb, et al. (52)	125-140°	---	600 Hz	---	Absolute and Relative	---	Not included
Secomb et al. (53)	125-140°	---	600 Hz	---	Absolute and Relative	---	Not included
Secomb et al. (54)	--- Stated similar to Haff et al. (2005)	--- Stated similar to Haff et al. (2005)	600 Hz	---	Absolute and Relative	---	Not included
Tran et al. (68)	--- Stated similar to Haff et al. (1997)	--- Stated similar to Haff et al. (1997)	600 Hz	---	Absolute and Relative (Assumed Net due to the values)	4 <sup>th</sup> Order Butterworth low pass filter 10 Hz	
Spiteri et al. (58)	140°	140°	2000 Hz	---	Relative	---	RFD <sub>30, 50, 90, 100</sub>
Sjokvist et al. (57)	--- States in line with Stone et al. (2004)				Absolute and Relative	---	Not included
Welch et al. (70)	No specific detail provided other than bar positioned at mid-thigh				Relative	---	Not included
Wang et al. (69)	Self-selected	Self-selected	1000 Hz	---	Net	---	PRFD (20 ms window)

							RFD <sub>30, 50, 90, 100, 150, 200, 250</sub>
Mangine et al. (44)	Self-selected	Self-selected	1000 Hz	---	Net	---	PRFD (20 ms window) RFD <sub>30, 50, 90, 100, 150, 200, 250</sub>
Halperin et al. (34)	130-140°	Not stated	1000 Hz	---	---	---	Not included
Dos'Santos et al. (22)	Self-selected	Self-selected	2000 Hz (down-sampled to 1500, 1000 & 500 Hz)	75 N	Absolute	20 ms moving average	RFD <sub>100</sub> RFD <sub>150</sub> RFD <sub>200</sub>
Bartolomei et al. (4)	140°	125°	1000 Hz	---	Absolute	---	PRFD (20 ms window)
James et al. (38)	141.9 ± 4.3°	139.2 ± 4.1°	1000 Hz down sampled to 100 Hz to compare to strain gauge	20 N	Net	4 <sup>th</sup> Order Butterworth low pass filter 10 Hz	PRFD (20 ms window) RFD <sub>30, 50, 90, 100, 150, 200, 250</sub>
De Witt et al. (18)	144 ± 3°	137 ± 3°	1000 Hz	---	--- Assumed Net due to the values	---	PRFD (20 ms window) RFD <sub>30, 50, 90, 100, 150, 200, 250</sub>
Dos'Santos, Thomas et al. (24)	137-146° †	140-149° †	1000 Hz	40 N	Absolute	---	Not included
Dos'Santos, et al. (21)	Self-selected	Self-selected	1000 Hz	2.5% BW, 5% BW, 10% BW, >75 N, <b>5 SD BW</b>	Absolute	---	RFD <sub>100</sub> RFD <sub>150</sub> RFD <sub>200</sub>
Beckham et al. (8)	125°	125° & 145°	1000 Hz	---	Absolute & AS	2 <sup>nd</sup> Order Butterworth low pass filter 10 Hz	Not included

Oranchuk et al. (50)	135-145°	---	1000 Hz	2.5% of mean body mass, based on force-time data	Relative	4 <sup>th</sup> Order Butterworth filter, with 20 Hz cut-off	PRFD (20 ms window)
Dobbin et al. (20)	140°	Self-selected, shoulder above the bar (as described by Thomas et al., 2015)	1200 Hz	---	Net relative and AS	---	Not included
Beattie et al. (5)	131 ± 9°	---	1000 Hz	---	Relative	---	Not included
Dos'Santos et al. (26)	145°	145° & 175°	1000 Hz	5 SD BW	Net	Unfiltered	PRFD RFD <sub>100</sub> RFD <sub>150</sub> RFD <sub>200</sub>
Leary et al. (42)	142 ± 7°	146 ± 11°	1000 Hz	---		Rectangular smoothing with a moving half-width of 12	PRFD RFD <sub>30, 50, 90, 100, 150, 200, 250</sub>

--- = not stated

¥ = Incorrectly cites joint angles 'in line with previous research' when the referenced studies used different joint angles

Net Force = Gross Force – Body Weight

PRFD = Peak Instantaneous RFD (the greatest rate of change in force between two tangential points; the window differs based on sampling frequency)

Mean force (Change in force / change in time from onset of force production to time to peak force)

RFD<sub>100</sub> = subscript numbers refer to the epoch for mean RFD

\*Based on knee angle achieved during the 2<sup>nd</sup> pull phase of the clean for each individual

#Published abstract

¥ Self-selected to replicate the start of the second pull

BW = Body weight (during the initial period of quiet standing), SD = standard deviation

167 Numerous authors have suggested that the posture adopted during the IMTP should replicate  
168 the start of the second pull phase of the clean, (30, 31, 33, 60); however, only two studies  
169 have actually assessed the participants knee joint angles during the clean and then adopted  
170 these angles during the IMTP (30, 31). This is most likely due to time and practicality of  
171 assessing specific joint angles during the clean prior to performing the IMTP, especially when  
172 assessing large squads of athletes. Interestingly, hip joint angles were not reported within  
173 these two studies (30, 31).

174 Due to the variety of knee and hip joint angles reported within the literature, Comfort et al. (11)  
175 investigated a range of knee ( $120^\circ$ ,  $130^\circ$ ,  $140^\circ$ ,  $150^\circ$ ) and hip ( $125^\circ$ ,  $145^\circ$ ) joint angles, along  
176 with self-selected posture (knee  $133\pm 3^\circ$ , hip  $138\pm 4^\circ$ ) based on the athletes preferred position  
177 to start the second pull of a clean, which is what the posture adopted during the IMTP was  
178 originally based on (33). The results of the study indicated that there were no significant or  
179 meaningful differences in PF, PRFD or impulse between postures, although the preferred  
180 (self-selected) posture demonstrated the highest reliability and the lowest measurement error.  
181 In contrast, Beckham et al. (6) found that powerlifters produced greater PF during an isometric  
182 testing with a vertical torso compared to a deadlift-specific body position at the same bar  
183 height, described as being a “relatively straight legged position and somewhat bent over the  
184 bar”. The authors suggested that the upright position may have provided a mechanical  
185 advantage and a posture more optimal for force production against the bar. In another study,  
186 Beckham et al. (8) compared the effects of different hip joint angles ( $125^\circ$  vs.  $145^\circ$ ), while  
187 standardizing the knee joint angle ( $125^\circ$ ) reporting meaningful and significantly higher PF and  
188 force at different epochs (50, 90, 200, 250 ms) in the more upright ( $145^\circ$ ) position, especially  
189 in subjects with greater experience in performing weightlifting exercises and their derivatives,  
190 in contrast to Comfort et al. (11). Interestingly, Beckham et al. (8) reported small changes in  
191 joint angles throughout the execution of the test and based on these observations recommend  
192 that in the future researchers and practitioners should adopt standardized knee and hip angles  
193 of  $120\text{-}135^\circ$  and  $140\text{-}150^\circ$ , respectively.

194 More recently, Dos'Santos et al. (26) compared hip joint angles of 145° and 175° with a  
195 standardized knee joint angle of 145°, finding greater time specific force values and RFD at  
196 predetermined epochs, with a 145° hip angle (Table 5). The hip angle of 175° previously  
197 reported by Kraska et al. (41) and replicated by Beckham et al. (6) actually refer to trunk angle  
198 relative to vertical, to ensure an upright trunk (forward lean of 5° from vertical), exhibiting an  
199 upright trunk as previously described (30, 31, 33, 60) rather than a 175° hip angle as used by  
200 Dos'Santos et al. (26). The authors of a recent meta-analysis also highlight the fact the  
201 practitioners should carefully consider the specific protocol, including joint angles, to ensure  
202 repeatability of the measures (27).

203 While adopting standardized knee and hip angles during the IMTP may seem logical, this  
204 practice may place athletes in a sub-optimal pulling position, due to the range of angles  
205 reported across individuals for the second pull phase of the clean (30, 31). Therefore, it is best  
206 to consider the individual athletes' appropriate second pull position and then quantify the knee  
207 and hip angles. This practice allows for the individual athlete's anthropometrics to be  
208 considered and allows them to assume an optimal pulling position, in line with the range of  
209 joint angles recommended by Beckham et al. (8). Once the pulling position is established then  
210 it is recommended that practitioners and researchers ensure that the individual starting  
211 postures are replicated between trials and testing sessions. Joint angles should be assessed  
212 prior to the commencement of the pull due to slight changes in joint angles during the pull (8).

213 Haff et al. (32) suggest using minimal pre-tension prior to initiation of the pull, as this is likely  
214 to impact both time specified force and RFD, with Dos'Santos et al. (26) recently reporting that  
215 the 175° hip angle results in significantly higher 'body weight' due to increased pre-tension,  
216 compared to a 145° hip angle, which may have contributed to in the differences in time specific  
217 force values and RFD that were reported. Similarly, Maffioletti et al. (43) suggested that pre-  
218 tension is undesirable when assessing isometric RFD, albeit with a focus on single joint  
219 assessment; it would, therefore, be advantageous to visually inspect the force-time data pre

220 and post isometric pull, to ensure that there are no differences in force, which should represent  
221 body weight.

222 Interestingly, numerous authors state that they have adopted the postures previously reported  
223 by other researchers, but in fact report different angles to those stated in the studies that they  
224 cite, or cite multiple researchers who reported different postures (Table 5). These differing  
225 postures are most likely related to individual athlete anthropometric profiles. It is therefore  
226 important that researchers carefully report and justify their choice of joint angles, but more  
227 importantly, standardize these between trials and testing sessions.

228 Other researchers have used strain gauge based equipment, with the handle attached via a  
229 chain (16, 17, 37, 38, 48) with a range of sampling frequencies (100-133 Hz (17, 37, 38)) and  
230 joint angles (knee 120-130° (17), 142±4°(38), 143±7° (37), 160° (48); hip 139±4° (38), 144±5°  
231 (37)). However, findings of two research groups that compared strain gauge systems to a  
232 force platform demonstrated that the strain gauge significantly underestimated PF, by ~8%  
233 (38) to ~10% (20). Additionally, James et al. (38) found that measures of RFD did not meet  
234 acceptable standards of reliability. While such systems can measure PF, which can be ratio  
235 or allometrically scaled, there does not seem to be an effective way to accurately measure or  
236 calculate RFD, and are therefore not recommended if practitioners have access to a force  
237 platform.

238

### 239 **Recommendations for Correct IMTP Assessment**

240 Due to the noticeable variations in assessment procedures, including posture, sampling  
241 frequency, and methods of calculating specific variables (namely use of different sampling  
242 frequencies, onset thresholds, and the method for the calculation of RFD), we suggest  
243 appropriate standardization of all testing procedures for the IMTP. Such standardization  
244 should permit more meaningful comparisons of individual performances between testing  
245 sessions, comparisons between athletes and more effective comparisons between published

246 studies. Standardization should also include the verbal cues as attentional focus has been  
247 shown to affect force production, with an external focus of 'push as hard and fast as possible'  
248 resulting in greater PF compared to an internal focus (34).

249

## 250 **Recommended Testing Procedures**

251 Prior to initiation of IMTP testing, the bar height necessary to obtain the correct body position  
252 should be determined. This should be an iterative process in which the athlete starts with a  
253 bar height that allows the athlete to assume a body position that replicates the start of the  
254 second pull position during the clean. The bar height should then be adjusted up or down to  
255 allow the athlete to obtain the optimal knee (125-145°) and hip (140-150°) angles (6, 8, 26).  
256 The body position should be very similar to the second pull of the clean and the clean grip  
257 mid-thigh pull exercise (19): upright torso, slight flexion in the knee resulting in some  
258 dorsiflexion, shoulder girdle retracted and depressed, shoulders above or slightly behind the  
259 vertical plane of the bar, feet roughly centred under the bar approximately hip width apart,  
260 knees underneath and in front of the bar, and thighs in contact with the bar (close to the  
261 inguinal crease dependent on limb lengths) (Figure 2). When making joint measurements, the  
262 athlete should ensure that no tension is applied to the bar but that all "slack" (e.g. elbow flexion,  
263 shoulder girdle elevation/protraction) is removed from the body, as this would result in a  
264 change in joint angles during the maximal effort which is undesirable (8).

265

266

267

268

269



270

271 Figure 2: Correct posture for the isometric mid-thigh pull, illustrating an upright trunk,  
272 replicating the start position of the second pull of the clean


273

274 While the use of a “self-selected” body position is likely beneficial to efficiency of testing, it is  
275 not recommended without ensuring that the hip and knee joint angles fall within the ranges  
276 recommended above, due to the influence of body positioning on force generation (6, 8, 26).  
277 The bar height used and joint angles obtained should be recorded so that repeated  
278 measurements can be standardized and therefore replicate the individuals’ body position  
279 between session, ensuring that differing results in subsequent testing are not the result of  
280 changed body position (8, 26). It is also considered best practice to measure the individuals  
281 grip width and foot position and standardize these for individuals across sessions (unless  
282 working with youth athletes where changes in stature as a result of maturation may require  
283 increased stance and grip width) as each can affect body positioning relative to the bar (19).  
284 After the bar height and posture have been established, a short familiarization session of  
285 submaximal trials is recommended approximately 48 hours prior to testing (e.g. 3 x 3 second  
286 trials, each of 50-, 75- and 90% of perceived maximum effort). While a consensus on the



287 optimal amount of familiarization has not yet been reached, nearly all IMTP studies use some  
288 familiarization.


289 Athletes should complete some manner of standard generalized warm-up (62). While there is  
290 variability in the generalized warm-up chosen among studies, most studies use a warm-up  
291 that incorporates clean derivatives, such as the dynamic mid-thigh pull, and should thus be a  
292 component of the standard warm-up (7, 21, 24, 32, 33). Submaximal trials of the IMTP are  
293 also recommended prior to maximal effort trials (e.g. 3 seconds each of: 50% maximal effort,  
294 75% maximal effort, 90% maximal effort, separated by 60 seconds rest). During this time, the  
295 athlete should be secured to the bar using lifting straps and athletic tape to ensure that grip  
296 strength is not a limiting factor (Figure 3) (30, 33).



297 Figure 3: Standardized Warm Up Procedure

298 For each of the maximal effort trials, standardized instructions should be given to the athlete  
299 of some iteration of “push your feet into the ground as fast and as hard as possible” to ensure  
300 that both maximal RFD and PF are obtained (10, 34). It is essential that athletes understand  
301 that the focus is to drive the feet directly into the force platform and not attempt to pull the bar  
302 with the arms, or rise up on to their toes. The athlete should get into the correct body position  
303 for the IMTP, using just enough pre-tension to achieve the correct body position and remove  
304 “slack” from the body, but without any more pre-tension than is necessary to get the “quiet  
305 standing” necessary for a stable force baseline (43). This can be verified by monitoring the


306 athlete's body positioning and ensuring the force trace created by the athlete is both similar to  
 307 body mass and steady, with trials where a change in force >50 N occurs during this period  
 308 rejected (21). This should be explained to the athletes and they should be encouraged to stay  
 309 as still as possible during this period to accurately determine body weight and onset threshold.  
 310 A countdown of "3, 2, 1, *PULL!*" gives the athlete sufficient warning to be ready to give a  
 311 maximum effort and provides at least one second of quiet standing to enable the identification  
 312 of the onset of the pull (Figure 5a). Strong verbal encouragement from researchers and  
 313 teammates ensures that the athlete gives a maximum effort (9). A minimum of two trials should  
 314 be collected, provided that each of those trials have no errors by the athlete (e.g.  
 315 countermovement, excessive pre-tension, leaning on the bar prior to the pull (Figure 4). With  
 316 increasing PF, additional trials should be performed, until the PF values of the trials are  
 317 separated by <250 N (30, 33). It is noted, however, that a percentage of peak force may be  
 318 advantageous as an absolute value will affect stronger and weaker athletes differently,  
 319 although the exact effect of this has not been investigated.



**Acceptable trials <250 N difference in peak force, minimal pre-tension (<50 N) or countermovement at the start**

Figure 4: Standardized isometric mid-thigh pull testing procedure

321 Visual inspection of the force-time curves during testing can easily be used to determine if the  
322 trials are acceptable, or if additional trials should be performed. In addition to the trials being  
323 within 250 N between attempts, trials should be repeated if there is not a stable weighing  
324 period (clear fluctuation in the force-time data) or a clear countermovement prior to the  
325 initiation of the pull (Figure 5c), as this will interfere with accurate identification of the initiation  
326 of the pull (19), or if the PF occurs at the end of the trial (Figure 5b). It is also important to  
327 check that the force during the initial period of quiet standing (in the ready position, strapped  
328 to the bar, immediately prior to commencing the pull) represents body weight, and therefore  
329 no prior tension has been applied (Figure 5a) as this will interfere with pull onset identification  
330 (19).



331 Figure 5: Examples of acceptable and unacceptable isometric mid-thigh pull force-time traces

332 **Recommended Data Analysis and Reporting**

333 Collection of IMTP force-time data can be compiled accurately with a sampling frequency as  
 334 low as 500 Hz , but if higher sampling frequencies can be used then they are preferred as they  
 335 may increase the accuracy of time dependent measures (21). Specifically, the utilization of  
 336 frequencies  $\geq 1000$  Hz are recommended especially if early force-time variables are of interest  
 337 (e.g. force at 50 or 100 ms) (21). There are not enough data for a consensus regarding optimal  
 338 filtering and/or smoothing methods for the IMTP (23); although unfiltered data has been

339 suggested as optimal for analysis of countermovement jump performance (61) and where  
340 possible, unfiltered data for isometric testing (23, 43). It is therefore suggested that unfiltered  
341 and non-smoothed data is used for subsequent analysis (23), as most of the RFD and impulse  
342 characteristics are dependent upon an accurate determination of the start of the pull (21),  
343 although data from portable force platforms may exhibit greater 'noise' and warrant smoothing.  
344 Accurate identification of the start of the inflection point is often achieved using automated  
345 methods - we recommend using 5 standard deviations of body weight during an initial one  
346 second weighing period prior to the (usually one second) of quiet standing (in the ready  
347 position, strapped to the bar, immediately prior to commencing the pull) as the threshold for  
348 determining the onset of the pull (21), although this may vary with technical idiosyncrasies of  
349 different force platforms (e.g. noise magnitude). Trials that do not have a stable baseline force  
350 trace during the weighing period (change in force >50 N) should be rejected and subsequently  
351 another trial should be performed (21, 43) (Figure 5). To facilitate this stable period, it is  
352 essential to enforce and practice this during the warm-up / familiarization trials.

353 It is recommended that time-specific RFD epochs (50-, 100-, 150-, 200- and 250 ms commonly  
354 reported) should be used when using the IMTP as a sport performance diagnostic tool as  
355 these are not only reliable (32), but can be selected specific to the durations relevant to the  
356 specific sporting tasks, such as ground contact time during acceleration or peak running  
357 speeds. In contrast, maximal strength capabilities can be inferred from PF (Table 1).

358 When reporting results from IMTP testing, it is important that the hip and knee angles used by  
359 each athlete, to establish the bar height, be reported (8, 26). Such standardization of posture  
360 between trials and testing sessions ensures that data is comparable between sessions, groups  
361 of athletes and studies (8, 26). While there is no consensus as to the superiority of either net  
362 or gross force values for the IMTP, it is important that researchers report whether body weight  
363 was or was not included in the force and impulse values reported (7). Other methodological  
364 considerations, such as the method for identifying the onset of the pull (and threshold) (21),  
365 methods used for smoothing/filtering force platform data (23), sampling frequency and other

366 aspects of analysis (22), such as the exponent used for allometric scaling, should be reported,  
367 as each are important for accurately interpreting results from the study.

368

369

370

## 371 **References**

- 372 1. Bailey CA, Sato K, Alexander R, Chiang CY, and Stone M. Isometric force production  
373 symmetry and jumping performance in collegiate athletes. *Journal of Trainology* 2: 1-  
374 5, 2013.
- 375 2. Bailey CA, Sato K, Burnett A, and Stone MH. Carry-over of force production  
376 symmetry in athletes of differing strength levels. *J Strength Cond Res.* 29: 3188-  
377 3196, 2015.
- 378 3. Bailey CA, Sato K, Burnett A, and Stone MH. Force-production asymmetry in male  
379 and female athletes of differing strength levels. *Int J Sports Physiol Perform.* 10: 504-  
380 508, 2015.
- 381 4. Bartolomei S, Sadres E, Church DD, Arroyo E, Iii JAG, Varanoske AN, Wang R,  
382 Beyer KS, Oliveira LP, Stout JR, and Hoffman JR. Comparison of the recovery  
383 response from high-intensity and high-volume resistance exercise in trained men.  
384 *Eur J Appl Physiol* 117: 1287-1298, 2017.
- 385 5. Beattie K, Carson BP, Lyons M, and Kenny IC. The Effect of maximal- and explosive-  
386 strength training on performance indicators in cyclists. *Int J Sports Physiol Perform*  
387 12: 470-480, 2017.
- 388 6. Beckham G, Lamont H, Sato K, Ramsey M, Haff GG, and Stone M. Isometric  
389 strength of powerlifters in key positions of the conventional deadlift. *Journal of*  
390 *Trainology* 1, 2012.
- 391 7. Beckham G, Mizuguchi S, Carter C, Sato K, Ramsey M, Lamont H, Hornsby G, Haff  
392 G, and Stone M. Relationships of isometric mid-thigh pull variables to weightlifting  
393 performance. *J Sports Med Phys Fitness* 53: 573-581, 2013.
- 394 8. Beckham GK, Sato K, Mizuguchi S, Haff GG, and Stone MH. Effect of body position  
395 on force production during the isometric mid-thigh pull. *J Strength Cond Res.* 32(1):  
396 48-56: 18.
- 397 9. Belkhiria C, De Marco G, and Driss T. Effects of verbal encouragement on force and  
398 electromyographic activations during exercise. *J Sports Med Phys Fitness* 58: 750-  
399 757, 2018.
- 400 10. Bembem MG, Clasey JL, and Massey BH. The effect of the rate of muscle contraction  
401 on the force-time curve parameters of male and female subjects. *Res Q Exerc Sport*  
402 61: 96-99, 1990.
- 403 11. Comfort P, Jones PA, McMahon JJ, and Newton R. Effect of knee and trunk angle on  
404 kinetic variables during the isometric mid-thigh pull: test-retest reliability. *Int J Sports*  
405 *Physiol Perform* 10: 58-63, 2015.
- 406 12. Comfort P and McMahon JJ. Reliability of maximal back squat and power clean  
407 performances in inexperienced athletes. *J Strength Cond Res.* 29: 3089-3096, 2015.
- 408 13. Comfort P and Pearson SJ. Scaling--which methods best predict performance? *J*  
409 *Strength Cond Res* 28: 1565-1572, 2014.
- 410 14. Comfort P, Thomas C, Dos'Santos T, Jones PA, Suchomel TJ, and McMahon JJ.  
411 Comparison of methods of calculating dynamic strength index. *Int J Sports Physiol*  
412 *Perform:* 1-20, 2017.

- 413 15. Cormie P, McGuigan MR, and Newton RU. Developing maximal neuromuscular  
414 power: Part 1--biological basis of maximal power production. *Sports Med* 41: 17-38,  
415 2011.
- 416 16. Crewther BT, Carruthers J, Kilduff LP, Sanctuary CE, and Cook CJ. Temporal  
417 associations between individual changes in hormones, training motivation and  
418 physical performance in elite and non-elite trained men. *Biol Sport* 33: 215-221,  
419 2016.
- 420 17. Davis GR, Gallien GJ, Moody KM, LeBlanc NR, Smoak PR, and Bellar D. Cognitive  
421 function and salivary DHEA levels in physically active elderly african american  
422 women. *Int J Endocrinol* 2015: 6, 2015.
- 423 18. De Witt JK, English KL, Crowell JB, Kalogera KL, Guilliams ME, Nieschwitz BE,  
424 Hanson AM, and Ploutz-Snyder LL. Isometric mid-thigh pull reliability and relationship  
425 to deadlift 1RM. *J Strength Cond Res*. 32 (2): 528-533. 2018.
- 426 19. DeWeese BH, Serrano AJ, Scruggs SK, and Burton JD. The midhigh pull: proper  
427 application and progressions of a Weightlifting movement derivative. *Strength Cond J*  
428 35: 54-58, 2013.
- 429 20. Dobbin N, Hunwicks R, Jones B, Till K, Highton J, and Twist C. Criterion and  
430 construct validity of an isometric mid-thigh pull dynamometer for assessing whole  
431 body strength in professional rugby league players. *Int J Sports Physiol Perform*.  
432 13(2): 235-239. 2018.
- 433 21. Dos'Santos T, Jones PA, Comfort P, and Thomas C. Effect of different onset  
434 thresholds on isometric mid-thigh pull force-time variables. *J Strength Cond Res* 31:  
435 3467-3473, 2017.
- 436 22. Dos'Santos T, Jones PA, Kelly J, McMahon JJ, Comfort P, and Thomas C. Effect of  
437 sampling frequency on isometric midhigh-pull kinetics. *Int J Sports Physiol Perform*  
438 11: 255-260, 2016.
- 439 23. Dos'Santos T, Lake JP, Jones PA, and Comfort P. Effect of low pass filtering on  
440 isometric mid-thigh pull kinetics. *J Strength Cond Res*. 32: 983-989, 2018.
- 441 24. Dos'Santos T, Thomas C, Comfort P, McMahon JJ, Jones PA, Oakley NP, and  
442 Young AL. Between-session reliability of isometric mid-thigh pull kinetics and  
443 maximal power clean performance in male youth soccer players. *J Strength Cond*  
444 *Res* Published ahead of print, 2017.
- 445 25. Dos'Santos T, Thomas C, Jones PA, and Comfort P. Assessing muscle strength  
446 asymmetry via a unilateral stance isometric mid-thigh pull. *Int J Sports Physiol*  
447 *Perform*. 12(4): 505-511. 2017.
- 448 26. Dos'Santos T, Thomas C, Jones PA, McMahon JJ, and Comfort P. The effect of hip  
449 joint angle on isometric mid-thigh pull kinetics. *J Strength Cond Res*. 31(10):2748-  
450 2757. 2017 .
- 451 27. Drake D, Kennedy R, and Wallace E. The validity and responsiveness of isometric  
452 lower body multi-joint tests of muscular strength: a systematic review. *Sports Med*  
453 *Open* 3: 23, 2017.
- 454 28. Faigenbaum AD, McFarland JE, Herman RE, Naclerio F, Ratamess NA, Kang J, and  
455 Myer GD. Reliability of the one-repetition-maximum power clean test in adolescent  
456 athletes. *J Strength Cond Res* 26: 432-437, 2012.
- 457 29. Gescheit DT, Cormack SJ, Reid M, and Duffield R. Consecutive days of prolonged  
458 tennis match play: performance, physical, and perceptual responses in trained  
459 players. *Int J Sports Physiol Perform*. 10: 913-920, 2015.
- 460 30. Haff GG, Carlock JM, Hartman MJ, Kilgore JL, Kawamori N, Jackson JR, Morris RT,  
461 Sands WA, and Stone MH. Force-time curve characteristics of dynamic and isometric  
462 muscle actions of elite women olympic weightlifters. *J Strength Cond Res* 19: 741-  
463 748, 2005.
- 464 31. Haff GG, Jackson JR, Kawamori N, Carlock JM, Hartman MJ, Kilgore JL, Morris RT,  
465 Ramsey MW, Sands WA, and Stone MH. Force-time curve characteristics and  
466 hormonal alterations during an eleven-week training period in elite women  
467 weightlifters. *J Strength Cond Res* 22: 433-446, 2008.

- 468 32. Haff GG, Ruben RP, Lider J, Twine C, and Cormie P. A comparison of methods for  
469 determining the rate of force development during isometric mid-thigh clean pulls. *J*  
470 *Strength Cond Res* 29: 386-395, 2015.
- 471 33. Haff GG, Stone M, O'Bryant HS, Harman E, Dinan C, Johnson R, and Han K-H.  
472 Force-time dependent characteristics of dynamic and isometric muscle actions. *J*  
473 *Strength Cond Res*. 11: 269-272, 1997.
- 474 34. Halperin I, Williams KJ, Martin DT, and Chapman DW. The effects of attentional  
475 focusing instructions on force production during the isometric midhigh pull. *J*  
476 *Strength Cond Res*. 30: 919-923, 2016.
- 477 35. Helms ER, Zinn C, Rowlands DS, Naidoo R, and Cronin J. High-protein, low-fat,  
478 short-term diet results in less stress and fatigue than moderate-protein, moderate-fat  
479 diet during weight loss in male Weightlifters: A pilot study. *Int J Sport Nutr Exerc*  
480 *Metab*. 25: 163-170, 2015.
- 481 36. Hornsby W, Gentles J, MacDonald C, Mizuguchi S, Ramsey M, and Stone M.  
482 Maximum strength, rate of force development, jump height, and peak power  
483 alterations in Weightlifters across five months of training. *Sports* 5: 78, 2017.
- 484 37. James LP, Beckman EM, Kelly VG, and Haff GG. The neuromuscular qualities of  
485 higher and lower-level mixed martial arts competitors. *Int J Sports Physiol Perform*.  
486 12(5): 612-620. 2017.
- 487 38. James LP, Roberts LA, Haff GG, Kelly VG, and Beckman EM. Validity and reliability  
488 of a portable isometric mid-thigh clean pull. *J Strength Cond Res* 31: 1378-1386,  
489 2017.
- 490 39. Kawamori N, Rossi SJ, Justice BD, Haff EE, Pistilli EE, O'Bryant HS, Stone MH, and  
491 Haff GG. Peak force and rate of force development during isometric and dynamic  
492 mid-thigh clean pulls performed at various intensities. *J Strength Cond Res*. 20: 483-  
493 491, 2006.
- 494 40. Khamoui AV, Brown LE, Nguyen D, Uribe BP, Coburn JW, Noffal GJ, and Tran T.  
495 Relationship between force-time and velocity-time characteristics of dynamic and  
496 isometric muscle actions. *J Strength Cond Res* 25: 198-204, 2011.
- 497 41. Kraska JM, Ramsey MW, Haff GG, Fethke N, Sands WA, Stone ME, and Stone MH.  
498 Relationship between strength characteristics and unweighted and weighted vertical  
499 jump height. *Int J Sports Physiol Perform*. 4: 461-473, 2009.
- 500 42. Leary BK, Statler J, Hopkins B, Fitzwater R, Kesling T, Lyon J, Phillips B, Bryner RW,  
501 Cormie P, and Haff GG. The relationship between isometric force-time curve  
502 characteristics and club head speed in recreational golfers. *J Strength Cond Res* 26:  
503 2685-2697, 2012.
- 504 43. Maffiuletti NA, Aagaard P, Blazevich AJ, Folland J, Tillin N, and Duchateau J. Rate of  
505 force development: physiological and methodological considerations. *Eur J Appl*  
506 *Physiol* 116: 1091-1116, 2016.
- 507 44. Maigne GT, Hoffman JR, Wang R, Gonzalez AM, Townsend JR, Wells AJ, Jajtner  
508 AR, Beyer KS, Boone CH, Miramonti AA, LaMonica MB, Fukuda DH, Ratamess NA,  
509 and Stout JR. Resistance training intensity and volume affect changes in rate of force  
510 development in resistance-trained men. *Eur J Applied Physiol*. 116: 2367-2374,  
511 2016.
- 512 45. McGuigan M and Winchester JB. The relationship between isometric and dynamic  
513 strength in collegiate football players. *J Sports Sci Med* 7: 101-105, 2008.
- 514 46. McGuigan MR, Newton MJ, Winchester JB, and Nelson AG. Relationship between  
515 isometric and dynamic strength in recreationally trained men. *J Strength Cond Res*  
516 24: 2570-2573, 2010.
- 517 47. McGuigan MR, Winchester JB, and Erickson T. The importance of isometric  
518 maximum strength In college wrestlers. *J Sports Sci Med*. 5: 108-113, 2006.
- 519 48. Moran J, Sandercock GRH, Ramírez-Campillo R, Wooller J-J, Logothetis S,  
520 Schoenmakers PPJM, and Parry DA. Maturation-related differences in adaptations to  
521 resistance training in young male swimmers. *J Strength Cond Res*. 32(1): 139-149.  
522 2018.



- 523 49. Nuzzo JL, McBride JM, Cormie P, and McCaulley GO. Relationship between  
524 countermovement jump performance and multijoint isometric and dynamic tests of  
525 strength. *J Strength Cond Res* 22: 699-707, 2008.
- 526 50. Oranchuk DJ, Robinson TL, Switaj ZJ, and Drinkwater EJ. Comparison of the hang  
527 high-pull and loaded jump squat for the development of vertical jump and isometric  
528 force-time characteristics. *J Strength Cond Res*. Publish Ahead of Print, 2017.
- 529 51. Painter KB, Haff GG, Ramsey MW, McBride J, Triplett T, Sands WA, Lamont HS,  
530 Stone ME, and Stone MH. Strength gains: block versus daily undulating periodization  
531 weight training among track and field athletes. *Int J Sports Physiol Perform* 7: 161-  
532 169, 2012.
- 533 52. Secomb JL, Farley OR, Lundgren L, Tran T, King A, Nimphius S, and Sheppard J.  
534 Associations between the performance of scoring manouvres and lower-body  
535 strength and power in elite surfers. *Int J Sports Sci Coach* 10: 911-918, 2015.
- 536 53. Secomb JL, Lundgren LE, Farley OR, Tran TT, Nimphius S, and Sheppard JM.  
537 Relationships between lower-body muscle structure and lower-body strength, power,  
538 and muscle-tendon complex stiffness. *J Strength Cond Res* 29: 2221-2228, 2015.
- 539 54. Secomb JL, Nimphius S, Farley OR, Lundgren L, Tran T, and Sheppard J.  
540 Relationships between lower-body muscle structure and, lower-body strength,  
541 explosiveness and eccentric leg stiffness in adolescent athletes. *J Sports Sci Med*  
542 14: 691-697, 2015.
- 543 55. Seitz LB, Reyes A, Tran TT, de Villarreal ES, and Haff GG. Increases in lower-body  
544 strength transfer positively to sprint performance: a systematic review with meta-  
545 analysis. *Sports Med* 44 1693-1702, 2014.
- 546 56. Sheppard J, Chapman D, and Taylor K. An evaluation of a strength qualities  
547 assessment method for the lower body. *JASC* 19: 4-10, 2011.
- 548 57. Sjokvist J, Sandbakk O, Willis SJ, Andersson E, and Holmberg HC. The effect of  
549 incline on sprint and bounding performance in cross-country skiers. *J Sports Med*  
550 *Phys Fitness* 55: 405-414, 2015.
- 551 58. Spiteri T, Newton RU, and Nimphius S. Neuromuscular strategies contributing to  
552 faster multidirectional agility performance. *J Electromyogr Kinesiol* 25: 629-636,  
553 2015.
- 554 59. Spiteri T, Nimphius S, Hart NH, Specos C, Sheppard JM, and Newton RU.  
555 Contribution of strength characteristics to change of direction and agility performance  
556 in female basketball athletes. *J Strength Cond Res* 28: 2415-2423, 2014.
- 557 60. Stone MH, Sands WA, Carlock J, Callan S, Dickie D, Daigle K, Cotton J, Smith SL,  
558 and Hartman M. The importance of isometric maximum strength and peak rate-of-  
559 force development in sprint cycling. *J Strength Cond Res* 18: 878-884, 2004.
- 560 61. Street G, McMillan S, Board W, Rasmussen M, and Heneghan JM. Sources of error  
561 in determining countermovement jump height with the impulse method. *J Appl*  
562 *Biomech* 17: 43-54, 2001.
- 563 62. Suchomel TJ, Lamont HS, and Moir GL. Understanding vertical jump potentiation: a  
564 deterministic model. *Sports Med* 46: 809-828, 2016.
- 565 63. Suchomel TJ, Nimphius S, and Stone MH. The importance of muscular strength in  
566 athletic performance. *Sports Med* 46: 1419-1449, 2016.
- 567 64. Thomas C, Comfort P, Chiang CY, and Jones PA. Relationship between isometric  
568 mid-thigh pull variables and sprint and change of direction performance in collegiate  
569 athletes. *Journal of Trainology* 4: 6-10, 2015.
- 570 65. Thomas C, Dos'Santos T, Comfort P, and Jones P. Between-session reliability of  
571 common strength- and power-related measures in adolescent athletes. *Sports* 5: 15,  
572 2017.
- 573 66. Thomas C, Jones PA, and Comfort P. Reliability of the dynamic strength index in  
574 collegiate athletes. *Int J Sports Physiol Perform* 10: 542-545, 2015.
- 575 67. Thomas C, Jones PA, Rothwell J, Chiang CY, and Comfort P. An Investigation into  
576 the relationship between maximum isometric strength and vertical jump performance.  
577 *J Strength Cond Res* 29: 2176-2185, 2015.

- 578 68. Tran TT, Lundgren L, Secomb J, Farley ORL, Haff GG, Seitz LB, Newton RU,  
579 Nimphius S, and Sheppard JM. Comparison of physical capacities between  
580 nonselected and selected elite male competitive surfers for the national junior team.  
581 *Int J Sports Physiol Perform* 10: 178-182, 2015.
- 582 69. Wang R, Hoffman JR, Tanigawa S, Miramonti AA, La Monica MB, Beyer KS, Church  
583 DD, Fukuda DH, and Stout JR. Isometric mid-thigh pull correlates with strength,  
584 sprint, and agility performance in collegiate rugby union players. *J Strength Cond  
585 Res* 30: 3051-3056, 2016.
- 586 70. Welch N, Moran K, Antony J, Richter C, Marshall B, Coyle J, Falvey E, and Franklyn-  
587 Miller A. The effects of a free-weight-based resistance training intervention on pain,  
588 squat biomechanics and MRI-defined lumbar fat infiltration and functional cross-  
589 sectional area in those with chronic low back. *BMJ Open Sport & Exercise Medicine*  
590 1, 2015.
- 591 71. West DJ, Owen NJ, Jones MR, Bracken RM, Cook CJ, Cunningham DJ, Shearer DA,  
592 Finn CV, Newton RU, Crewther BT, and Kilduff LP. Relationships between force-time  
593 characteristics of the isometric midhigh pull and dynamic performance in  
594 professional rugby league players. *J Strength Cond Res* 25: 3070-3075, 2011.
- 595 72. Whittington J, Schoen E, Labounty LL, Hamdy R, Ramsey MW, Stone ME, Sands  
596 WA, Haff GG, and Stone MH. Bone mineral density and content of collegiate  
597 throwers: influence of maximum strength. *J Sports Med Phys Fitness* 49: 464-473,  
598 2009.
- 599 73. Winchester J, McGuigan MR, Nelson AG, and Newton M. The relationship between  
600 isometric and dynamic strength in college aged males. *J Strength Cond Res* 24: 1,  
601 2010.
- 602 74. Winchester JB, McBride JM, Maher MA, Mikat RP, Allen BK, Kline DE, and  
603 McGuigan MR. Eight weeks of ballistic exercise improves power independently of  
604 changes in strength and muscle fiber type expression. *J Strength Cond Res* 22:  
605 1728-1734, 2008.

606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631

632 Author Biographies:

633

634 Paul Comfort is a Reader in Strength and Conditioning and the programme leader for the  
635 Masters in Strength and Conditioning at the University of Salford.

636 George Beckham is an assistant professor in the Kinesiology Department at California State  
637 University, Monterey Bay.

638 Tom Dos'Santos is a Doctoral student in Biomechanics and Strength and Conditioning at the  
639 University of Salford.

640 Stuart Guppy is a Master's candidate in the area of Sports Science at Edith Cowan  
641 University.

642 Michael H. Stone is Professor, Graduate Coordinator and Exercise and Sport Science  
643 Laboratory Director in the Department of Sport, Exercise, Recreation and Kinesiology at  
644 East Tennessee State University.

645 G. Gregory Haff is the Course Coordinator for the Masters of Exercise Science (Strength  
646 and Conditioning) at Edith Cowan University and served as the President of the National  
647 Strength and Conditioning Association from 2015-2018.