



University of
Salford
MANCHESTER

The association between *Toxoplasma gondii* infection and postpartum blues

Gao, J-M, He, Z-H, Xie, Y-T, Hide, G, Lai, D-H and Lun, Z-R

<http://dx.doi.org/10.1016/j.jad.2019.03.013>

Title	The association between <i>Toxoplasma gondii</i> infection and postpartum blues
Authors	Gao, J-M, He, Z-H, Xie, Y-T, Hide, G, Lai, D-H and Lun, Z-R
Type	Article
URL	This version is available at: http://usir.salford.ac.uk/id/eprint/50675/
Published Date	2019

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: usir@salford.ac.uk.




Figure S1A. Quantitative comparison between *Toxoplasma gondii* immunoglobulin (IgG) optical density (OD) and Hamilton Depression Scale (HAMD) score in all new mothers (n=475). There is no significant correlation between IgG optical density and HAMD score ($p = 0.872$).




Figure S1B. Quantitative analysis of all *Toxoplasma gondii* seropositive new mothers (n=27) by comparison of immunoglobulin (IgG) optical density (OD) and Hamilton Depression Scale (HAMD) score in the same mothers. There is no significant correlation between IgG optical density and HAMD score ($p= 0.214$).