



University of
Salford
MANCHESTER

Exact soliton solutions of the nonlinear Helmholtz equation: communication

Chamorro-Posada, P, McDonald, GS and New, GHC

<http://dx.doi.org/10.1364/JOSAB.19.001216>

Title	Exact soliton solutions of the nonlinear Helmholtz equation: communication
Authors	Chamorro-Posada, P, McDonald, GS and New, GHC
Type	Article
URL	This version is available at: http://usir.salford.ac.uk/18330/
Published Date	2002

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: usir@salford.ac.uk.

JOSA COMMUNICATIONS

The JOSA Communications section is for short papers reporting new results that are deemed to be especially significant or of wide interest. A Communication must be no longer than four printed pages, including title and abstract. Page proofs are provided to authors.

Exact soliton solutions of the nonlinear Helmholtz equation: communication

P. Chamorro-Posada

Área de Teoría de la Señal y Comunicaciones, Departamento de Ingeniería Eléctrica, Electrónica, Computadores y Sistemas, Universidad de Oviedo Edificio Departamental 4, Campus de Viesques, 33204-Gijón, Spain

G. S. McDonald

Photonics and Nonlinear Science Group, Joule Physics Laboratory, School of Sciences, University of Salford, Salford M5 4WT, UK

G. H. C. New

Quantum Optics and Laser Science Group, The Blackett Laboratory, Imperial College of Science, Technology, and Medicine, Prince Consort Road, London SW7 2BW, UK

Received March 9, 2001

Exact analytical soliton solutions of the nonlinear Helmholtz equation are reported. A lucid generalization of paraxial soliton theory that incorporates nonparaxial effects is found. © 2002 Optical Society of America
OCIS codes: 190.3270, 190.5530, 240.4350.

1. INTRODUCTION

Investigations of the propagation of optical beams in Kerr-type media are usually based on the nonlinear Schrödinger equation (NSE), which is particularly appealing because of its analytical properties. The main drawback of this approach comes from the limitations imposed by the paraxial approximation that is implicit in the NSE. To overcome these difficulties we developed techniques based on a nonparaxial NSE (NNSE) that is derived without the use of the paraxial approximation; exact analytical solutions were derived for bright two-dimensional solitons, and their properties were thoroughly analyzed.¹ Further results were presented in recent publications^{2,3} and have been widely disseminated at conferences.^{4–7}

In a recent paper,⁸ analytical solutions that correspond to soliton solutions of the nonlinear Helmholtz equation were presented. The bright and dark solutions for $N + 1$ dimensions were reported as new, and the authors then concentrated attention on the highly physical two-dimensional solutions. In this Communication we address key questions that were not fully answered in Ref. 8.

2. RESULTS

The two-dimensional nonlinear Helmholtz equation,

$$\frac{\partial^2 E}{\partial z^2} + \frac{\partial^2 E}{\partial x^2} + k^2 E + \gamma |E|^2 E = 0, \quad (1)$$

where $E(x, z)$ is the electric field, $k = n_0 \omega/c$, and $\gamma = 2n_0 n_2 \omega^2/c^2$, is transformed into the NNSE:

$$\kappa \frac{\partial^2 u}{\partial \zeta^2} + i \frac{\partial u}{\partial \zeta} + \frac{1}{2} \frac{\partial^2 u}{\partial \xi^2} + |u|^2 u = 0, \quad (2)$$

when it is written in terms of the field envelope $A(x, z)$, where $E(x, z) = A(x, z) \exp(jkz)$ and the normalizations $\zeta = z/L_D$, $\xi = \sqrt{2}x/w_0$, and $u(\xi, \zeta) = (kn_2 L_D / n_0)^{1/2} A(\xi, \zeta)$ have been employed. w_0 is a reference beam width, $L_D = k \omega_0^2/2$ is the corresponding diffraction length, and $\kappa = 1/(k \omega_0)^2$ is a measure of the degree to which an input beam that propagates along the ζ axis deviates from being paraxial. In the paraxial limit, when the first term in Eq. (2) is neglected the NSE is recovered. A detailed account of the physical properties of the NNSE and its solutions can be found in Refs. 1–7. In what follows, we focus only on those results that are relevant to the study presented in Ref. 8.

The fundamental bright-soliton solution of the nonlinear Helmholtz equation, in terms of the normalizations of the NNSE, can be written as a lucid generalization of a paraxial soliton:

$$u(\xi, \zeta) = \eta \operatorname{sech} \left[\frac{\eta(\xi + V\zeta)}{\sqrt{1 + 2\kappa V^2}} \right] \\ \times \exp \left[i \left(\frac{1 + 2\kappa\eta^2}{1 + 2\kappa V^2} \right)^{1/2} \left(-V\xi + \frac{\zeta}{2\kappa} \right) \right] \\ \times \exp \left(\frac{-i\zeta}{2\kappa} \right), \quad (3)$$

where η , V , and κ are the amplitude, transverse velocity, and nonparaxial parameters, respectively. When nonparaxial effects are negligible, i.e., when $\kappa \rightarrow 0$, $\kappa\eta^2 \rightarrow 0$, and $\kappa V^2 \rightarrow 0$, one recovers the well-known paraxial soliton solution. The width of the nonparaxial soliton is given by $\xi_0 = (1 + 2\kappa V^2)^{1/2}/\eta$. The soliton's area is thus proportional to $\xi_0 \eta = (1 + 2\kappa V^2)^{1/2}$ and depends on both the transverse velocity and the actual size of the beam (through κ). The latter property reflects the fact that the solutions of the NNSE are invariant under transformations

$$\xi = \frac{\xi' + V\zeta'}{(1 + 2\kappa V^2)^{1/2}}, \quad \zeta = \frac{-2\kappa V\xi' + \zeta'}{(1 + 2\kappa V^2)^{1/2}}, \quad (4)$$

$$u(\xi, \zeta) = \exp \left\{ i \left(\frac{V\xi'}{(1 + 2\kappa V^2)^{1/2}} \right. \right. \\ \left. \left. + \frac{1}{2\kappa} \left[1 - \frac{1}{(1 + 2\kappa V^2)^{1/2}} \right] \zeta' \right) \right\} u'(\xi', \zeta'), \quad (5)$$

from which the well-known Galilean transformation invariance of the NSE is recovered in the appropriate paraxial limit. The transformations of Eqs. (4) and (5) correspond to a rotation of the solution in the original (unscaled) coordinate system by an angle θ , given by $\sec \theta = (1 + 2\kappa V^2)^{1/2}$ (Refs. 1 and 2); this allows the analysis to be extended to common nonparaxial situations in which off-axis soliton beams are considered. It has further been shown² that these transformations can be used to predict the long-term evolution of nonparaxial solitons when they are used in conjunction with analytical paraxial techniques.⁹

With respect to the conservation properties of the NNSE, we have generalized the first conserved quantity of the NSE, $\int_{-\infty}^{+\infty} |u(\xi, \zeta)|^2 d\xi = C$, where C is a constant, to the framework of the NNSE. This quantity has a correspondence to the conservation of an energy flow that is given by

$$\int_{-\infty}^{+\infty} \left[\frac{1}{2\kappa} + \frac{\partial \phi(\xi, \zeta)}{\partial \zeta} \right] |u(\xi, \zeta)|^2 d\xi = C', \quad (6)$$

where C' is another constant. This exact result generalizes a previously published (approximate) expression for which only fast on-axis phase variations were taken into account.^{10,11} The NSE conservation law is recovered from Eq. (6) in the paraxial limit.

Considering the effect that exact solutions of the NSE have had on a wide range of physical systems in which solitonlike solutions exert a dominant influence, the results given above, along with those of Refs. 1–8, are likely to underpin a broad class of exciting new research topics in nonlinear optics in which higher-order effects and related geometries are considered.

P. Chamorro-Posada's e-mail address is pedro@tsc.uniovi.es.

REFERENCES

1. P. Chamorro-Posada, G. S. McDonald, and G. H. C. New, "Non-paraxial solitons," *J. Mod. Opt.* **45**, 1111–1121 (1998).
2. P. Chamorro-Posada, G. S. McDonald, and G. H. C. New, "Propagation properties of non-paraxial spatial solitons," *J. Mod. Opt.* **47**, 1877–1886 (2000).
3. P. Chamorro-Posada, G. S. McDonald, and G. H. C. New, "Non-paraxial beam propagation methods," *Opt. Commun.* **192**, 1–12 (2001).
4. P. Chamorro-Posada, G. S. McDonald, and G. H. C. New, "Nonparaxial solitons," presented at the UK National Conference of Quantum Electronics, Cardiff, UK, 1997.
5. P. Chamorro-Posada, G. S. McDonald, and G. H. C. New, "Exact analytical nonparaxial optical solitons," presented at the 1998 International Conference on Quantum Electronics, San Francisco, Calif.
6. P. Chamorro-Posada, G. S. McDonald, and G. H. C. New, "Nonparaxial solitons: accurate numerical methods and new results," presented at the 1998 European Conference on Quantum Electronics, Glasgow, UK.
7. P. Chamorro-Posada, G. S. McDonald, G. H. C. New, J. Noon, and S. Chávez-Cerda, "Fundamental properties of $(1+1)D$ and $(2+1)D$ nonparaxial optical solitons," presented at the UK National Conference of Quantum Electronics, Manchester, UK, 1999.
8. T. A. Laine and A. T. Friberg, "Self-guided waves and exact solutions of the nonlinear Helmholtz equation," *J. Opt. Soc. Am. B* **17**, 751–757 (2000).
9. J. Satsuma and N. Yajima, "Initial value problems of one-dimensional self-modulation of nonlinear waves in dispersive media," *Suppl. Prog. Theor. Phys.* **55**, 284–306 (1974).
10. N. Akhmediev, A. Ankiewicz, and J. M. Soto-Crespo, "Does the nonlinear Schrödinger equation correctly describe beam propagation?" *Opt. Lett.* **18**, 411–413 (1993).
11. J. M. Soto-Crespo and N. Akhmediev, "Description of the self-focusing and collapse effects by a modified nonlinear Schrödinger equation," *Opt. Commun.* **101**, 223–230 (1993).