



University of
Salford
MANCHESTER

Snell's law for nonlinear beams

Christian, JM, McDonald, GS, Sanchez-Curto, J and Chamorro-Posada, P

Title	Snell's law for nonlinear beams
Authors	Christian, JM, McDonald, GS, Sanchez-Curto, J and Chamorro-Posada, P
Publication title	
Publisher	
Type	Conference or Workshop Item
USIR URL	This version is available at: http://usir.salford.ac.uk/id/eprint/18447/
Published Date	2010

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: library-research@salford.ac.uk.

Snell's Law for Nonlinear Beams

J M Christian¹, G S McDonald¹, J Sánchez-Curto² and P Chamorro-Posada²

¹ Materials & Physics Research Centre, University of Salford, M5 4WT, UK

² ETSI Telecomunicación, Universidad de Valladolid, Spain

The refraction of plane waves at a uniform boundary between dissimilar linear dielectric materials is perhaps one of the oldest and best well-understood phenomenon in optics. In contrast, the behaviour of a beam with a finite transverse cross-section is much less well understood. The situation is even more complicated when the two materials in question are nonlinear in nature. Seminal analyses some two decades ago considered scalar spatial optical solitons incident on the boundary between two dissimilar Kerr-type media [1]. While these analyses were highly instructive, they were based upon a governing nonlinear Schrödinger equation whose central tenet was that angles of incidence, reflection and refraction must be negligibly small. In practice, this is not entirely satisfactory since one would like these angles to be unconstrained.

To this end, we have recently developed a theory describing the refraction of soliton beams incident *at any angle* on a Kerr-type interface [2]. Our latest efforts have been to extend our Kerr analyses to a much wider class of power-law materials, of which the Kerr nonlinearity is a particular case. By a curious twist of fate, it turns out that the full nonlinear-beams problem can be described by a simple generalization of the trivially-familiar Snell's law. Figure 1 shows excellent agreement between theoretical predictions and full simulations of the interface problem.




Figure 1. (Left) Typical interface geometry and (right) predictions of soliton refraction using the new Snell law, $\gamma n_{01} \cos \theta_i = n_{02} \cos \theta_r$, where γ is a universal function of system and beam parameters.

References

- [1] A B Aceves, J V Moloney, and A C Newell, Phys Rev A **39**, 1809 (1989).
- [2] J Sanchez-Curto, P Chamorro-Posada and G S McDonald, Opt Lett **32**, 1126 (2007).