



University of  
**Salford**  
MANCHESTER

# Nonparaxial refraction and giant Goos-Hänchen shifts at nonlinear optical interfaces

McCoy, EA, Christian, JM and McDonald, GS

<b>Title</b>	Nonparaxial refraction and giant Goos-Hänchen shifts at nonlinear optical interfaces
<b>Authors</b>	McCoy, EA, Christian, JM and McDonald, GS
<b>Publication title</b>	
<b>Publisher</b>	Bristol Centre for Complexity Sciences
<b>Type</b>	Conference or Workshop Item
<b>USIR URL</b>	This version is available at: <a href="http://usir.salford.ac.uk/id/eprint/23000/">http://usir.salford.ac.uk/id/eprint/23000/</a>
<b>Published Date</b>	2012

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: [library-research@salford.ac.uk](mailto:library-research@salford.ac.uk).

# **Nonparaxial refraction and giant Goos-Hänchen shifts at nonlinear optical interfaces**

E. A. McCoy, J. M. Christian, and G. S. McDonald  
*Materials & Physics Research Centre,  
University of Salford, Greater Manchester M5 4WT, UK*

## **KEYWORDS:**

spatial solitons, self-focusing, optical refraction, nonlinear materials, Goos-Hänchen shift

## **THEME 1: NONLINEAR DYNAMICS**

### **APPLICATION THEME 1: PHYSICAL SCIENCES**

The scattering of spatial optical solitons (self-localizing beams of laser light) at the interface between two dissimilar materials is a problem of fundamental importance in nonlinear photonics. Theoretical analyses must take into account a highly complex interplay between diffraction, self-lensing, finite beam waists, and discontinuities in both the linear and nonlinear properties of the host medium at the boundary. Over the past three decades, various research groups worldwide have resorted to simplified mathematical descriptions based on the universal nonlinear Schrödinger equation [A.B. Aceves *et al.*, Phys. Rev. A vol. 39, 1809 (1989)].

Our approach deploys the nonlinear Helmholtz equation [J. Sánchez-Curto *et al.*, Phys. Rev. A vol. 85, 013836 (2012)]. We have been able to relax the strong angular constraint that is inherent to essentially all previously-published works in this arena. More specifically, we can now solve the class of problem where beam angles of incidence, reflection, and refraction may be arbitrarily large.

A compact law governing arbitrary-angle refraction will be discussed. Theoretical predictions are in excellent agreement with those obtained from exhaustive numerical simulations. Striking examples will also be given of Goos-Hänchen (GH) shifts (a phenomenon whereby, close to the critical angle of incidence, the reflected beam undergoes a displacement along the interface) [F. Goos and H. Hänchen, Ann. Phys. vol. 1, 333 (1947)]. Such shifts are an inherent property of beam-interface interactions, and they can be strongly enhanced in the presence of nonlinearity. We will report what we believe to be the largest GH shifts uncovered to date.