



University of
Salford
MANCHESTER

Helmholtz dark spatial solitons in waveguides with defocusing saturable materials

Lundie, MJ, Christian, JM, McDonald, GS and Chamorro-Posada, P

Title	Helmholtz dark spatial solitons in waveguides with defocusing saturable materials
Authors	Lundie, MJ, Christian, JM, McDonald, GS and Chamorro-Posada, P
Publication title	
Publisher	Institute of Physics
Type	Conference or Workshop Item
USIR URL	This version is available at: http://usir.salford.ac.uk/id/eprint/23004/
Published Date	2012

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: library-research@salford.ac.uk.

Helmholtz dark spatial solitons in waveguides with defocusing saturable materials

M. J. Lundie¹, J. M. Christian¹, G. S. McDonald¹, and P. Chamorro-Posada²

¹ *Materials & Physics Research Centre, University of Salford,
Greater Manchester M5 4WT, U.K.*

² *ETSI Telecomunicación, Universidad de Valladolid,
Campus Miguel Delibes Paseo Belén 15, E-47011 Valladolid, Spain*

Angular configurations play a fundamental role in essentially all nonlinear photonic architectures: from beam multiplexing applications, to scattering at a single interface, to evolution inside patterned optical structures. Equations of the nonlinear Helmholtz type are ideally suited to describing scalar oblique-propagation contexts. Knowledge of their exact solitons facilitates novel device designs, and the pursuit of these classes of solution is a key research objective of our collaboration.

Saturation under high-intensity illumination is a property of many photonic materials. Phenomenological descriptions of a saturable refractive index must go beyond polynomial-type expansions in the (local) light intensity [e.g., the cubic-quintic approximation (Pushkarov *et al.*, *Quantum Electron.* **11**, 471 (1979)), which eventually break down. However, such approaches almost always result in a governing equation that does not possess exact soliton solutions. A notable exception is the model proposed by Wood *et al.* [*Opt. Commun.* **69**, 156 (1988)].

We will present, for the first time, exact dark spatial solitons for a Helmholtz equation with a self-defocusing saturable nonlinearity. These novel solutions have been obtained by deploying a unified combination of analytical techniques (symmetry reduction, coordinate transformations, and direct integration). Multi-parameter asymptotic analysis recovers the predictions of conventional (paraxial) theory [Krolikowski and Luther-Davies, *Opt. Lett.* **18**, 188 (1993)], while convergence to its Kerr counterpart [Chamorro-Posada and McDonald, *Opt. Lett.* **28**, 825 (2003)] has been found in the limit of low light intensities. Computations involving perturbed initial-value problems have demonstrated that Helmholtz saturable dark solitons are highly robust nonlinear waves surrounded by wide basins of attraction.