



University of
Salford
MANCHESTER

Helmholtz spatial solitons: towards a theory of ultranarrow optical beams

Walsh, TM, McDonald, GS and Christian, JM

Title	Helmholtz spatial solitons: towards a theory of ultranarrow optical beams
Authors	Walsh, TM, McDonald, GS and Christian, JM
Type	Conference or Workshop Item
URL	This version is available at: http://usir.salford.ac.uk/id/eprint/23018/
Published Date	2012

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: usir@salford.ac.uk.

Helmholtz spatial solitons: towards a theory of ultranarrow optical beams

T. M. Walsh, G. S. McDonald, and J. M. Christian

Materials & Physics Research Centre, University of Salford, U.K.

Email: t.m.walsh@edu.salford.ac.uk

Keywords: scalar solitons, ultranarrow solitons, order-of-magnitude analysis

Spatial optical solitons are self-localizing and self-stabilizing beams of light that propagate with a stationary (invariant) intensity profile. Such nonlinear wave states may arise when a beam's natural tendency to diffract is exactly compensated by local refractive-index changes that are written into the host medium by the beam itself. The standard theory of 'broad' spatial solitons is based upon a Schrödinger-type picture, with all its advantages and disadvantages [2]. Our Group has refined this approach, lifting some key (e.g., angular) limitations of conventional models by retaining more natural Helmholtz-type governing equations [2]. However, there is a clear need to understand how beams behave when the envelope waist (w_0) becomes 'narrow' on the wavelength (λ) scale (see figure 1); such considerations are important in the progressive miniaturization of optical devices. Various authors throughout the world have considered this class of problem, but their approaches are often restricted in some quite subtle ways. We have recently taken steps toward constructing the first Helmholtz-based theory for ultranarrow beams by deploying an order-of-magnitude analysis of nonlinear vector Maxwell's equations. This approach appears to be fully-consistent with Helmholtz broad-beam theory [2], and lacks the unphysical features that can creep into other ultranarrow-beam models. Semi-analytical bright and dark solitons have already been derived, leading to some surprising conclusions.




Figure 1. Schematic diagram illustrating broad (left) and ultranarrow (right) beams (note that the parameter $\varepsilon \sim \lambda/w_0$).

References

- [1] Y. S. Kivshar, *Opt. Quant. Electron.* **30**, 571 (1998); Y. S. Kivshar and B. Luther-Davies, *Phys. Rep.* **298**, 81 (1998).
- [2] P. Chamorro-Posada, G. S. McDonald, and G. H. C. New, *J. Mod. Opt.* **45**, 1111 (1998).