



University of  
**Salford**  
MANCHESTER

# Algebraic soliton refraction: New wave contexts for nonlinear Snell's law

Yates, RF, Christian, JM and McDonald, GS

<b>Title</b>	Algebraic soliton refraction: New wave contexts for nonlinear Snell's law
<b>Authors</b>	Yates, RF, Christian, JM and McDonald, GS
<b>Publication title</b>	
<b>Publisher</b>	
<b>Type</b>	Conference or Workshop Item
<b>USIR URL</b>	This version is available at: <a href="http://usir.salford.ac.uk/id/eprint/29300/">http://usir.salford.ac.uk/id/eprint/29300/</a>
<b>Published Date</b>	2013

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: [library-research@salford.ac.uk](mailto:library-research@salford.ac.uk).

# Algebraic soliton refraction: new wave contexts for nonlinear Snell's law


R. F. Yates, J. M. Christian, E. A. McCoy, and G. S. McDonald

*Materials & Physics Research Centre, University of Salford, U.K.*

Email: [r.f.yates@edu.salford.ac.uk](mailto:r.f.yates@edu.salford.ac.uk)

Keywords: algebraic solitons, Snell's law, dual power-law nonlinearity, *energy theme*

The refraction of light waves at planar interfaces is a problem of fundamental interest to the optics community. To date, our Snell's law-type analyses have considered only families of *hyperbolic solitons* – robust nonlinear beams that are strongly (exponentially) localized in the transverse direction [1,2]. Here, we take the first steps toward understanding refraction of waves with much weaker transverse localization: *algebraic solitons* [3]. The stability of algebraic solitons in homogeneous systems has been investigated elsewhere [4], and those earlier results have provided a guide for material-parameter regimes in which to investigate refraction phenomena. Key research highlights include the derivation of an inhomogeneous Helmholtz governing equation for describing nonlinear beam refraction at the interface between two dissimilar dual power-law optical materials. A novel Snell's law will be reported for generalized hyperbolic solitons, with the new result for algebraic solitons emerging as a limit that maps directly onto the familiar law describing plane-wave refraction at *linear interfaces*. Computer simulations have started to test theoretical predictions in fully-nonlinear regimes, and we are also investigating Goos-Hänchen shifts (see Fig. 1).



**Figure 1.** A selection of computer simulations illustrating the Goos-Hänchen shift (denoted by  $\zeta_0$ ) for hyperbolic solitons in various power-law optical materials [2] where the nonlinearity exponent  $q$  is varied but the incidence angle remains fixed.

## References

- [1] J. Sánchez-Curto, P. Chamorro-Posada, and G. S. McDonald, *Opt. Lett.* **32**, 1126 (2007).
- [2] J. M. Christian *et al.*, *J. Atom. Mol. Opt. Phys.* **2012**, 137967 (2012).
- [3] R. W. Micallef, V. V. Afansjev, Y. S. Kivshar, and J. D. Love, *Phys. Rev. A* **54**, 2936 (1996).
- [4] J. M. Christian, G. S. McDonald, and P. Chamorro-Posada, *J. Phys. A: Math. Theor.* **43**, 085212 (2010).