



University of
Salford
MANCHESTER

Size of the abductor hallucis muscle in older women with hallux valgus

Mickle, Karen J and Nester, Christopher J

<http://dx.doi.org/10.1186/1757-1146-7-S1-A57>

Title	Size of the abductor hallucis muscle in older women with hallux valgus
Authors	Mickle, Karen J and Nester, Christopher J
Type	Article
URL	This version is available at: http://usir.salford.ac.uk/32990/
Published Date	2014

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: usir@salford.ac.uk.



MEETING ABSTRACT

Open Access

Size of the abductor hallucis muscle in older women with hallux valgus

Karen J Mickle^{1*}, Christopher J Nester²

From 4th Congress of the International Foot and Ankle Biomechanics (i-FAB) Community Busan, Korea. 8-11 April 2014

Toe deformities are highly prevalent in older people with up to 74% of older men and women having some degree of hallux valgus [1]. Despite the well documented hypotheses that atrophied, or weak toe flexor muscles are associated with the formation of toe deformities [2], there has been little evidence to support this theory. Only one study has directly compared the toe flexor strength of individuals with toe deformities to those without, revealing that older people with hallux valgus have reduced hallux strength compared to those without the deformity [3]. Therefore, to further investigate the pathomechanics of hallux valgus, this study aimed to determine whether the size of the abductor hallucis muscle differed in older women with and without hallux valgus deformity.

Forty-four older adults (60+ years) were recruited to participate in the study. Each participant had their feet assessed by the Chief Investigator (KJM), with hallux valgus severity rated using the Manchester Scale [4]. The abductor hallucis muscle was imaged using a GE Venue 40 US with a 6-9 MHz transducer [5]. Muscle cross-sectional area (CSA) was measured using Image J software with the assessor blinded to group allocation. Ten participants (all women) were classified as having moderate or severe hallux valgus and their muscle size was compared to 10 age and BMI matched women without any hallux deformity.

The older women with moderate-severe hallux valgus were found to have a significantly reduced cross-sectional area of the abductor hallucis muscle ($p < 0.05$; Figure 1). This may suggest that muscle weakness and atrophy is associated with the development or progression of hallux valgus, however further longitudinal studies are required to confirm this notion. Further research is also required to determine whether strengthening the toe flexor muscles




Figure 1 Mean (\pm SD) abductor hallucis CSA in women with and without hallux valgus deformity. * indicates significant difference.

results in hypertrophic changes to muscle morphology and these results highlight the need to investigate whether strengthening the intrinsic toe muscles could reduce the incidence and severity of toe deformities.

Authors' details

¹Biomechanics Research Laboratory, University of Wollongong, NSW, 2522, Australia. ²Centre for Health Sciences Research, University of Salford, Salford, M6 6PU, UK.

Published: 8 April 2014

References

1. Dunn JE, et al: Prevalence of foot and ankle conditions in a multiethnic community sample of older adults. *Am. J. Epidemiol* 2004, **159**(5):491-8.
2. Myerson MS, Shereff MJ: The pathological anatomy of claw and hammer toes. *J. Bone Joint Surg. Am* 1989, **71-A**(1):45-9.
3. Mickle KJ, et al: ISB Clinical Biomechanics Award 2009: Toe weakness and deformity increase the risk of falls in older people. *Clin. Biomech* 2009, **24**:787-791.
4. Garrow AP, et al: The grading of hallux valgus. The Manchester Scale. *J. Am. Podiatr. Med. Assoc* 2001, **91**(2):74-8.
5. Mickle K, et al: Reliability of ultrasound to measure morphology of the toe flexor muscles. *J Foot Ankle Res* 2013, **6**(1):12.

doi:10.1186/1757-1146-7-S1-A57

Cite this article as: Mickle and Nester: Size of the abductor hallucis muscle in older women with hallux valgus. *Journal of Foot and Ankle Research* 2014 **7**(Suppl 1):A57.

* Correspondence: kmickle@uow.edu.au

¹Biomechanics Research Laboratory, University of Wollongong, NSW, 2522, Australia

Full list of author information is available at the end of the article