



University of
Salford
MANCHESTER

Evidence for passive chemical camouflage in the parasitic mite *Varroa destructor*

Kather, R, Drijfhout, FP, Shemilt, S and Martin, SJ

<http://dx.doi.org/10.1007/s10886-015-0548-z>

Title	Evidence for passive chemical camouflage in the parasitic mite <i>Varroa destructor</i>
Authors	Kather, R, Drijfhout, FP, Shemilt, S and Martin, SJ
Type	Article
URL	This version is available at: http://usir.salford.ac.uk/34977/
Published Date	2015

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: usir@salford.ac.uk.

1 EVIDENCE FOR PASSIVE CHEMICAL CAMOUFLAGE IN THE
2 PARASITIC MITE *Varroa destructor*

3
4
5 RICARDA KATHER¹, FALKO P. DRIJFHOUT², SUE SHEMILT²
6 AND STEPHEN J. MARTIN^{3*}
7

8
9 ¹*Department of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2TN,*
10 *UK*

11
12 ²*Chemical Ecology Group, School of Physical and Geographical Sciences, Lennard-*
13 *Jones Laboratory, Keele University, Keele ST 5 5BG, UK*

14
15 ³*School of Environment and Life Sciences, The University of Salford, Manchester M5*
16 *4WT, UK*

17
18 *corresponding author

19
20 **Key words-**Varroa; Honey bees; Cuticular hydrocarbons; Mimicry; Camouflage

33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63

Abstract - Social insect colonies provide a stable and safe environment for their members. Despite colonies been heavily guarded, parasites have evolved numerous strategies to invade and inhabit these hostile places. Two common strategies are chemical mimicry via biosynthesis of the hosts' odour or chemical camouflage were compounds are acquired straight from the host. The ectoparasitic mite *Varroa destructor* feeds on the heamolymph of its honeybee host *Apis mellifera* and uses chemical mimicry to remain undetected as it lives on the adult host during its phoretic phase or while reproducing on the honeybee brood.. During the mite life cycle it switches between host adults and brood, which requires it to adjust its profile to mimic the very different odours of honeybee brood and adults. In a series of transfer experiments using adult bees and pupae, we tested whether *V. destructor* does this by synthesising compounds or using chemical camouflage. We show that *V. destructor* required direct access to the host cuticle to mimic its odour and was unable to synthesise host-specific compounds itself. Mites use chemical camouflage to mimic the host odour, even when dead, indicating a passive physico-chemical mechanism of the parasite cuticle. The chemical profile of *V. destructor* was adjusted within three to nine hours after switching hosts, demonstrating that passive camouflage is a highly efficient, fast and flexible way for the mite's to adapt to a new host's profile when moving between different host life stages, or host colonies.

64 **INTRODUCTION**

65 Colonies of social insects are a popular target of parasites, as they represent a stable
66 micro-climate, as well as, a concentrated source of food and other resources (Wilson
67 1971; Hölldobler and Wilson 1990). The main challenge for these parasites is to
68 overcome the intricate system of defence that social insects have evolved to defend their
69 colonies against such invaders. The parasite needs to enter the colony unscathed and
70 remain unharmed whilst trying to exploit its resources. Cuticular hydrocarbons (CHCs),
71 excreted as part of the insect's cuticular lipid layer, are used by many insects to recognise
72 and identify individuals around them (Howard and Blomquist 2005). In social insects
73 CHCs are species- (Martin and Drijfhout 2009) and colony-specific (Martin et al. 2013),
74 and colony members compare their own CHC profiles to those around them to detect
75 potential nest invaders; the first line of colony defence.

76 Many social parasites have evolved strategies to evade this chemical recognition
77 system (Dettner and Liepert 1994; Lenoir et al. 2001; Bagnères and Lorenzi 2010) either
78 through biosynthesis of host CHCs (chemical mimicry *sensu stricto*: Akino et al. 1999;
79 Howard et al. 1982, 1990; Lenoir et al. 1997; Lenoir et al., 2001), or by reducing their
80 own CHC profile to undetectable levels (chemical insignificance: wasps: Lorenzi and
81 Bagnères 2002; parasitoids: Kroiss et al. 2009), using a combination of both (Jeral et al.
82 1997; Uboni et al. 2012) or acquiring compounds straight from the host (chemical
83 camouflage: beetles: Vander Meer and Wojcik 1982; parasitoid: Akino et al. 1999;
84 spiders: von Beeren and Witte 2012). Cini et al. (2011) showed that it is CHCs in
85 particular rather than other, co-occurring compounds that are important in the detection of
86 the parasite by the host.

87 Many parasites interact with their host inside the nest and use chemical mimicry
88 to be perceived and treated as if they were a member of the colony (Lenoir et al. 2001).
89 Some parasites mimic a particular caste (Akino and Yamaoka 1998) or gender (Hojo et
90 al. 2009) of their host to increase acceptance and trick the host into feeding them.
91 Nevertheless, very little is known about the chemical strategies used by parasites that
92 exclusively feed and reproduce on their hosts. The ectoparasitic mite *Varroa destructor* is
93 a major pest of the European honey bee *Apis mellifera* and has a very close relationship
94 with its host. This mite uses 'drifting' bees to infest new colonies and, once inside the

95 colony, it switches to nurse bees to increase its likelihood of reaching the brood area,
96 where it reproduces (Kraus et al. 1986). To reproduce the mite moves from the adult
97 nurse bee onto a final staged larva. Once the brood cell has been sealed the mite emerges
98 from beneath the larva and starts its reproductive cycle feeding on the developing pupa.
99 When the pupae emerges from the brood cell as adult bee, the foundress mite and her
100 mature offspring are also released. The males soon perish but the females enter their
101 phoretic phase, feeding on the haemolymph of the adults (Rosenkranz et al., 2010). There
102 are no behavioural interactions between the mite and its host, but instead *V. destructor*
103 blends in with the bee's body chemistry to become chemically invisible to the host as the
104 mite is carried around the colony (Nation et al. 1992). Honeybee CHC profiles vary
105 depending on the age and gender of the bee (Nation et al. 1992, Arnold 2000, Aumeier et
106 al. 2002, Kather et al. 2010) and results published by Nation et al. (1992) and Martin et
107 al. (2001) suggest that *V. destructor* adjusts its mimicry to match these differences as it
108 switches hosts, which indicates that the mite does not use chemical insignificance to
109 remain undetected in the host colony.

110 Here we investigated the mechanism through which *V. destructor* is able to mimic
111 the CHC profile of its host and tested whether the mite uses chemical camouflage or
112 biosynthesis to achieve its chemical mimicry. Because of the high death rate of bees
113 associated with radio labelling of CHCs in *A. mellifera* (Gary Blomquist and Falko
114 Drijfhout, *pers. comm.*), we used the speed at which *V. destructor* adjusts its mimicry to a
115 new host as an indicator for whether any changes in CHC profile were due to
116 biosynthesis of compounds (true chemical mimicry) or the mite transferring compounds
117 straight from the host (chemical camouflage). Studies have shown that once biosynthesis
118 of new compounds has started it takes one to several days for these compounds to reach
119 the cuticle (de Renobales et al. 1988, Ichinose and Lenoir 2009). Therefore, we assumed
120 that if *V. destructor* synthesised compounds itself it would take at least one day to detect
121 any significant change in the mite's CHC profile. However, if the mite transferred host
122 compounds onto its own cuticle, we can expect to see changes in the mite's CHC profile
123 within a few hours of exposing it to the new host. Fortunately, the CHC profiles of adult
124 bees are rich in alkenes whereas pupae profiles are rich in methylalkanes, thus allowing
125 easy quantification of changes in the mite's cuticular chemistry when mites are moved

126 between these two honeybee life stages. We therefore, measured the speed at which the
127 mite was able to adjust its profile to that of its new host.

128 In addition, we predicted that if *V. destructor* used chemical camouflage, it would
129 need direct physical contact with the host's cuticular lipid layer to be able to transfer it
130 onto its own cuticle. However, if mites synthesised their CHC, they may be able to adjust
131 their profiles without direct cuticular contact with the host or access to host's hemolymph.
132 Therefore, we conducted a series of additional experiments where we again moved mites
133 from adult bees onto pupae, but this time we restricted their access to the host cuticular
134 lipid layer or the host hemolymph or both to test if this affected the ability of the mites to
135 adjust their profile.

136 Finally if *V. destructor* uses chemical camouflage to mimic its host, how is this
137 achieved given that several other parasites actively groom their host to transfer
138 compounds but such behaviour is not seen in *V. destructor*. To investigate this we
139 transferred dead mites that had their own CHC profile removed chemically onto bee and
140 fly pupae and observed if mites could still acquire the pupa's CHC profile.

141

142 **METHODS**

143 *Sample Collection* - For all experiments, mites were collected from adult bees (i.e. nurse
144 bees) by removing brood frames from a hive and covering the bees on the frame with
145 icing sugar. Brood frames were returned to the hive and a Varroa board was inserted at
146 the bottom of the hive to catch the fallen mites. These were removed from the board
147 using a fine moist brush, which was cleaned with water and dried after each mite. Mites
148 were gently wiped to remove excess sugar and placed in Eppendorf tubes. As expected,
149 mites collected this way had a CHC profile very similar to that of adult (nurse) bees (Fig.
150 1) and were used as a starting point for all treatment groups.

151 Adult bees and pupae of the same hive were collected prior to sugaring to allow
152 for mite-bee comparison later on. The pupae collected were of the white-eye stage i.e. 3-4
153 days after the cell had been sealed. All treatment groups were kept at 34°C and 70%
154 humidity. During experiments, mites were kept in 0.6ml microcentrifuge tubes that had
155 been prepped with a wet cotton ball at the bottom to maximise humidity inside the vial
156 and a hole was created in the lid to allow for oxygen exchange. Each treatment group

157 contained eight mites per replicate and ten replicates per group. At the end of the
158 experiment, mites were stored at -20°C for chemical analysis later. Only mites that were
159 still alive at the end of the experiment were used for chemical analysis (refer to Table 1
160 for the total number of mites per treatment group that were used for chemical analysis).
161 The CHC profiles of bee pupae were also quantified to control for any changes in mite
162 CHC chemistry caused by inter-pupae differences rather than due to a treatment effect.

163

164 *Speed of host CHC acquisition* - The first experiment investigated the speed at which *V.*
165 *destructor* is able to adjust its chemical mimicry to that of a new host. Mites were
166 transferred from adult bees (alkene rich) onto pupae (methylalkanes rich), and kept on the
167 pupae for 20 mins, 1 hr, 3 hrs, 9 hrs or 27 hrs to measure the speed at which *V. destructor*
168 acquires a new host profile. A total of 460 mites were collected, of which 60 were frozen
169 straight away to serve as starting point (0 hrs control). The remaining 400 mites were
170 transferred to microcentrifuge tubes for the various time periods, with one pupa added to
171 each tube.

172

173 *Limited access to host cuticular lipid layer* - The second experiment tested whether *V.*
174 *destructor* needs access to the host cuticular lipid layer or access to the host hemolymph
175 to adjust its mimicry. Mites were moved from adult bees to pupae and limited in their
176 access to either a) the host's lipid layer or b) the host's hemolymph and lipid layer. A
177 total of 400 mites were transferred to adult bees kept in bee cages and allowed to feed for
178 24 hrs to control for any effects of starvation during the experiment. After 24 hrs, the
179 mites were transferred to microcentrifuge tubes which had been prepped with four rows
180 of holes along the sides of the tube. This allowed mites to perceive the smell of any pupae
181 placed immediately around the vial without allowing direct access to these pupae. The
182 tubes were then placed in a petri dish, with 2-3 tubes per dish.

183 Four treatment groups were created: 1) Mites were kept on one pupa per tube to
184 imitate a natural situation (control group), where mites have access to the host's cuticular
185 lipid layer and its hemolymph. 2) Mites were able to smell the host but were denied
186 access to its hemolymph and cuticular lipid layer. In this group, pupae were placed
187 around the (perforated) tubes which prevented the mites touching the pupae. 3) Mites had

188 access to host hemolymph but not the host cuticular lipid layer. To achieve this, the mites
189 were kept on a pupa that had been stripped off its cuticular lipid layer by washing the
190 pupae in high-performance liquid chromatography (HPLC) grade hexane for 3 min. and
191 leaving it to dry on a clean glass plate for 30 min. We had already confirmed by GC-MS
192 that all CHCs had successfully been removed. Because the washed pupae were now
193 odourless, unwashed pupae were placed around the microcentrifuge tubes containing the
194 washed pupae and mites to ensure that the pupaa odour was present. In the tubes the
195 mites still fed on the washed pupa and there were no obvious differences in mite
196 behaviour between this treatment group and the control group. 4) The final mite group
197 was kept in full isolation with no pupae added to or placed around the tube, to serve as
198 negative control (full isolation from the host). All treatment groups were left for 18 hours
199 before mites and pupae were retrieved and frozen prior to chemical analysis.

200 There was clear behavioural evidence that mites were able to perceive the odour
201 of the pupae placed around the tubes, as mites started to accumulate around the holes
202 along the tube wall, as soon as pupae were placed in the petri dish. This behaviour was
203 not observed in the isolated mite group, where no pupae were placed around the tube.
204

205 *Limited access to host hemolymph* - To create a scenario in which mites had access to the
206 host's cuticular lipid layer but could not feed on the host hemolymph, we maintained *V.*
207 *destructor* on mature fly pupae (*Calliphora vomitoria*) that had been stripped off their
208 own CHCs (washed in HPLC grade hexane for 3 min.) and then spiked the fly pupae with
209 the odour of a bee pupae. Each fly pupa was covered with the extract of one bee pupa
210 (washed in HPLC grade hexane for 15 min.) by gradually dripping the extract onto the
211 fly's cuticle and leaving it to dry over a 30 min. period. (Fig. 1b). The cuticle of a mature
212 fly pupae is too thick for the mite to penetrate; hence mites can only access the artificial
213 lipid layer but not the hemolymph. For this experiment a total of 220 mites were
214 collected, of which 60 were frozen straight away to serve as starting points. 80 mites were
215 kept on fly pupae and the remaining 80 were kept on bee pupae (positive control) in
216 microcentrifuge tubes prepped with a wet cotton ball and one hole in the lid as described
217 above. All treatment groups were left for 18 hours before mites and pupae were retrieved
218 and frozen prior to chemical analysis.

219

220 *Active versus passive transfer of host CHCs* – To investigate whether *V. destructor* uses
221 passive CHC transfer as an alternative strategy to actively grooming the host, we tested
222 whether dead mites were still able to mimic a host odour. Mites were killed by freezing at
223 -20°C for 1 hr. They were then defrosted for 10 mins, then stripped of their CHCs by
224 immersing them in HPLC grade hexane for 3 min. and leaving them to dry for 30 min. 80
225 of these dead mites were transferred to petri dishes containing white-eye pupae. The dead
226 mites were placed onto pupae with their legs touching the host cuticle to imitate the
227 natural position of the mite on the host. As positive control, 80 live mites were transferred
228 to petri dishes containing white-eye pupae. Mites were again left for 18 hrs and frozen for
229 chemical analysis. A sub-set of washed mites tested by GC-MS confirmed that all CHCs
230 had been successfully removed from the dead mites prior to the transfer onto the host.

231

232 *Chemical and Statistical Analysis* - Samples were extracted in HPLC hexane containing a
233 C_{20} standard (1mg/100ml hexane). Each bee sample was extracted in 0.5 ml hexane, and
234 mite samples were extracted in pools of six mites in 300 μl hexane per sample. Samples
235 were left at room temperature for 15 min., before transferring 30 μl of extract to a glass
236 insert and left to evaporate overnight.

237 Samples were re-suspended in 30 μl hexane and analysed on an HP6890GC
238 (equipped with an HP-5MS column; length: 30m; ID: 0.25mm; film thickness: 0.25 μm)
239 connected to an HP5973 MSD (quadrupole mass spectrometer with 70-eV electron
240 impact ionization). Samples were injected in the splitless mode. The oven was
241 programmed from 70°C to 200°C at $40^{\circ}\text{C}/\text{min}$ and then from 200°C to 320°C at $25^{\circ}\text{C}/\text{min}$
242 and, finally, held for 5 min at 350°C . The carrier gas helium was used at a constant flow
243 rate of 1.0ml min^{-1} . Compounds were identified using standard MS databases, diagnostic
244 ions and Kovats indices.

245 The peak area of each compound was determined by manual integration of each
246 total ion chromatogram (TIC), which was then translated into actual concentration
247 (mg/ml hexane) using the peak of the C_{20} standard. Compounds which on average
248 contributed less than 1% to the overall chemical profile (i.e. n-alkanes + alkenes +

249 methylalkanes) were excluded from the analysis. Compounds were grouped into three
250 main chemical classes: *n*-alkanes, alkenes and methylalkanes.

251 Because the CHC profiles of adult bees and bee pupae mainly differ in their
252 alkene:methylalkane ratio, we focused our analysis on these two CHC families and
253 excluded the *n*-alkanes. For the speed of host CHC acquisition, a regression analysis was
254 conducted to investigate whether the alkene:methylalkane ratio changed significantly
255 during the 27- hour-period after the mite had been placed onto the new host. For all other
256 experiments, one-way ANOVA's and *post-hoc* Tukey tests were carried out to test for
257 significant changes in the alkene:methylalkane ratio between treatment groups. The
258 assumptions associated with doing an ANOVA test were checked prior to doing the
259 analysis. All tests were performed using the statistical software R (v 2.81).

260

261

262

263 RESULTS

264 *Speed of host CHC acquisition* - As expected, the CHC profiles of honey bee pupae had a
265 high methylalkane: alkene ratio that was reversed in the adult bees (Fig. 2a). Hence, mites
266 collected from adult bees, at the beginning of the experiment, had high concentrations of
267 alkenes but low levels of methylalkane. When mites were then transferred onto a pupa,
268 their methylalkane levels increased followed by a drop in alkene concentration over the
269 27 hrs during which mites were exposed to the new host. This led to a significant drop in
270 the alkene: methylalkane ratio (Fig. 2b) over time, which levelled off after 9 hours of
271 exposure to the host (Polynomial Regression: $y=1.5 - 3.795 x + 3.265 x^2$, $F=26$, d.f. = 2,
272 52, $p < 0.0001$). A significant increase in mite methylalkane concentration was already
273 visible after the first 20 min. of being on the new host (ANOVA, $F=4$, d.f.=5,49,
274 $p<0.001$), which was followed by a second increase after 3 hrs (ANOVA, $F=4$, d.f.=5,49,
275 $p<0.004$) (Fig.2b). Alkene concentration remained constant for the first 3 hours and then
276 dropped significantly (ANOVA, $F=10$, d.f.=5,49, $p<0.001$).

277

278 *Limited access to host cuticular lipid layer* - The treatment groups of the second
279 experiment differed significantly in their alkene: methylalkane ratio (ANOVA, $F=30.31$,

280 d.f.=4,43, $p < 0.001$) (Fig. 3a). As expected, mites that were kept on (unwashed) bee
281 pupae, and hence had access to both host hemolymph and the host's cuticular lipid layer,
282 had significantly higher levels of methylalkane compared to mites kept on adult bees
283 (*post-hoc* Tukey test: $p < 0.001$). The second mite group (Odour), which was able to
284 smell the host through the perforated tube but had no direct physical access to host
285 hemolymph or cuticle, was unable to increase their methylalkane levels so their alkene:
286 methylalkane ratio remained the same as mites kept in full isolation (*post-hoc* Tukey test:
287 $p = 0.92$). When mites only had access to host hemolymph by keeping them on washed
288 pupae (Pupa + Odour – CHCs), these were also unable to increase their methylalkane
289 concentration to match the CHC profile of the host. Instead their alkene: methylalkane
290 ratio was again the same as mites kept in full isolation (*post-hoc* Tukey test: $p = 0.27$)
291 (Fig 3a). When mites were isolated from the host pupae the concentration of all CHCs
292 decreased (Fig. 3a). Therefore, access to host odour or host hemolymph alone was
293 insufficient for mites to mimic their new host profile.

294

295 *Limited access to host hemolymph* - Mites were able to increase their methylalkane levels
296 when given an artificial lipid layer (fly pupa spiked with the CHC profile of one bee
297 pupa) even though access to host hemolymph was denied. Their alkene:methylalkane
298 ratio was the same as that of mites kept on bee pupae (ANOVA, $F=86.15$, d.f.=2,22, $p <$
299 0.001 ; *post-hoc* Tukey test: $p = 0.31$) and significantly different from that of mites kept
300 on adult bees (*post-hoc* Tukey test: $p < 0.001$) (Fig. 3b). This indicates that mites are able
301 to adjust their chemical profile as long as they have direct access to the host's cuticular
302 lipid layer, even if host hemolymph cannot be accessed.

303

304 *Active versus passive transfer of host CHCs* - Dead (washed) mites were still able to
305 increase their methylalkane levels to the degree that they were significantly different in
306 their alkene:methylalkane ratio compared to mites kept on adult bees (ANOVA, $F=52.01$,
307 d.f.=2,22, $p < 0.001$; *post-hoc* Tukey test: $p < 0.001$) (Fig. 3c). Even though dead mites
308 were similar in methylalkane concentration to live control mites (ANOVA, $F=2.33$,
309 d.f.=1,16, $p = 0.15$), they had significantly higher levels of alkene, which led to a
310 significant difference in alkene:methylalkane ratio in these two groups (*post-hoc* Tukey

311 test: $p < 0.001$). This disproportionate increase in alkene levels was not due to alkene
312 residues left after washing mites, because washed mites that were analysed straight away
313 (Control Mites) had no CHCs left on their cuticle after washing. The fact that dead mites
314 were still able to adsorb host CHCs indicates that at least the uptake of host CHCs is a
315 passive process.

316

317 **DISCUSSION**

318 The results indicate that *V. destructor* uses chemical camouflage to mimic the odour of its
319 bee host. Without access to the host's cuticular lipid layer the mite was unable to adjust
320 its mimicry to a new host, even when access to host hemolymph was provided. This
321 finding suggests that *V. destructor* is unable to synthesize host-specific CHCs but instead
322 there is a transfer of host CHCs from the host's cuticle to that of the mite; this even
323 occurs when the mite is dead. This is further supported that changes in the mites profile
324 was observed after just 20 mins. The rapid adsorption of a host's CHCs has been observed
325 in a number of parasites and can be facilitated by an active process such as grooming the
326 host (ants: Lenoir et al. 1997; Franks et al. 1990; spiders: von Beeren and Witte 2012;
327 silverfish: von Beeren et al. 2011), in addition to any simple passive processes were close
328 contact with the host cuticle is required (cockroaches: Everaerts et al. 1997; termites:
329 Vauchot et al. 1998; beetles: Vander Meer and Wojcik 1982). For example, the
330 myrmecophile 'shampoo' ant *Formicoxenus provancheri* grooms its host, *Myrmica*
331 *alaskensis*, around 45% of its time inside the nest (Lenoir et al. 2001), while the
332 myrmecophile beetle *Myrmecaphodius excavaticollis* passively adsorbs host CHCs
333 without any obvious grooming behaviour (Vander Meer and Wojcik 1982). Further
334 evidence for the passive transfer of CHCs has come from using artificially mixed
335 colonies of *Reticulitermes* termites, where *R. santonensis* and *R. lucifugus grassei*
336 acquired each other's CHCs without any allogrooming taking place (Vauchot et al. 1998).
337 In both studies, *M. excavaticollis*, *R. santonensis* and *R. lucifugus grassei* are all able to
338 take up host compounds even if dead individuals were used, thus excluding the
339 possibility of biosynthesis or behavioural acquisition as relevant factors.

340 Our findings suggest that *V. destructor* also uses passive transfer (i.e., transfer of
341 host CHCs without the parasite actively grooming or in any way interacting with the

342 host) of host CHCs to camouflage itself on the host, because dead mites were still able to
343 adsorb compounds when placed on the host. The mite's legs are too short to spread host
344 CHCs across its cuticle via grooming and, hence, passive CHC transfer is an efficient
345 strategy to facilitate the mite's chemical mimicry. When switching host stages, passive
346 chemical camouflage allows the mite CHC profile to rapidly adjust within a few hours to
347 match the chemical profile of its new host. So, when moving from an adult bee to a bee
348 pupa, the mite quickly adopts the profile of its new host. This process is aided by the fact
349 that the mite naturally seems to lose compounds when these are not provided through
350 contact with the host cuticular lipid layer, as was apparent when mites were kept in full
351 isolation. This was also observed in the myrmecophile beetle *Myrmecaphodius*
352 *excavaticollis* (Vander Meer and Wojcik 1982), as the termites *Reticulitermes*
353 *santonensis* and *R. lucifugus grassei* (Vauchot et al. 1998), where CHC transfer could
354 already be observed after just two hours of cohabitation, but acquired CHCs started to
355 decrease in concentration as soon as individuals were isolated from their host.

356 It was surprising that dead *V. destructor* mites preferentially adsorb alkenes
357 compared to live mites. Alkenes have a lower melting temperature in comparison to
358 methylalkanes (Gibbs 2002) and, thus, it is possible that these transfer more readily than
359 methylalkanes. If it was easier for mites to adsorb alkenes compared to methylalkanes,
360 this would have also been observed in the live mites unless the adjustment of the exact
361 alkene:methylalkane ratio is actively regulated by the mite. This result certainly requires
362 further investigation.

363 There is some evidence to suggest that the ability to passively adsorb CHCs may
364 be species-specific in arthropods. When *Reticulitermes santonensis* and *R. l. grassei* are
365 kept in (artificial) mixed colonies, the latter takes up more compounds of the former than
366 *vice versa* (Vauchot et al. 1998). A similar case was reported by Vienne et al. (1990 cited
367 in Vauchot et al., 1998), whereas *Formica selysi* acquired more allospecific CHC than
368 *Myrmica rubida* when these two ant species lived as a mixed colony. The fact that there
369 seems to be a species-specific ability of taking up CHCs indicates that there may be
370 differences in cuticular physiology or chemistry that allow some species to take up CHCs
371 more readily than others. If this occurs in non-parasitic arthropods, it is not surprising that
372 a range of parasitic arthropods have evolved the ability to soak up enough host CHCs to

373 achieve chemical camouflage. The physico-chemical mechanism that allows them to do
374 this however remains elusive.

375 Our results confirmed the findings reported in Nation et al. (1992) that the
376 chemical mimicry of *V. destructor* changes as the parasite switches hosts. The mite *V.*
377 *destructor* is the first case known to us where a parasite frequently switches its
378 camouflage as it moves between host stages. Because groups of individuals within an *A.*
379 *mellifera* colony can vary significantly in their CHC profile, due to task (Kather et al.
380 2011) or life stage, a passive chemical camouflage is likely to be the best strategy to
381 facilitate a quick adjustment in the parasite's camouflage to match the new host with
382 minimal energetic cost. The mite also has a number of appendages such as suckers, hairs
383 and its crab-like carapace (Rosenkranz et al., 2010) that allow it to hold on to the host
384 during the transition time and, this way, these appendages buy the mite time until it is
385 fully blended in with the host's CHC profile, which only takes a few hours. This study
386 helps explain why *V. destructor* has become one of the most widespread and successful
387 ecto-parasitic pests of honey bees.

388

389 **ACKNOWLEDGEMENTS** - We thank the Kings Lynn, Sheffield and Barnsley
390 Beekeepers for providing mites and Roger Butlin of Sheffield University for comments
391 on the manuscript. This research was funded by funding from BBSRC (BB/G017077/1),
392 the East Anglian Beekeepers (EARS) and NERC (NE/F018355/1).

393

394

395

396

397

398

399

400

401

402

403

404 **REFERENCES**

- 405 AKINO, T. and YAMAOKA, R. 1998. Chemical mimicry in the root aphid parasitoid
406 *Paralipsis eikoeae* Yasumatsu (Hymenoptera: Aphidiidae) of the aphid-attending ant
407 *Lasius sakagami* Yamauchi & Hayashida (Hymenoptera: Formicidae). *Chemoecology*
408 8:153–161.
- 409
- 410 AKINO, T., KNAPP, J.J., THOMAS, J.A. and ELMES, G.W. 1999. Chemical mimicry and
411 host specificity in the butterfly *Maculinea rebeli*, a social parasite of *Myrmica* ant
412 colonies. *Proc. Roy. Soc. B* 266:1419–1426.
- 413
- 414 ARNOLD, G., QUENET, B. and MASSON, C. 2000. Influence of Social Environment on
415 Genetically Based Subfamily Signature in the Honeybee. *J. Chem. Ecol.* 26: 2321–2333.
- 416
- 417 AUMEIER, P., ROSENKRANZ, P. and FRANCKE, W. 2002. Cuticular volatiles, attractivity of
418 worker larvae and invasion of brood cells by *Varroa* mites. A comparison of Africanized
419 and European honey bees. *Chemoecology* 12: 65–75.
- 420
- 421 BAGNÈRES, A-G. and LORENZI, M.C. 2010. Chemical deception/mimicry using cuticular
422 hydrocarbons. In: G.J. Blomquist and A-G. Bagnères (Eds.) *Insect Hydrocarbons:*
423 *Biology, Biochemistry, and Chemical Ecology*. Cambridge, University Press. pp. 282-
424 324.
- 425
- 426 CINI, A., BRUSCHINI, C., SIGNOROTTI, L., PONTIERI, L., TURILLAZZI, S. and CERVO, R.
427 2011. The chemical basis of host nest detection and chemical integration in a cuckoo
428 paper wasp. *J. Experimental Biol.* 214: 3698–3703.
- 429
- 430 DETTNER, K. and LIEPERT, C. 1994. Chemical mimicry and camouflage. *Ann. Rev.*
431 *Entomol.* 39: 129–154.
- 432
- 433 EVERAERTS, C., FARINE, J. P. and BROSSUT, R. 1997. Changes of species specific
434 cuticular hydrocarbon profiles in the cockroaches *Nauphoeta cinerea* and *Leucophaea*

435 *maderae* reared in heterospecific groups. *Entomologia Experimentalis et Applicata* 85:
436 145–150.
437
438 FRANKS, N., BLUM, M. S., SMITH, R. and ALLIES, A. B. 1990. Behavior and chemical
439 disguise of cuckoo ant *Leptothorax kutteri* in relation to its host *Leptothorax acervorum*.
440 *J. Chem. Ecol.* 16: 1431–1444.
441
442 GIBBS, A. 2002. Lipid melting and cuticular permeability: new insights into an old
443 problem. *J. Insect Physiol.* 48: 391-400.
444
445 HOJO, M. K., WADA-KATSUMATA, A., AKINO, T., YAMAGUCHI, S., OZAKI, M. and
446 YAMAOKA, R. 2009. Chemical disguise as particular caste of host ants in the ant inquiline
447 parasite *Niphanda fusca* (Lepidoptera: Lycaenidae). *Proc. Roy. Soc. B* 276: 551–558.
448
449 HÖLLDOBLER, B. and WILSON, E.O. 1990. *The Ants*. Cambridge, Harvard University
450 Press.
451
452 HOWARD, R.W., MCDANIEL, C. A. and BLOMQUIST, G. J. 1982. Chemical mimicry as an
453 integrating mechanism for three termitophiles associated with *Reticulitermes virginicus*
454 (Banks). *Psyche* 89: 157-167.
455
456 HOWARD, R.W., STANLEY-SAMUELSON, D.W. and AKRE, R. D. 1990. Biosynthesis and
457 Chemical Mimicry of Cuticular Hydrocarbons from the Obligate Predator, *Microdon*
458 *albicomatus* Novak (Diptera: Syrphidae) and Its Ant Prey, *Myrmica incompleta*
459 Provancher (Hymenoptera: Formicidae). *J. Kansas Entomol. Soc.* 63: 437-443.
460
461 HOWARD, R.W. and BLOMQUIST, G.J. 2005. Ecological, behavioral, and biochemical
462 aspects of insect hydrocarbons. *Ann. Rev. Entomol.* 50: 371–393.
463

464 ICHINOSE, K and LENOIR, A. 2009. Ontogeny of hydrocarbon profiles in the ant
465 *Aphanogaster senilis* and effects of social isolation. *Comptes Rendues Biologies* 332:
466 697-703.
467

468 JERAL, J.M., BREED, M.D. and HIBBARD, B. E. 1997. Thief ants have reduced quantities
469 of cuticular compounds in a ponerine ant, *Ectatomma ruidum*. *Physiol. Entomol.* 22: 207–
470 211.
471

472 KATHER, R., DRIJFHOUT, F. P. and MARTIN, S. J. 2011. Task group differences in
473 cuticular lipids in the honey bee *Apis mellifera*. *J. Chem. Ecol.* 37:205-212.
474

475 KRAUS, B. KOENIGER, N. and FUCHS, S. 1986. Unterscheidung zwischen Bienen
476 verschiedenen Alters durch *Varroa jacobsoni* Oud. und Bevorzugung von
477 Ammenbienen Volk. *Apidologie* 17: 257–266.
478

479 KROISS, J., SCHMITT, T. and STROHM, E. 2009. Low level of cuticular hydrocarbons in a
480 parasitoid of a solitary digger wasp and its potential for concealment. *Entomol. Sci.* 12:
481 9–16.
482

483 LENOIR, A., MALOSSE, C. and YAMAOKA, R. 1997. Chemical mimicry between parasitic
484 ants of the genus *Formicoxenus* and their host *Myrmica* (Hymenoptera, Formicidae).
485 *Biochem. Systematics Ecol.* 25: 379–389.
486

487 LENOIR, A., D’ETTORRE, P., ERRARD, C. and HEFETZ, A. 2001. Chemical ecology and
488 social parasitism in ants. *Ann. Rev. Entomol.* 46: 573–599.
489

490 LORENZI, M.C. and BAGNÈRES, A.G. 2002. Concealing identity and mimicking hosts: a
491 dual chemical strategy for a single social parasite? (*Polistes atrimandibularis*,
492 Hymenoptera: Vespidae). *Parasitology* 125: 507–512.
493

494 MARTIN, C. SALVY, M., PROVOST, É.M., BAGNÉRES, A.G., ROUX, M., CRAUSER, D.,
495 CLEMENT, J.L. and LE CONTE, Y. 2001. Variations in chemical mimicry by the
496 ectoparasitic mite *Varroa jacobsoni* according to the developmental stage of the host
497 honey-bee *Apis mellifera*. *Insect Biochem. Mol. Biol.* 31:365-379.

498
499 MARTIN, S. J., and DRIJFHOUT, F. P. 2009. A review of ant cuticular hydrocarbons *J.*
500 *Chem. Ecol.* 35:1151-1161.

501
502 MARTIN, S. J., VITIKAINEN, E., SHEMILT, S., DRIJFHOUT, F. P. and SUNDSTROM, L. 2013.
503 Sources of variation in cuticular hydrocarbons in the ant *Formica exsecta*? *J. Chem. Ecol.*
504 39: 1415-1423.

505
506 NATION, J.L. SANFOR, M.T. and MILNE, K. 1992. Cuticular hydrocarbons from *Varroa*
507 *jacobsoni*. *Experimental Appl. Acarol.* 16: 331–344.

508
509 DE RENOBABLES, M., NELSON, D. R., MACKAY, M. E., ZAMBONI, A. C. and BLOMQUIST, G.
510 J. 1988. Dynamics of hydrocarbon biosynthesis and transport to the cuticle during pupal
511 and early adult development in the cabbage looper *Trichoplusia ni* (Lepidoptera:
512 Noctuidae). *Insect Biochem.* 18: 607-613.

513
514 ROSENKRANZ, P., AUMEIER, P. and ZIEGELMANN, B. 2010. Biology and control of *Varroa*
515 *destructor*. *J. Invert. Pathol.* 103: S96–S119.

516
517 UBONI, A., BAGNÈRES, A-G., CHRISTIDÈS, J-P. and LORENZIET, M. G. 2012.
518 Cleptoparasites, social parasites and a common host: Chemical insignificance for visiting
519 host nests, chemical mimicry for living in. *J. Insect Physiol.* 58: 1259–1264.

520
521 VANDER MEER, R.K. and WOJCIK, D.P. 1982. Chemical Mimicry in the Myrmecophilous
522 Beetle *Myrmecaphodius excavaticollis*. *Science* 218: 806–808.

523

524 VAUCHOT, B., PROVOST, E., BAGNÈRES, A-G., RIVIERE, G., ROUX, M. and CLÉMENT, J-L.
525 1998. Differential Adsorption of Allospecific Hydrocarbons by the cuticles of Two
526 Termite Species, *Reticulitermes santonensis* and *R. lucifugus grassei*, Living in a Mixed
527 Colony. *J. Insect Physiol.* 44: 59-66.
528
529 VON BEEREN, C., SCHULZ, S. HASHIM, R. and WITTE, V. 2011. Acquisition of chemical
530 recognition cues facilitates integration into ant societies. *BMC Ecol.* 11:30.
531
532 VON BEEREN, C., HASHIM, R. and WITTE, V. 2012. The social integration of a
533 myrmecophilous spider does not depend exclusively on chemical mimicry. *J. Chem.*
534 *Ecol.* 38: 262–271.
535
536 WILSON, E.O. 1971. *The Insect Societies*. Harvard, Belknap Press.
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553

554 **Table 1** Total number of mite samples (n) per treatment group across all experiments. For
 555 each mite sample, a pool of six mites were extracted and treated as one sample in the data
 556 analysis.

557
 558

Experiment	Treatment Group	n
<i>Experiment 1</i>		
(Speed of host CHC acquisition)		
	0 hrs	10
	20 min.	8
	1 hr	10
	3 hrs	8
	9 hrs	10
	27 hrs	9
<i>Experiment 2</i>		
(Limited access to host cuticular lipid layer)		
	Mites (Adult Bee)	10
	Mites (Bee Pupa)	8
	Mites (Pupa + Odour – CHCs)	10
	Mites (Odour)	6
	Full Isolation	10
<i>Experiment 3</i>		
(Limited access to host hemolymph)		
	Mites (Adult Bee)	6
	Mites (Fly Pupa + Bee Pupa CHCs)	10
	Mites (Bee Pupa)	9
<i>Experiment 4</i>		
(Active vs. passive transfer)		
	Mites (Adult Bee)	10
	Dead Mites (Bee Pupa)	8
	Live Mites (Bee Pupa)	10

559
 560
 561
 562
 563
 564
 565

566 **Fig. 1** CHC profiles of a) a honey bee pupa and b) a washed fly pupa spiked with the
567 CHC extract of one bee pupa. The main CHCs are: 1) n-C₂₃, 2) C_{25:1}, 3) n-C₂₅, 4) 9-, 11-,
568 13-meC₂₅, 5) C_{27:1}, 6) n-C₂₇, 7) 9-, 11-, 13-meC₂₇, 8) C_{29:1}, 9) n-C₂₉, 10) 9-, 11-, 13-
569 meC₂₉, 11) C_{31:1}, 12) n-C₃₁, 13) 9-, 11-, 13-meC₃₁, 14) C_{33:1}.

570


571 **Fig. 2** Speed of chemical mimicry. Mites were collected from adult bees (time: 0hrs) and
572 kept on pupae for different time intervals. Bee profiles are shown in panel a) and mite
573 profiles in panel b). Mites acquired pupa-specific methylalkanes and lost adult-specific
574 alkenes within 3-9 hours of exposure to the new host, leading to a drop in alkene-
575 methylalkane ratio. Error bars show one standard deviation and sample sizes are given in
576 Table 1.

577

578 **Fig. 3** Alkene and methylalkane concentrations of mite CHC profiles across experiments
579 2 to 4. In each panel 'Adult bees' represents the starting CHC profile of mites i.e. kept on
580 adult bees, and 'Bee pupae' is the positive control i.e. mites kept on untreated pupae. a)
581 *experiment 2*; mites have access to the host cuticular layer. Pupa + Odour - CHC - are
582 mites on washed bee pupae i.e. CHC removed but surrounded by unwashed pupae to
583 provide host odour; Odour- are mites only exposed to pupae odour, and Full Isolation- is
584 the negative control since mites were isolated from the host. b) *experiment 3*; excludes
585 access to host haemolymph; Fly Pupa + Bee Pupa CHCs - are mites kept on fly pupae
586 spiked with a bee pupa CHC profile. c) *experiment 4*; transfer of host CHCs was active or
587 passive; Dead mites- their CHC profile removed and placed on living pupa, Live mites
588 also had their CHC removed but not killed, while control mites had their CHC removed
589 and were isolated from pupa and showed that chemical washing removed all the mites
590 CHC. For each experiment, treatment groups differed in their alkene:methylalkane ratio if
591 they have different letters, whereas groups with the same letter were similar in their
592 alkene:methylalkane ratio. Error bars show the standard error and samples sizes are given
593 in Table 1.

594

595



596


597 Fig. 1

598

599

600

601



602

603

604 Fig. 2


605

606

607

608

609



610 Fig. 3