



University of  
**Salford**  
MANCHESTER

# Postural control during quiet standing and voluntary stepping response tasks in individuals post-stroke : a case-control study

Moisan, G, Chayasit, P, Boonsinsukh, R, Nester, CJ and Hollands, K

<http://dx.doi.org/10.1080/10749357.2021.1943803>

<b>Title</b>	Postural control during quiet standing and voluntary stepping response tasks in individuals post-stroke : a case-control study
<b>Authors</b>	Moisan, G, Chayasit, P, Boonsinsukh, R, Nester, CJ and Hollands, K
<b>Publication title</b>	Topics in Stroke Rehabilitation
<b>Publisher</b>	Taylor & Francis
<b>Type</b>	Article
<b>USIR URL</b>	This version is available at: <a href="http://usir.salford.ac.uk/id/eprint/61685/">http://usir.salford.ac.uk/id/eprint/61685/</a>
<b>Published Date</b>	2021

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: [library-research@salford.ac.uk](mailto:library-research@salford.ac.uk).

1 **May 21st 2021**

2 **Postural control during quiet standing and voluntary stepping response tasks in individuals**  
3 **post-stroke: a case-control study**

4 Gabriel Moisan<sup>1-2</sup>, Pornprom Chayasit<sup>3</sup>, Rumpa Boonsinsukh<sup>3</sup>, Christopher James Nester<sup>2</sup>,  
5 Kristen Hollands<sup>2</sup>

6 Affiliations : <sup>1</sup>Department of Human Kinetics, Université du Québec à Trois-Rivières, Trois-  
7 Rivières, Canada, <sup>2</sup>School of Health Sciences, University of Salford, Salford, UK, <sup>3</sup>Faculty of  
8 Physical Therapy, Srinakharinwirot University, Nakhon Nayok, Thailand.

9 **Correspondence Address:**

10 Gabriel Moisan DPM PhD

11 3351 Boulevard Des Forges

12 Trois-Rivières, Québec, Canada, G9A 5H7

13 Email : [gabriel.moisan@uqtr.ca](mailto:gabriel.moisan@uqtr.ca)

14 Phone number: 1-819-376-5011 #3606

15

16

17

18

19

20

21 **Abstract**

22 **Background:** Postural control impairments following a stroke have impact on mobility, reduce  
23 independence and increase the risk of falls. Assessing these impairments during tasks  
24 representative of real-life situations, such as quiet standing (QS) and voluntary stepping response  
25 (VSR) will enhance our understanding of how the postural control system is impaired in individuals  
26 post-stroke (IPS). It will also inform the development of a more targeted and effective rehabilitation  
27 to prevent falls in IPS.

28 **Objectives:** Identify the postural control impairments encountered by IPS during QS and VSR.

29 **Methods:** Twenty IPS and sixteen healthy controls were recruited to perform QS and VSR tasks  
30 while ground reaction forces and whole-body motion were measured. Displacement and speed  
31 variation of the COM, centre of pressure (COP) displacement and spatiotemporal data were  
32 calculated and compared between groups.

33 **Results:** During QS, IPS exhibited greater maximal COP displacement in mediolateral direction,  
34 COM displacement in vertical direction and COM speed excursions compared to controls. During  
35 VSR, IPS exhibited smaller step length, braking force, posterior foot placement in relation to the  
36 pelvis and COM anteroposterior excursion compared to controls. IPS presented less static and  
37 dynamic postural stability compared to controls.

38 **Conclusions:** Greater postural sway during QS, smaller anteroposterior COM displacement before  
39 losing balance and altered voluntary recovering steps during VSR could place IPS at more risk of  
40 falling when they face a postural challenge in the community. These novel results will improve the  
41 current knowledge base and should be considered in IPS rehabilitation.

42 **Keywords:** biomechanics, balance control, stroke, postural control, quiet standing, voluntary

43 stepping response

44 **Word count:** 3016

45

## 46 **Introduction**

47           Stroke is a serious global health problem affecting 33 million in 2010<sup>1</sup> rising to 77 million  
48 by 2030.<sup>2</sup> It is the leading cause of disability in Europe and the United States of America.<sup>3, 4</sup> One  
49 of the numerous disabilities following a stroke is impairment in postural control, which has impact  
50 on mobility, reduce independence and increase the risk of falls.<sup>5</sup> All of these factors are known to  
51 affect self-efficacy, quality of life and the ability to maintain previous life roles.<sup>6</sup> Individuals post-  
52 stroke (IPS) exhibit asymmetrical load distribution and greater postural sway compared to healthy  
53 counterparts.<sup>7</sup> Also, during stance, their centre of pressure (COP) demonstrates larger and faster  
54 movements making them more at risk of falling.<sup>8</sup> Most previous studies focused on quiet standing  
55 (QS) tasks to assess the postural control impairments in IPS.<sup>8, 9</sup> However, other daily situations,  
56 known to be impaired in IPS, will challenge their postural control system such as bending the trunk  
57 to reach out for an object and taking a protective step following a perturbation.<sup>10-13</sup> During  
58 simulations of a slip or a trip, IPS exhibit many deficits known to increase their risk of falling,  
59 namely alterations in body movements and velocity and inefficient compensatory steps to avoid  
60 falling.<sup>12, 13</sup> Taking an efficient protective step after a perturbation is important, as using a multistep  
61 response is predictive of falls in daily life.<sup>13, 14</sup> Little attention has been paid to self-induced  
62 perturbation tasks that do not include a simulated trip or slip in previous studies. It is imperative to  
63 better understand the dynamic postural impairments during these challenging situations as they  
64 could also place IPS at risk of losing balance and fall. Furthermore, these tasks that do not include  
65 a simulated slip or trip could be more easily implemented in clinical contexts to evaluate postural  
66 control deficits or response to treatments in IPS, as they do not require specialised equipment.  
67 Ultimately, improving our knowledge about static and dynamic postural control impairments in

68 IPS could inform treatment targeted toward the specific deficits experienced by these individuals  
69 and could help increase their independence, participation in society and prevent falls.

70 Also, most previous studies analysed the COP trajectory to investigate postural control in  
71 IPS.<sup>8, 9, 15</sup> However, analysing the displacement of the centre of mass (COM) could also provide  
72 additional useful information about their postural control impairments<sup>16</sup> as it is the only variable  
73 that characterises body sway and have been identified as a strong predictor of risks of falls during  
74 highly challenging slip or trip-related perturbation tasks.<sup>12, 13</sup> However, its movement during less  
75 challenging postural control tasks, such as voluntary stepping response (VSR) and QS tasks,  
76 remains unclear for IPS. Thus, further studies are needed to determine how COM movements are  
77 altered in IPS during these tasks especially as it could be a good indicator of postural instability for  
78 these individuals.<sup>16</sup>

79 Thus, the main objective of this study was to identify the static and dynamic postural control  
80 impairments encountered by IPS using COP, COM and spatiotemporal outcomes. It was  
81 hypothesised that IPS will exhibit significant balance impairments known to increase the risk of  
82 falls, such as increased COP and COM displacement during static and dynamic postural control  
83 tasks and smaller recovering steps.

## 84 **Materials and methods**

### 85 Participants

86 Twenty IPS and 16 healthy age-matched adults (controls) were recruited to participate to  
87 this case-control study (Table 1) from July 2017 to December 2019. Sample size was determined  
88 based on the results of preliminary data. In this study, we sought to include IPS, at any time  
89 following a stroke, with at least the minimum level of mobility to safely take part to the protocol.

90 Thus, the results would be generalisable to the widest possible group of IPS with minimum mobility  
91 levels, meaning they might encounter some of the balance challenges the study posed in everyday  
92 life. For this reason, we included anyone who was able to stand independently and who had  
93 sufficient dynamic postural control to be able to safely take part to the study. Potential participants  
94 were excluded if they had cognitive deficits that affected the understanding of instructions or  
95 provision of informed consent (as indicated by Mini-Mental State Examination (MMSE) < 24),  
96 had visual problem that cannot be corrected with glasses, or had other neurological, cardiovascular,  
97 or musculoskeletal conditions (e.g. lower extremities amputation, injury, osteoporosis or etc.) that  
98 could have impeded their ability to perform any instrument testing without any braces or orthoses.  
99 IPS were included if they suffered of a stroke of any aetiology and controls were included if they  
100 had no self-reported health issues which may have affected their balance.

101 IPS were invited to take part through community stroke support and exercise groups in  
102 Greater Manchester, United Kingdom, whereas controls were recruited by poster advertisements  
103 and email invitations to University staff and students, participants of previous studies who have  
104 agreed to be contacted for further studies and carers/partners of IPS. The University of Salford,  
105 College of Health and Social Care Research Ethics Committee approved the study, and all  
106 participants provided written informed consent. The manuscript also conforms with the STROBE  
107 guidelines.

#### 108 Instrumentations

109 Kinematic data were collected with ten 100Hz cameras (Vicon, Oxford Metrics) with  
110 reflective markers attached on the participants' anterior superior iliac spines, posterior superior  
111 iliac spines, lateral thighs, lateral and medial femoral epicondyles, lateral legs, lateral malleolus,

112 2nd metatarsal heads, calcaneal tubercles, forehead, backhead, C7, T10, jugular notch, xiphoid  
113 process, right back, acromions, lateral humeral epicondyles, lateral arms, lateral forearms, left and  
114 right distal of radius, left and right ulnar head, and the 2nd metacarpal heads according to the Plug-  
115 in gait full body kinematic model. Kinetic data were collected at a sampling rate of 1 000 Hz with  
116 two force plates (AMTI, USA) embedded in the floor and synchronised with the kinematic data.

### 117 Experimental protocol

118 First, demographic data regarding age, sex, weight, height, number of falls in the last 12  
119 months and stroke duration (IPS only) were collected by self-report. Participants also filled the  
120 consent form and the Activities-specific Balance Confidence (ABC) scale.<sup>17</sup> Then, Mini-Mental  
121 Status Exam (MMSE),<sup>18</sup> Fugl-Meyer motor Assessment Lower extremity (FMA-LE) and sensation  
122 (FMA-S) scales (IPS only),<sup>19</sup> Dynamic Gait Index (DGI)<sup>20</sup> and Mini-Balance Evaluation-Systems  
123 test (mini-BESTest)<sup>21</sup> (only the anticipatory and reactive postural control sections) were  
124 administered to participants.

125 A calibration trial was recorded during relaxed standing in which participants placed their  
126 feet in a natural, self-selected posture, attempting equal weight bearing on both feet. A comfortable  
127 position was used as it allows for a more practical and realistic evaluation of postural control of  
128 IPS without changing postural stability compared to a standardised position.<sup>22</sup> Then, participants  
129 had to stand on a force plate for 30 seconds and try to move as little as possible with both feet  
130 always in contact with the force plate (QS task) until an audio cue appears as a signal to start leaning  
131 forward with whole body until they feel they are losing balance and take a step to prevent  
132 themselves from falling (VSR task). These two tasks were chosen as they can easily be  
133 implemented in clinical contexts with minimal equipment. The VSR task was previously used by



134 members of our research team to efficiently determine postural control differences between IPS  
135 and controls.<sup>23</sup> All participants were barefoot during all testing trials. No support of any kind was  
136 permitted during data collection. Three to five practice trials were allowed to promote familiarity  
137 with the test and ensure response stability. Then, ten trials were collected for each participant.  
138 Resting was permitted as needed to prevent fatigue. To assure participants' safety and prevent a  
139 fall during the tasks, a research assistant stood beside them to provide support as needed.

#### 140 Data processing

141 Biomechanical data were processed using Visual3D software (C-motion, Inc.,  
142 Germantown, MD, USA). Kinematic data were low-pass filtered using a 4<sup>th</sup> order Butterworth filter  
143 with a cut-off frequency of 10 Hz and individual segment coordinate systems were defined using  
144 anatomical markers. The variables measured during the QS task were the anterior/posterior (AP)  
145 and medial/lateral (ML) COP displacement and AP/ML/vertical maximal COM displacement and  
146 speed excursion (maximal subtracted by minimal speed). During the VSR task, the variables  
147 measured were step length and width (normalised to leg length), step duration, AP/ML/vertical  
148 COM excursion (between initial forward leaning and toe off), braking force (normalised to body  
149 mass), anterior foot placement in relation to pelvis (AFPP) and posterior foot placement in relation  
150 to pelvis (PFPP). Step length and width were respectively calculated using the stepping limb's heel  
151 and medial malleolus markers. A negative value for the step width refers to a step taken towards  
152 the other foot. A negative value for the COM excursion in ML and vertical directions respectively  
153 refers to the COM moving towards the stepping limb and the ground. The braking force was  
154 calculated by dividing the maximal AP force, measured using the force plate under the stepping  
155 foot, by the mass of the participant. AFPP and PFPP were respectively defined as the AP absolute  
156 distance between the COM and the heel marker of the stepping and support limbs.

157 Statistical analysis

158 IBM SPSS v.27.0.0.1 was used to compare the descriptive and biomechanical data of the  
159 IPS and the control groups using Mann Whitney tests for data that showed abnormal distribution  
160 and independent t tests for data that showed normal distribution according to Shapiro–Wilk’s test  
161 ( $p<0.05$ ). Cohen’s d effect sizes and 95% confidence intervals were calculated to compare the  
162 biomechanical data between the IPS and control groups. The level of statistical significance was  
163 set at  $p<0.05$  and  $d\geq 0.50$  for all analyses.

164 **Results**

165 There was no statistically significant difference in age, weight, height, MMSE score and  
166 falls in the last 12 months between the IPS and the control groups. Greater ABC, mini BESTest  
167 and DGI scores ( $p<.001$ ) were observed for the control compared to the IPS group (see Table 1).  
168 Individualised information is available in Supplementary material.

169 During the QS task, the IPS group exhibited greater COP displacement in ML direction  
170 (2.99 (CI=2.56-3.42) vs 2.30 cm (CI=1.73-2.64),  $d=0.92$ ,  $p=0.011$ ), COM displacement in vertical  
171 direction (0.59 (CI=0.47-0.71) vs 0.32 cm (CI=0.25-0.39),  $d=1.24$ ,  $p=0.001$ ) as well as COM speed  
172 excursion in AP (4.27 (CI=3.49-5.04) vs 3.03 cm/s (CI=2.53-3.52),  $d=0.90$ ,  $p=0.012$ ), ML (3.06  
173 (CI=2.49-3.63) vs 1.99 cm/s (CI=1.50-2.49),  $d=0.97$ ,  $p=0.007$ ) and vertical (2.35 (CI=1.51-3.18)  
174 vs 1.33 cm/s (CI=0.78-1.88),  $d=0.68$ ,  $p=0.030$ ) directions compared to the control group. No  
175 between-group differences were observed for COP displacement in AP direction and COM  
176 displacement in AP and ML directions. All between-group comparisons during the QS task are  
177 presented in the Table 2.

178            Only 19 of the 20 IPS completed the VSR task as one participant did not feel confident  
179 enough and was afraid of falling. The IPS group exhibited smaller step length (49.19 (CI=40.67-  
180 57.71) vs 89.84% (CI=56.31-123.37),  $d=-0.92$ ,  $p<0.001$ ), braking force (1.40 (CI=1.02-1.78) vs  
181 2.91 N/kg (CI=2.41-3.41),  $d=-1.81$ ,  $p<0.001$ ), PFPP (29.79 (CI=24.47-35.10) vs 43.18 cm  
182 (CI=38.84-47.52),  $d=-1.37$ ,  $p<0.001$ ) and COM excursion in AP direction (15.76 (CI=12.71-18.81)  
183 vs 20.49 cm (CI=17.90-23.07),  $d=-0.83$ ,  $p=0.020$ ) compared to the control group. No between-  
184 group differences were observed for step width and duration, COM excursion in ML and vertical  
185 directions and for AFPP. All between-group comparisons during the VSR task are presented in the  
186 Table 2.

## 187 **Discussion**

188            The main objective of this study was to identify the static and dynamic postural control  
189 impairments encountered by IPS using COP, COM and spatiotemporal outcomes. It was  
190 hypothesised that IPS would exhibit significant balance impairments known to increase the risk of  
191 falls, such as increased COP and COM displacements and speed excursions during static and  
192 dynamic postural control tasks and smaller recovering steps. Our results provide new insights about  
193 static and dynamic postural control deficits in IPS compared to healthy counterparts. Consistent  
194 with our hypotheses, greater COM vertical displacement was observed during QS, revealing more  
195 postural sway in IPS, known to increase the risk of falls.<sup>24</sup> It is the first study to measure COM  
196 movements to analyse postural control in IPS during QS and thus our results could not be directly  
197 compared with those of previous studies. However, Yu et al.<sup>16</sup> observed greater COM acceleration  
198 and scalar distance between COP and COM in IPS compared to controls during QS, suggesting  
199 impaired postural control. We also observed greater COM speed excursion in all planes during QS.  
200 Greater COM displacement is a strong predictor of falls in IPS during slip-related perturbation

201 tasks<sup>12, 13</sup> and the inability to control the dynamic COM state (velocity and position) is a causative  
202 factor of falls.<sup>13, 25</sup> The greater COM displacement and speed excursions observed in our study are  
203 novel and will add to the current body of knowledge by revealing that dynamic COM state is altered  
204 in IPS not only during highly challenging dynamic tasks, but also during easier static task. These  
205 results could perhaps be explained by the stroke-related impairments (e.g., proprioceptive  
206 impairments, muscle weakness, limb paralysis, post-stroke brain damage affective cognitive  
207 processing and sensorimotor integration) and thus greater postural sway results from these deficits.  
208 For example, greater postural sway, identified with COP measurements, was observed in IPS with  
209 impaired ankle proprioception.<sup>26</sup> Further studies are needed to better understand how the COM  
210 movements and speed excursions are altered in IPS during QS and to correlate these impairments  
211 with clinical tests.

212         IPS also exhibited greater COP displacement in ML direction during QS, which is  
213 consistent with a previous study that found greater ML RMS COP displacement for IPS compared  
214 to healthy counterparts.<sup>15</sup> ML stability seems to be mostly related to plantar cutaneous  
215 mechanoreceptors activity,<sup>27</sup> which gives information to the central nervous system about how the  
216 foot is positioned on the ground and how the body is leaning over the feet.<sup>28</sup> As many as 89% of  
217 IPS will exhibit somatosensory deficits,<sup>29</sup> known to be detrimental to postural control for this  
218 population.<sup>30</sup> Deficits in plantar cutaneous sensation may be one of the underlying factors  
219 explaining the greater ML COP displacement observed in this study and perhaps the high incidence  
220 of falls experienced by IPS. Surprisingly, no between-group statistically significant difference in  
221 COM ML displacement was found even though COM and COP movements are related during QS.  
222 However, the increased COM ML displacement in IPS compared to controls observed in our study  
223 almost reached statistical significance ( $p=0.072$ ) and the effect size was moderate ( $d=0.73$ ). With

224 a greater number of participants, this difference could perhaps reach statistical significance and  
225 thus change this result. Future studies should try to correlate the loss of plantar cutaneous sensation  
226 with postural control ability in IPS.

227         During the VSR task, IPS exhibited a smaller AP COM excursion compared to controls.  
228 This result suggests that IPS lose balance with less COM anterior displacement and could place  
229 them at greater risk of falling following a postural perturbation. No comparison could be made with  
230 previous studies, as it is the first to measure the COM movements during a forward leaning task.  
231 However, Portnoy et al.<sup>8</sup> observed a greater minimal anterior distance of the COP from the base of  
232 support and a smaller AP COP displacement during a functional reach task. These are consistent  
233 with our results and suggest that IPS lose balance with less anterior body displacement. The greater  
234 AP COM excursion found in our study could perhaps explain the shorter recovering step, the  
235 smaller PFPP and braking forces observed in IPS compared to controls. As COM travels less  
236 anteriorly, a shorter step is required to recover balance. As there was no difference in step duration,  
237 it can be hypothesised that the AP momentum of the stepping leg was decreased for IPS, explaining  
238 the smaller braking forces for these individuals. On the other hand, if this recovering step is too  
239 short, it may be inadequate to recover balance and force the IPS to use a multistep response to  
240 recover, which is known to predispose the individuals with poor postural control to fall.<sup>14</sup> Our  
241 results are consistent with those of Gray et al.,<sup>22</sup> which found that IPS took longer to initiate a  
242 shorter lateral voluntary step compared to controls. Preventing a fall often requires increasing the  
243 body's base of support with a quick and fast step to slow down the momentum of the body's  
244 COM.<sup>22</sup> The inability of IPS to react to external or self-induced perturbations with an efficient step  
245 place them at greater risk of falling. In light of our results, we suggest that clinicians treating IPS  
246 use QS and VSR tasks to quantify their postural control deficits.

247           The first limitation that should be considered for this study is that the data collection session  
248 took place in a highly controlled environment. The participants knew when and how they were  
249 going to lose balance and that the risks of falling were minimal during the VSR task. In real life  
250 situations, falls generally occurs during unexpected perturbations with little time to prepare or react  
251 <sup>31</sup>. The results of our study may not be representative of these situations. As IPS may experience  
252 fear of falling, they may have taken a conservative approach in taking the step during VSR. Future  
253 studies should investigate if similar postural control deficits are observed during unexpected  
254 postural perturbations. The second limitation of this study is that IPS may have experienced more  
255 fatigue than their healthy counterparts during the data collection as they present decreased physical  
256 fitness.<sup>32</sup> Physical exertion is known to increase postural sway in IPS.<sup>33</sup> To prevent IPS from  
257 experiencing fatigue, rest periods were given as needed to the participants. The third limitation is  
258 the difference in male/female ratio between the groups. Previous studies suggested slight  
259 differences in postural control between genders in other population.<sup>34, 35</sup> However, it remains  
260 unclear whether gender affects postural control of IPS during static and dynamic postural stability  
261 tasks.

## 262 **Conclusion**

263           IPS present less static and dynamic postural stability compared to healthy counterparts. The  
264 postural control impairments are observed when standing upright as well as when leaning forward  
265 and taking a step to recover and avoid falling. Greater postural sway during QS, highlighted by  
266 greater COM vertical displacement and COM speed excursions, as well as smaller AP maximal  
267 COM displacement before losing balance and altered recovering steps during VSR likely place IPS  
268 at more risk of falling when they face a postural challenge in the community. These results will  
269 inform more targeted and effective treatments to prevent falls in IPS.

270 **Declaration of interest statement**

271 The authors have no conflict of interest in this study.

272 **Funding**

273 This study was funded by the Canadian Institutes of Health Research (CIHR) and the Fonds  
274 de Recherche du Québec-Santé (FRQS) through scholarship received by the first author and  
275 Thailand Research Fund through the Royal Golden Jubilee PhD program (grant number  
276 PHD/0076/2558) received by the second author.

277 **References**

- 278 1. Feigin VL, Forouzanfar MH, Krishnamurthi R, et al. Global and regional burden of stroke  
279 during 1990-2010: findings from the Global Burden of Disease Study 2010. *Lancet*. Jan 18  
280 2014;383(9913):245-54.
- 281 2. Strong K, Mathers C, Bonita R. Preventing stroke: saving lives around the world. *The*  
282 *Lancet Neurology*. 2007/02/01/ 2007;6(2):182-187.
- 283 3. Benjamin EJ, Blaha MJ, Chiuve SE, et al. Correction to: Heart Disease and Stroke  
284 Statistics—2017 Update: A Report From the American Heart Association. *Circulation*. 201703  
285 2017;135(10):e646-e646.
- 286 4. Béjot Y, Bailly H, Durier J, Giroud M. Epidemiology of stroke in Europe and trends for  
287 the 21st century. *La Presse Médicale*. 201612 2016;45(12):e391-e398.
- 288 5. Geurts AC, de Haart M, van Nes IJ, Duysens J. A review of standing balance recovery  
289 from stroke. *Gait Posture*. Nov 2005;22(3):267-81.
- 290 6. Schmid AA, Van Puymbroeck M, Altenburger PA, et al. Balance and balance self-  
291 efficacy are associated with activity and participation after stroke: a cross-sectional study in  
292 people with chronic stroke. *Arch Phys Med Rehabil*. Jun 2012;93(6):1101-7.
- 293 7. Kamphuis JF, de Kam D, Geurts AC, Weerdesteyn V. Is weight-bearing asymmetry  
294 associated with postural instability after stroke? A systematic review. *Stroke Res Treat*.  
295 2013;2013:692137.
- 296 8. Portnoy S, Reif S, Mendelboim T, Rand D. Postural control of individuals with chronic  
297 stroke compared to healthy participants: Timed-Up-and-Go, Functional Reach Test and center of  
298 pressure movement. *Eur J Phys Rehabil Med*. Oct 2017;53(5):685-693.
- 299 9. Mehdizadeh H, Taghizadeh G, Ghomashchi H, et al. The effects of a short-term memory  
300 task on postural control of stroke patients. *Top Stroke Rehabil*. Oct 2015;22(5):335-41.
- 301 10. Chern JS, Lo CY, Wu CY, Chen CL, Yang S, Tang FT. Dynamic postural control during  
302 trunk bending and reaching in healthy adults and stroke patients. *Am J Phys Med Rehabil*. Mar  
303 2010;89(3):186-97.

- 304 11. Inness EL, Mansfield A, Lakhani B, Bayley M, McIlroy WE. Impaired reactive stepping  
305 among patients ready for discharge from inpatient stroke rehabilitation.(Research  
306 Report)(Report). *Phys Ther.* 2014;94(12):1755.
- 307 12. Patel PJ, Bhatt T. Fall risk during opposing stance perturbations among healthy adults and  
308 chronic stroke survivors. *Exp Brain Res.* Feb 2018;236(2):619-628.
- 309 13. Salot P, Patel P, Bhatt T. Reactive Balance in Individuals With Chronic Stroke:  
310 Biomechanical Factors Related to Perturbation-Induced Backward Falling. *Phys Ther.* Mar  
311 2016;96(3):338-47.
- 312 14. Hilliard MJ, Martinez KM, Janssen I, et al. Lateral balance factors predict future falls in  
313 community-living older adults. *Arch Phys Med Rehabil.* Sep 2008;89(9):1708-13.
- 314 15. Marigold DS, Eng JJ. The relationship of asymmetric weight-bearing with postural sway  
315 and visual reliance in stroke. *Gait Posture.* Feb 2006;23(2):249-55.
- 316 16. Yu E, Abe M, Masani K, et al. Evaluation of postural control in quiet standing using  
317 center of mass acceleration: comparison among the young, the elderly, and people with stroke.  
318 *Arch Phys Med Rehabil.* Jun 2008;89(6):1133-9.
- 319 17. Powell LE, Myers AM. The Activities-specific Balance Confidence (ABC) Scale. *J*  
320 *Gerontol A Biol Sci Med Sci.* Jan 1995;50a(1):M28-34. doi:10.1093/gerona/50a.1.m28
- 321 18. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for  
322 grading the cognitive state of patients for the clinician. *J Psychiatr Res.* Nov 1975;12(3):189-98.  
323 doi:10.1016/0022-3956(75)90026-6
- 324 19. Gladstone DJ, Danells CJ, Black SE. The fugl-meyer assessment of motor recovery after  
325 stroke: a critical review of its measurement properties. *Neurorehabil Neural Repair.* Sep  
326 2002;16(3):232-40. doi:10.1177/154596802401105171
- 327 20. Jonsdottir J, Cattaneo D. Reliability and validity of the dynamic gait index in persons with  
328 chronic stroke. *Arch Phys Med Rehabil.* Nov 2007;88(11):1410-5.  
329 doi:10.1016/j.apmr.2007.08.109
- 330 21. Tsang CS, Liao LR, Chung RC, Pang MY. Psychometric properties of the Mini-Balance  
331 Evaluation Systems Test (Mini-BESTest) in community-dwelling individuals with chronic  
332 stroke. *Phys Ther.* Aug 2013;93(8):1102-15. doi:10.2522/ptj.20120454
- 333 22. Gray VL, Yang C-I, Fujimoto M, McCombe Waller S, Rogers MW. Stepping  
334 characteristics during externally induced lateral reactive and voluntary steps in chronic stroke.  
335 *Gait Posture.* 2019/06/01/ 2019;71:198-204.
- 336 23. Chayasit P, Hollands K, Hollands M, Boonsinsukh R. Characteristics of Voluntary-  
337 induced Stepping Response in Persons with Stroke compared with those of healthy Young and  
338 Older Adults. *Gait Posture.* Oct 2020;82:75-82. doi:10.1016/j.gaitpost.2020.08.125
- 339 24. Sackley CM. Falls, sway, and symmetry of weight-bearing after stroke. *Int Disabil Stud.*  
340 Jan-Mar 1991;13(1):1-4.
- 341 25. Pavol MJ, Runtz EF, Pai Y-C. Young and Older Adults Exhibit Proactive and Reactive  
342 Adaptations to Repeated Slip Exposure. *J Gerontol A Biol Sci Med Sci.* 2004;59(5):M494-M502.
- 343 26. Niam S, Cheung W, Sullivan PE, Kent S, Gu X. Balance and physical impairments after  
344 stroke. *Arch Phys Med Rehabil.* Oct 1999;80(10):1227-33.
- 345 27. Okubo J, Watanabe I, Baron JB. Study on influences of the plantar mechanoreceptor on  
346 body sways. *Agressologie: revue internationale de physio-biologie et de pharmacologie*  
347 *appliquees aux effets de l'agression.* 1980;21(D):61.
- 348 28. Kavounoudias A, Roll R, Roll JP. The plantar sole is a 'dynamometric map' for human  
349 balance control. *Neuroreport.* Oct 5 1998;9(14):3247-52.



350 29. Connell L, Lincoln N, Radford K. Somatosensory impairment after stroke: frequency of  
351 different deficits and their recovery. *Clin Rehabil.* 2008;22(8):758-767.

352 30. Parsons SL, Mansfield A, Inness EL, Patterson KK. The relationship of plantar cutaneous  
353 sensation and standing balance post-stroke. *Top Stroke Rehabil.* Oct 2016;23(5):326-32.

354 31. Weerdesteyn V, de Niet M, van Duijnhoven HJ, Geurts AC. Falls in individuals with  
355 stroke. *J Rehabil Res Dev.* 2008;45(8):1195-213.

356 32. Persson CU, Hansson PO, Lappas G, Danielsson A. Physical Activity Levels and Their  
357 Associations With Postural Control in the First Year After Stroke. *Phys Ther.* Sep  
358 2016;96(9):1389-96.

359 33. Carver T, Nadeau S, Leroux A. Relation between physical exertion and postural stability  
360 in hemiparetic participants secondary to stroke. *Gait Posture.* Apr 2011;33(4):615-9.

361 34. Ozcan Kahraman B, Kahraman T, Kalemci O, Salik Sengul Y. Gender differences in  
362 postural control in people with nonspecific chronic low back pain. *Gait Posture.* Jul 2018;64:147-  
363 151.

364 35. Blaszczyk JW, Beck M, Sadowska D. Assessment of postural stability in young healthy  
365 subjects based on directional features of posturographic data: vision and gender effects. *Acta*  
366 *Neurobiol Exp (Wars).* 2014;74(4):433-42.

367

**Table 1 Demographic data**

	<b>Individuals post-stroke</b>	<b>Controls</b>
<b>Age (yrs)</b>	67.4 (11.5)	68.9 (8.1)
<b>Gender ratio (M/F)</b>	(16/4)	(7/9)
<b>Height (m)</b>	1.70 (0.09)	1.67 (0.09)
<b>Weight (kg)</b>	81.1 (11.6)	71.6 (15.9)
<b>MMSE (/30)</b>	28.6 (1.8)	29.3 (1.3)
<b>ABC (/100)</b>	63.0 (24.9)*	98.5 (2.2)*
<b>Mini BESTest (/12)</b>	5.2 (4.0)*	11.7 (0.8)*
<b>DGI (/24)</b>	15.4 (6.3)*	23.8 (0.6)*
<b>Falls last 12 months</b>	1.2 (3.2)	0.1 (0.3)
<b>Stroke duration (yrs)</b>	8.1 (9.3)	--
<b>FMA-LE (/34)</b>	22.4 (9.7)	--
<b>FMA-S (/12)</b>	11.2 (1.5)	--

368 Significant between-group differences ( $p < 0.001$ ) are identified with \*.

369 Results are expressed as mean (SD) except for gender ratio.

370 MMSE: Mini-Mental Status Exam, Mini BESTest: Mini-Balance Evaluation-Systems test, DGI:

371 Dynamic Gait Index, FMA-LE: Fugl-Meyer motor Assessment Lower extremity, FMA-S: Fugl-

372 Meyer motor Assessment sensation.

373

374  
375

**Table 2. Biomechanical parameters during QS and VSR tasks**

		Individuals post-stroke	Controls	Cohen's d	p value
<b>QS</b>	COP displacement AP (cm)	2.97 (2.41-3.52)	2.30 (1.82-2.77)	0.64	0.068
	COP displacement ML (cm)	2.99 (2.56-3.42)	2.18 (1.73-2.64)	0.92	0.011
	COM displacement AP (cm)	9.73 (5.25-14.20)	4.21 (2.12-6.30)	0.88	0.111
	COM displacement ML (cm)	2.24 (1.45-3.04)	1.37 (0.99-1.75)	0.73	0.072
	COM displacement vertical (cm)	0.59 (0.47-0.71)	0.32 (0.25-0.39)	1.24	0.001
	COM speed excursion AP (cm/s)	4.27 (3.49-5.04)	3.03 (2.53-3.52)	0.90	0.012
	COM speed excursion ML (cm/s)	3.06 (2.49-3.63)	1.99 (1.50-2.49)	0.97	0.007
	COM speed excursion vertical (cm/s)	2.35 (1.51-3.18)	1.33 (0.78-1.88)	0.68	0.030
<b>VSR</b>	Step length (%)	49.19 (40.67-57.71)	89.84 (56.31-123.37)	-0.92	<0.001
	Step width (%)	2.08 (0.44-3.71)	1.87 (-1.90-5.64)	0.04	0.220
	Step duration (ms)	281.14 (246.33-315.94)	297.16 (269.18-325.13)	-0.26	0.455
	COM excursion AP (cm)	15.76 (12.71-18.81)	20.49 (17.90-23.07)	-0.83	0.020
	COM excursion ML (cm)	-0.26 (-2.38-1.86)	-1.08 (-2.18-0.02)	0.23	0.500
	COM excursion vertical (cm)	-0.50 (-1.52-0.51)	-0.58 (-1.12--0.05)	0.05	0.888
	Braking force (N/kg)	1.40 (1.02-1.78)	2.91 (2.41-3.41)	-1.81	<0.001
	AFPP (cm)	14.54 (8.40-20.69)	17.45 (13.72-21.18)	-0.28	0.080
	PFPP (cm)	29.79 (24.47-35.10)	43.18 (38.84-47.52)	-1.37	<0.001

376 Results are displayed as means (95% confidence intervals). AFPP: Anterior foot placement in relation to pelvis, AP: Antero-posterior,  
377 COP: Centre of pressure, COM: Centre of mass, ML: Medio-lateral, PFPP: Posterior foot placement in relation to pelvis, QS: Quiet  
378 standing, VSR: Voluntary stepping response.

379  
380

Participant No.	Age (y)	Weight (kg)	Height (cm)	Sex (M/F)	Type of stroke	Stroke duration (y)	Hemiplegic side	Stepping foot	MMSE (/30)	ABC (/100)	FMA-LE (/34)	FMA-S (/12)	Mini BESTest (/12)	DGI (/24)
1	60	76.0	175.0	M	Hemorrhagic	6	R	R	30	56	7	12	1	N/A
2	71	82.6	170.0	M	Hemorrhagic	44	R	R	29	43	21	12	0	N/A
3	66	66.6	168.0	M	Hemorrhagic	8	R	R	30	81	26	12	5	N/A
4	62	88.9	170.0	M	Hemorrhagic	4	R	R	29	58	22	12	7	20
5	47	98.0	178.0	M	Hemorrhagic	7	L	L	30	79	20	10	6	12
6	45	95.0	173.0	M	Hemorrhagic	3	L	L	28	29	23	7	6	16
7	77	70.0	168.0	M	Ischemic	4	R	R	30	59	22	11	3	16
8	64	69.9	179.0	M	Hemorrhagic	3	L	L	29	89	28	10	8	19
9	84	77.1	179.0	M	Hemorrhagic	7	L	R	26	86	34	11	5	16
10	54	83.0	182.0	M	Ischemic	3	R	R	30	89	34	12	10	23
11	55	91.0	155.0	F	Hemorrhagic	9	R	L	28	55	4	12	1	9
12	80	77.0	162.5	F	Ischemic	3	R	L	27	77	30	10	6	15
13	80	65.0	156.5	M	Unknown	3	R	50%L, 50%R	29	100	32	12	11	24
14	75	102.0	183.0	M	Hemorrhagic	6	L	L	30	94	33	12	12	24
15	75	86.0	166.0	M	Ischemic	10	R	R	30	24	12	12	1	3
16	73	64.0	168.0	F	Hemorrhagic	9	L	L	30	18	20	12	2	10
17	81	91.0	157.0	F	Hemorrhagic	4	L	R	30	75	34	12	11	23
18	65	79.0	163.5	M	Ischemic	7	L	L	26	73	17	12	8	14
19	60	91.0	170.5	M	Hemorrhagic	3	L	N/A	27	36	4	8	0	6
20	75	68.0	174.5	M	Ischemic	10	R	L	24	38	24	12	1	12

**Supplementary material: Individualised information for individuals post-stroke**

**Abbreviations**

ABC: Activities-specific Balance Confidence scale

DGI: Dynamic Gait Index

F: Female

FMA-LE: Fugl-Meyer motor Assessment Lower extremity

FMA-S: Fugl-Meyer motor Assessment Sensation

L: Left

M: Male

Mini BESTest: Mini-Balance Evaluation Systems test

MMSE: Mini-Mental State Exam

N/A: Data not available

R: Right