



University of
Salford
MANCHESTER

Day surgery and general anaesthesia : what makes patients anxious?

Mitchell, MJ

Title	Day surgery and general anaesthesia : what makes patients anxious?
Authors	Mitchell, MJ
Publication title	Day Surgery Australia
Publisher	Australian Day Surgery Nurses Association
Type	Article
USIR URL	This version is available at: http://usir.salford.ac.uk/id/eprint/9735/
Published Date	2010

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: library-research@salford.ac.uk.

Day surgery and general anaesthesia: What makes patients anxious?

Abstract

For many patients the prospective of undergoing surgery and general anaesthesia is highly anxiety provoking. With the global rise in day surgery and limited nurse/ patient contact, anxiety has become a prominent issue. The aim of the study was to establish the degree of anxiety arising from elective day surgery and general anaesthesia and uncover specific anxiety provoking aspects. Day surgery patients (n=460) completed a questionnaire during recovery at home and return it by post. The majority of patients (85%) were anxious on the day of surgery and 50% desired a detailed level of information. Anaesthetic information provision, catastrophising and imminence of surgery were deemed to be reliable predictors of anxiety. The planned provision of anaesthetic information in advance of the day of surgery, emphasising 'controlled unconsciousness', provision of information to help limit catastrophising thoughts and assistance to reduce the impact of 'waiting' are recommended for the effective management of anxiety.

Keywords

Day surgery, general anaesthesia, anxiety, information provision and waiting.

Introduction

The purpose of the study was to investigate adult patient anxiety prior to elective, day surgery and general anaesthesia. This was considered necessary as little formal care for the anxious patient is currently provided in the elective day surgery setting ^[1, 2]. Indeed, the structured teaching of formal pre-operative anxiety care in nurse education in the United Kingdom is very limited ^[3, 4]. However, anxiety is a prominent issue prior to day surgery and general anaesthesia ^[5, 6], and with the predicted transfer of an increasing volume of elective surgery to day or 23 hour stay surgery in the coming years ^[7] the number experiencing pre-operative anxiety in day surgery units is likely to grow.

Literature Review

The volume and range of elective surgery now possible in day surgery facilities is growing and this rise is reflective of the global trend ^[8, 9]. Additionally, modern surgical techniques have made post-operative recovery quick with discharge home is achieved in a few short hours ^[10]. Moreover, such surgery is not minor 'lumps and bumps' but extensive surgery ^[11]. As the surgical assault on the body has diminished, the need for many once traditional physical aspects of surgical nursing care has faded ^[12]. Surgeons have fundamentally changed the way in which they practice and the once prominent features of post-operative nursing care are in rapid decline. "Surgery is now radically different from in 1948 or 1980, so it follows that surgery should not be provided in the same way as in 1948 or 1980." ^[13 p. 7]. The nursing profession must therefore carefully re-evaluate its place and role in this new era of surgical development.

Redundant tasks once central to post-operative surgical nursing have left a void, partially being filled by the adoption of quasi-medical tasks ^[14, 15]. However, in a review of the literature by Pearson et al. ^[16] it was concluded that no high quality evidence about the effectiveness of a nurse in the role of a surgical assistant existed. Indeed, the adoption of many medical tasks may have arisen in this manner, that is, as a result of political or financial pressures and not necessarily based upon nursing evidence ^[17, 18]. However, anxiety management in elective day surgery is a considerable nursing issue and much evidence is available to help inform practice ^[19-23].

Jakobsson et al. ^[24] suggested general anaesthesia to be the most prominent source of anxiety for patients prior to surgery. The use of 'needles' during induction ^[25] or a mask being placed over the face ^[26] lead to much anxiety. Gilmartin and Wright ^[1] interviewed 20 patients following day surgery and discovered that many had been anxious about waking up during their surgery or not waking afterwards. Hadjistavropoulos et al. ^[27] and Lee and Gin ^[28] recommend pre-operative education to ensure readiness for anaesthesia, educate about the method of induction of anaesthesia, reduce anxiety and explain post-operative care. Lack et al. ^[29] advocate general anaesthesia to be described not merely as being 'put to sleep' but as 'carefully controlled and supervised unconsciousness' allowing surgery to be performed painlessly. Van den Berg ^[30] interviewed 240 patients undergoing day surgery and statistically significantly more adults preferred inhalation induced anaesthesia to intravenous induction. Mitchell ^[22, 31] also established many patients undergoing regional and local anaesthesia were anxious with regard to being awake in theatre. A dental surgery study by Osborn & Sandler ^[32], demonstrated patients with raised anxiety required an increased amount of sedation to induce

and maintain a clinically acceptable level of sedation and were also more prone to intra-operative movements.

Patient anxiety prior to general anaesthesia is therefore of great concern to both patients and clinicians. As day and short stay surgery are now the main methods of elective surgical treatment throughout the World ^[9], it would be of great benefit when communicating with patients throughout the peri-operative period to have a clearer understanding of anxiety and knowledge of effective interventions.

A i m

To establish the degree of anxiety arising from elective day surgery and general anaesthesia and uncover specific anxiety provoking aspects.

M e t h o d

A questionnaire was constructed and used in a larger study examining the wider issues of anxiety associated with the clinical environment, hospital personnel and general and local/ regional anaesthesia (n=673) ^[31]. **JAN GA** However, this paper will only consider the questions relating to the anxiety provoking aspects of general anaesthesia associated with the clinical environment (Table 1). Anxiety prior to local/ regional anaesthesia has been reported elsewhere ^[22, 31]. The questionnaire was compiled using evidence gained from the literature and previously undertaken research in this field ^[4, 33, 34]. A pilot study was undertaken utilising the first 10% of respondents. This resulted in minor amendments prior to continuation of data collection. All items on the questionnaire were structured using a Likert Scale type format, for example, very anxious, a little anxious, made no difference, a little calm, very calm or never thought about it.

Data were collected from three public Day Surgery Units over a two-year period (2005 - 2007). All medical and nursing staff gave consent to the study prior to local Ethics Committee approval. A convenience sample of patients meeting the inclusion criteria (non life-threatening, intermediate surgery, no history of chronic physical or mental health, English speaking, 18 years upwards and not undergoing ophthalmic or dental surgery) were invited to participate. Ophthalmic and dental patients were excluded as they were deemed to experience additional anxieties resulting from the type of surgery, for example, loss of vision or possible alteration in physical appearance. Staff in each Day Surgery Unit invited patients on the day of surgery to take home the questionnaire. Questionnaires were completed at home 24 - 48 hours after surgery and return by mail in the pre-paid envelope provided.

R e s u l t s

Four hundred and sixty patients undergoing surgery and general anaesthesia took part. A variety of surgical procedures were undertaken with Gynaecological Surgery (25%), General Surgery (23%) and Orthopaedic Surgery being the most frequent. Participants' ages ranged from 18 years to 75 years with the average age being 46 years (271 females and 189 males). The number of patients experiencing anxiety on the day of surgery was 85% (Graph 1). The majority desired information about their planned surgery 1 - 4 weeks in advance (Graph 2) and 50% wished to receive a detailed level of information (Graph 3). Moreover, anxiety was largely generated by waiting on the day of surgery for the operation to commence and contemplating general anaesthesia (Graph 4).

A further examination of the data was undertaken using factor analysis. Exploratory factor analysis seeks to summarise all data uncovered (Table 1) and reduce findings into smaller

coherent groupings. Three groupings or factors were established from the identified questions (Table 1) and named 'Pre-operative Anaesthetic Information', 'Anaesthetic Catastrophising' and 'Imminence of Surgery'. Questions relating to anaesthetic information provision prior to surgery were given the title of Pre-operative Anaesthetic Information (Table 2) as during factor analysis all such aspects were positively related. Likewise, questions relating to potentially extreme events occurring whilst anaesthetised were given the overall title of Anaesthetic Catastrophising (Table 3). Finally, items relating to the impending start of surgery were given the title of Imminence of Surgery (Table 4).

These three new variables named Pre-operative Anaesthetic Information, Anaesthetic Catastrophising and Imminence of Surgery were entered into a multiple regression analysis. Multiple regression was employed to help determine if an overall level of anxiety can be predicted by the three new variables created during factor analysis (pre-operative anaesthetic information, anaesthetic catastrophising and imminence of surgery). This indeed was the case and Pre-operative Anaesthetic Information, Anaesthetic Catastrophising and Imminence of Surgery were deemed to be reliable predictors of increased anxiety on the day of surgery ^[23]. It can therefore be reliably stated that increased anxiety prior to day surgery and general anaesthesia can be predicted by Pre-operative Anaesthetic Information, Anaesthetic Catastrophising and Imminence of Surgery' (Tables 2, 3 & 4). Moreover, of the three, Imminence of Surgery evoked the most anxiety, that is, most apprehension arose from waiting for surgery to begin.

Discussion

First, a number of limitations may have influenced the results. Data collection was undertaken using a questionnaire that had not been previously employed. Therefore, questions could have been included which participants did not find relevant and other more relevant questions excluded. Participants may not therefore have been able to provide all the views they wished. However, the questionnaire was rigorously constructed from the available literature and previous research ^[35]. Second, only 41% of participants responded which could have limited the information obtained. Some patients may have wanted to contribute but during the first week may have been too poorly to complete the questionnaire. This is unfortunate but the reason why a larger sample was pursued.

The aim of the study was to establish the degree of anxiety arising from elective day surgery and general anaesthesia and uncover specific anxiety provoking aspects. Anxiety was expressed by 85% of patients (Graph 1). Such a high number alone may demonstrate the need for more effective, formal anxiety management prior to the day of surgery. Other researchers have also suggested anxiety to be a considerable issue in modern surgery ^[31, 36]. Anxiety arose from three distinct areas referred to here as Pre-operative Anaesthetic Information (Table 2), Anaesthetic Catastrophising (Table 3) and Imminence of Surgery (Table 4). All themes were derived from a combination of exploratory factor analysis and multiple regression. The results suggest anxiety about the gaining of sufficient anaesthetic information, erroneous beliefs concerning general anaesthesia and apprehension regarding the immediacy of surgery can reliably predict overall anxiety on the day of surgery.

Pre-operative Anaesthetic Information

Information provision has historically been viewed as fundamental to effective management of anxiety^[37]. In total, 87% patients desired to receive information between 1 - 4 weeks in advance of their surgery (Graph 2) and 50% required a detailed level of information (Graph 3). Anxiety may therefore have arisen from a real or perceived lack of information about surgery and/ or anaesthesia. Such findings appear to support previous evidence linking anxiety to a lack of sufficient information and a possible lack of a systematic approach to its delivery^[20, 38].

Much research has helped to extinguish the myth that patients are made more anxious if provided with additional information^[28, 39, 40]. Gilmartin^[41] interviewed 30 day surgery patients and concluded a large amount of patient information provision was required prior to surgery. In addition, a survey of 92 Swedish anaesthetists recommended mental preparation of patients and relatives (information, what to expect, emphasizing safety) should take place some days before surgery^[42]. The Royal College of Anaesthetists^[29] suggest when a patient agrees to surgery it is normally on the basis of information received from the surgeon about the intended surgical procedure. "Since the surgeon may not offer information about the anaesthetic technique, nor about the risks of anaesthesia, the patient generally agrees in principle to a procedure knowing very little about the anaesthetic that will be involved." (Lack et al. p. 15). Indeed, the patient may not even meet the anaesthetist until minutes before surgery when the conversation may focus (by necessity) on the biomedical issues with little psycho-social discussion^[43]. In such instances post-operative information regarding minimising nausea and vomiting (avoiding sitting up too quickly or drinking and eating immediately following surgery, taking slow deep breathes to reduce the sensation) may not be transferred to the patient^[44].

In a qualitative study^[45], it was further suggested information provision for home recovery was insufficient. Limited information once home was also highlighted in a review of the literature by Mitchell^[46]. Watt-Watson et al.^[47] reported 45% of patients did not receive sufficient information about their discharge medication. Following a survey of 116 day surgery patients, Bernier et al.^[48] provided an excellent pre-operative teaching guide which included 26 questions to be asked which covers most areas of patient concern. Finally, media-based interventions (video and written information) have proven effective in improving knowledge and reducing anxiety^[49].

Anaesthetic Catastrophising

Patients have been concerned about general anaesthesia for many decades^[50, 51] and with good reason. "Anaesthesia is an intrinsically hazardous undertaking with no inherent therapeutic benefit for the patient."^[52 p.1648] Kopp & Shafer^[53] state "During anesthesia the patient is controlled through interventions that reduce consciousness, eliminate language, alter memory, and limit autonomous actions." (p. 54). In the present study patients held many negative and unfounded beliefs about anaesthesia such as dying whilst anaesthetised, not being able to wake-up and waking during surgery. Anaesthetic catastrophising has been defined as a tendency to focus upon, and exaggerate, the negative aspects of anaesthesia, and the tendency to feel overwhelmed and unable to cope with or control the situation^[54].

Other researchers examining patient experience of anaesthesia have found similar results^[55]. Matthey et al^[56] revealed 47% of patients were very concerned with death, 45% with brain damage and 40% with intra-operative awareness. Fekrat et al^[57] surveyed 67 patients and 26

paired anaesthetists/ surgeons and concluded patient's major concerns were 'whether surgery would work', 'the body being damaged by surgery' and 'dying during anaesthesia'. Payne et al.^[58] conducted a study to determine patient preference for induction of anaesthesia, that is, induction via a mask or intravenously. Statistically significantly more patients preferred intravenous induction of anaesthesia as many did not like to have a mask over the face. Crockett et al.^[59] constructed a questionnaire to determine patient anxiety prior to anaesthesia. The questionnaire had 6 main themes - pre-occupation with morbid thoughts, outcome concerns, unconsciousness, loss of control, dependence on others and pain/ discomfort. Such catastrophising thoughts may therefore be quite common amongst patients awaiting general anaesthesia and again possibly reflect the limited level of knowledge.

Imminence of Surgery

Data from the present study indicated 59% of patients were anxious as a result of the wait prior to surgery (Graph 4). Such anxiety may have been exacerbated as 41% were also anxious about the anaesthetic and 31% by the 'thought of the unknown'. Indeed, this was evident in the 'imminence of surgery' grouping as this generated the most anxiety of the three themes. In support of such findings a review of the literature by Pearson et al.^[60] revealed waiting for surgery to commence also lead to greater anxiety. Freeman & Denham^[61] suggested waiting not only increased anxiety but reduced the quality of the day surgery patients' experience. In a qualitative study by Markovic et al.^[62] anxiety was again a common theme and associated with waiting, lack of a supportive relative, lack of information and having to walk to theatre. Carr et al.^[63] suggested the greatest level of anxiety occurred immediately prior to anaesthesia for 67% of in-patients. Khan & Bhutiani^[64] stated patients normally attend the hospital three times and waited an average of 41 - 53 weeks for day-case herniorrhaphy. A 'Walk-in-walk-out' (WIWO) clinic was therefore implemented which helped to limited anxiety, improve efficiency and reduce costs. "The patients in the WIWO clinic have their consultation and operation in one single visit and leave the hospital 2 - 3 hours after the operation."^[64 p. 2] To avoid waiting, patient inconvenience and increased anxiety a number of researchers have further recommended adoption of such 'one-stop' surgical practices^[65-68].

C o n c l u s i o n

Surgical practices have changed immeasurably throughout the World during the last two decades. The majority of elective surgery patients experience complex surgery and general anaesthesia on a day-case basis. As a result of such brief hospital stay the time available in which to develop the nurse/ patient relationship has become very restricted. However, an effective nurse/ patient relationship is vital in the management of patient anxiety, irrespective of how brief^[69]. Adequate information provision, reducing erroneous beliefs about anaesthesia and easing the time spent waiting are fundamental for effective anxiety management. Modern surgical nursing practices must strive to acknowledge such evidence by i) encouraging the wider teaching of such care in pre-registration programmes of study, ii) seeking to adopt such formal care in the clinical setting, and iii) by undertaking further studies to investigate this growing area of important patient care.

Implication for Practice

Information provided 1 - 4 weeks in advance of surgery for majority of patients.

Detailed level of information for majority of patients.

Both surgical and anaesthetic information provision.

Planned approach to information provision (formally documented).

Erroneous and unhelpful thoughts regarding anaesthesia explored in advance.

Emphasis upon "carefully controlled and supervised unconsciousness".

Emphasis upon close physical presence of the anaesthetist and nurse during surgery.

Close physical support, hand-holding and words of encouragement prior to induction

Simple methods of distraction, where appropriate

Attention to a 'patient friendly' ward and anaesthetic room environment.

Support from relative/ friend (where possible).

Increase in 'one-stop' practices.

TABLE 1
PRE-OPERATIVE APPREHENSION.

Pre-operative Apprehension
1) How would a nurse explaining your anaesthetic on the ward before going to theatre affect anxiety?
2) How would a visit from your anaesthetist before going to theatre affect your anxiety?
3) How would your anaesthetist explaining your anaesthetic before going to theatre affect anxiety?
4) How would your partner/ friend being with you in theatre before you are put to sleep affect your anxiety?
5) How would being told how long your anaesthetic will last affect your anxiety?
6) How would being told how soon you will be able to eat and drink again affect your anxiety?
7) How would always being told what was to happen next affect your anxiety?
8) How did waiting for your turn to go to theatre affect your anxiety?
9) How did having or the thought of having an injection affect your anxiety?
10) How did the thought of possibly needing a drip (intravenous infusion) affect your anxiety?
11) How did the thought of possibly removing any false item e.g. dentures or wig, affect your anxiety?
12) How did the thought of possibly needing an anaesthetic mask over your face affect your anxiety?
13) How did the thought of losing all control over your thoughts and movements affect your anxiety?
14) How did the thought of possibly waking up during your operation affect your anxiety?
15) How did the thought of having to trust strangers with your life affect your anxiety?
16) How did the thought of possibly dying while sleeping affect your anxiety?
17) How did the thought of possibly not waking up afterwards affect your anxiety?
18) How did having to walk to theatre affect your anxiety?
19) How did arriving at the theatre door affect your anxiety?
20) How did seeing medical equipment in the theatre room affect your anxiety?
21) How would the anaesthetist explaining things to you again in theatre before you are put to sleep affect your anxiety?
22) How would the nurse explaining things to you again in theatre affect your anxiety?
23) How would holding the hand of a nurse in theatre affect your anxiety?
24) How would your partner/ friend being with you in the recovery room very soon after waking up affect your anxiety?

TABLE 2
PRE-OPERATIVE ANAESTHETIC INFORMATION.

Pre-operative Anaesthetic Information
Nurse explaining the anaesthetic
Anaesthetist visiting
Anaesthetist explaining the anaesthetic
Informed how long anaesthetic will last
Informed where you can eat and drink
Informed what will happen next


TABLE 3
ANAESTHETIC CATASTROPHISING.

Anaesthetic Catastrophising
Thought of anaesthetic mask over the face
Thought of losing all control
Thought of waking up during surgery
Thought of having to trust strangers
Thought of dying while sleeping
Thought of not waking up afterwards


TABLE 4
IMMINENCE OF SURGERY.

Imminence of Surgery
Waiting your turn for surgery to commence
Removing false items
Arriving at the theatre door


GRAPH 1.
PATIENT RATINGS OF ANXIETY ON THE DAY OF SURGERY (n=460).




GRAPH 2.
PREFERRED TO RECEIVE INFORMATION (n=460).



GRAPH 3.
REQUIRED LEVEL OF PRE-OPERATIVE INFORMATION (n=460).



GRAPH 4.
MOST ANXIETY PROVOKING ASPECTS (n=460).



References

- 1) Gilmartin, J. and Wright, K. (2008) Day surgery: patients' felt abandoned during the preoperative wait. Journal of Clinical Nursing. 17 (18) 2418 - 2425.
- 2) Mitchell, M.J. (2007) Nursing research into modern day surgery: A literature review. Ambulatory Surgery. 13 (4) 1 - 29.
- 3) Fitzpatrick, E. and Hyde, A. (2006) Nurse-related factors in the delivery of preoperative patient education. Journal of Clinical Nursing. 15 (6) 671 - 677.
- 4) Mitchell, M.J. (2006) Nursing knowledge and the expansion of day surgery in the United Kingdom. Ambulatory Surgery. 12 (3) 131 - 138.
- 5) Rosén, S., Svensson, M. and Nilsson, U. (2008) Calm or not calm: The question of anxiety in the perianesthesia patient. Journal of PeriAnesthesia Nursing. 23 (4) 237 - 246.
- 6) Gardner, T.F., Nnadozie, M.U., Davis, B.A. and Kirk, S. (2005) Patient anxiety and patient satisfaction in hospital-based and freestanding ambulatory surgery centers. Journal of Nursing Care Quality. 20 (3) 238 - 243.
- 7) Baskerville, P. Surgical Education (*medical*) in Ambulatory Surgery. *7th International Congress on Ambulatory Surgery*. 2007. Amsterdam, Holland.
- 8) Gurusamy, K.S., Junnarkar, S., Farouk, M. and Davidson, B.R. (2008) Day-case versus overnight stay in laparoscopic cholecystectomy. Cochrane Database of Systematic Reviews. 1 (CD006798. DOI: 10.1002/14651858.CD006798.pub2.).
- 9) Toftgaard, C. (2009) Day surgery development. Ambulatory Surgery. 15 (1) 4 - 9.
- 10) Erkal, S. (2007) Patients' experiences at home after day case cystoscopy. Journal of Clinical Nursing. 16 (6) 1118 - 1124.
- 11) Smith, I. (2006) Dissecting the myths of day surgery: the anaesthetist's view. Journal of Perioperative Practice. 16 (5) 244 - 248.
- 12) Suhonen, R.A., Iivonen, M.K. and Välimäki, M.A. (2007) Day-case surgery patients' health-related quality of life. International Journal of Nursing Practice. 13 (2) 121 - 129.
- 13) Darzi, A. (2007) Saws and Scalpels to Lasers and Robots - Advances in Surgery. London: Department of Health.
- 14) Operating Theatre and Pre-operative Assessment Programme National Team. (2002) National good practice guidance on pre-operative assessment for day surgery. In: (N.H.S. Modernisation Agency Eds.). Step Guide to Improving Operating Theatre Performance. London: HMSO.
- 15) Kinley, H., Czoski-Murray, C., George, S., McCabe, C., Primrose, J., Reilly, C., Wood, R., Nicolson, P., Healy, C., Read, S., Norman, J., Janke, E., Alhameed, H., Fernandes, N. and Thomas, E. (2002) Effectiveness of appropriately trained nurses in preoperative assessment: Randomised controlled equivalence/ non-inferiority trial. British Medical Journal. 325 1323 - 1328.
- 16) Pearson, A., Richardson, M. and Cairns, M. (2004) Best practice in day surgery units: A review of the evidence. Ambulatory Surgery. 11 (1 - 2) 49 - 54.
- 17) Day, T. (2007) Respiratory assessment in the recovery unit: essential skills for the perioperative practitioner. Journal of Advanced Peri-operative Care. 3 (2) 41 - 49.
- 18) Woods, L.P. (1999) The contingent nature of advanced nursing practice. Journal of Advanced Nursing. 30 (1) 121 -128.
- 19) Prewett, A.T., Thakkar, H. and Found, P. (2008) An audit of patient satisfaction with anaesthetic services in a large day surgery unit. Journal of One-Day Surgery. 18 (1) 22 - 25.
- 20) Tse, K.-Y. and So, W.K.-W. (2008) Nurses' perceptions of preoperative teaching for ambulatory surgical patients. Journal of Advanced Nursing. 63 (6) 619 - 625.
- 21) Rhodes, L., Miles, G. and Pearson, A. (2006) Patient subjective experience and satisfaction during the peri-operative period in the day surgery setting: A systematic review. International Journal of Nursing Practice. 12 (4) 178 - 1192.
- 22) Mitchell, M.J. (2008) Conscious surgery: Influence of the environment on patient anxiety. Journal of Advanced Nursing. 64 (3) 261 - 271.

- 23) Mitchell, M.J. (2009) General anaesthesia and day case patient anxiety. Journal of Advanced Nursing. (In Press).
- 24) Jakobsson, J., Stomberg, M.W., Rawal, N., Brattwall, M. and Segerdahl, M. (2008) Day surgery for knee arthroscopy, open hernia repair and laparoscopic cholecystectomy anaesthesia routine and practice; The results from a Swedish nation-wide survey. Ambulatory Surgery. 14 (2) 1 - 23.
- 25) Ward, C., Varvinski, A. and Montgomery, J. (2007) Patients' choice of induction method: Do patients prefer being given a choice of their induction method? Journal of One-Day Surgery. 17 (2) 33 - 36.
- 26) van den Berg, A.A. (2003) Towards needleless induction of anaesthesia. Anaesthesia. 58 (8) 806 - 807.
- 27) Hadjistavropoulos, H.D., Dobson, J. and Boisvert, J.A. (2001) Information provision, patient involvement, and emotional support: prospective areas for improving anesthetic care. Canadian Journal of Anaesthesia. 48 (9) 864 - 870.
- 28) Lee, A. and Gin, T. (2005) Educating patients about anaesthesia: effect of various modes on patients' knowledge, anxiety and satisfaction. Current Opinion in Anaesthesiology. 18 (2) 205 - 208.
- 29) Lack, J.A., Rollin, A.-M., Thoms, G., White, L. and Williamson, C. (2003) Raising the Standard: Information for patients. London: RCoA and AAGBI.
- 30) van den Berg, A.A., Chitty, D.A., Jones, R.D., Sohel, M.S. and Shahan, A. (2005) Intravenous or inhaled induction of anesthesia in adults? An audit of preoperative patient preferences. Anesthesia and Analgesia. 100 (5) 1422 - 1424.
- 31) Mitchell, M.J. (2009) Patient anxiety and conscious surgery. Journal of Peri-operative Practice. 19 (6) 168 - 173.
- 32) Osborn, T.M. and Sandler, N.A. (2004) The effects of preoperative anxiety on intravenous sedation. Anesthesia Progress. 51 (2) 46 - 51.
- 33) Mitchell, M.J. (2000) Anxiety management: A distinct nursing role in day surgery. Ambulatory Surgery. 8 (3) 119 - 128.
- 34) Mitchell, M.J. (1997) Patients' perceptions of pre-operative preparation for day surgery. Journal of Advanced Nursing. 26 (2) 356 - 363.
- 35) Mitchell, M.J. (2005) Anxiety Management in Adult Day Surgery: A Nursing Perspective. London: Whurr.
- 36) Bauer, K.P., Dom, P.M., Ramirez, A.M. and O'Flaherty, J.E. (2003) Pre-operative intravenous midazolam: Benefits beyond anxiolysis. Journal of Clinical Anesthesia. 16 (3) 177 - 183.
- 37) Boore, J.R.P. (1978) Prescription for Recovery. London: Royal College of Nursing.
- 38) Lemos, P., Pinto, A., Morais, G., Pereira, J., Loureiro, R., Teixeira, S. and Nunes, C.S. (2009) Patient satisfaction following day surgery. Journal Of Clinical Anesthesia. 21 (3) 200 - 205.
- 39) Gillies, M.A.M. and Baldwin, F.J. (2001) Do patient information booklets increase peri-operative anxiety? European Journal of Anaesthesiology. 18 (9) 620 - 622.
- 40) Cowan, S.A., Engbaek, J., Kyst, A., Rosenstock, C.V., Jensby, K. and Jensen, L. (2000) Information, satisfaction and anxiety in connection with anaesthesia. A questionnaire study [Danish]. Ugeskrift for Laeger. 162 (9) 1231 - 1235.
- 41) Gilmartin, J. (2004) Day surgery: Patients' perceptions of a nurse-led pre-admission clinic. Journal of Clinical Nursing. 13 (2) 243 - 250.
- 42) Warrén Stomberg, M., Segerdahl, M., Rawal, N., Jakobsson, J. and Brattwall, M. (2008) Clinical practice and routines for day surgery in Sweden: Implications for improvement in nursing interventions. Journal of PeriAnesthesia Nursing. 23 (5) 311 - 320
- 43) Kindler, C.H., Szirt, L., Sommer, D., Hausler, R. and Langewitz, W. (2005) A quantitative analysis of anaesthetist-patient communication during the pre-operative visit. Anaesthesia. 60 (1) 53 - 59.
- 44) Royal College of Anaesthetists. (2006) Risk associated with your anaesthetic: Section 1 Feeling sick. London: RCOA.

- 45) Barthelsson, C., Lutzen, K., Anderberg, B. and Nordstrom, G. (2003) Patients' experiences of laparoscopic cholecystectomy in day surgery. Journal of Clinical Nursing. 12 (2) 253 - 259.
- 46) Mitchell, M.J. (2003) Impact of discharge from day surgery on patients and carers. British Journal of Nursing. 12 (7) 402 - 408.
- 47) Watt-Watson, J., Chung, F., Chan, V.W.S. and McGillion, M. (2004) Pain management following discharge after ambulatory same-day surgery. Journal of Nursing Management. 12 (3) 153 - 161.
- 48) Bernier, M.J., Sanares, D.C., Owen, S.V. and Newhouse, P.L. (2003) Pre-operative teaching received and valued in a day surgery setting. American Operating Room Nurses' Journal. 77 (3) 563 - 569.
- 49) Lee, A., Chui, P.T. and Gin, T. (2003) Educating patients about anesthesia: A systematic review of randomized controlled trials of media-based interventions. Anesthesia and Analgesia. 96 (5) 1424 - 1431.
- 50) Mackenzie, J.W. (1989) Daycase anaesthesia and anxiety: A study of anxiety profiles amongst patients attending a day bed unit. Anaesthesia. 44 (2) 437 - 440.
- 51) Gupta, A., Larsen, L., Sjoberg, I. and Lennmarken, C. (1994) Out-patients survey: A survey of anaesthesia care in a university hospital. Scandinavian Journal of Caring Sciences. 8 (2) 107 - 112.
- 52) Royston, D. and Cox, F. (2003) Anaesthesia: The patient's point of view. Lancet. 362 (9396) 1648 - 1658.
- 53) Kopp, V.J. and Shafer, A. (2000) Anesthesiologists and peri-operative communication. Anesthesiology. 93 (2) 548 - 555.
- 54) Faymonville, M.E., Mambourg, P.H., Joris, J., Vrijens, B., Fissette, J., Albert, A. and Lamy, M. (1997) Psychological approaches during conscious sedation. Hypnosis versus stress reducing strategies: A prospective randomized study. Pain. 73 (3) 361 - 367.
- 55) Spitellie, P.H., Holmes, M.A. and Dommino, K.B. (2002) Awareness during anesthesia. Anesthesiology Clinics of North America. 20 (3) 317 - 332.
- 56) Matthey, P., Finucane, B.T. and Finegan, B.A. (2001) The attitude of the general public towards preoperative assessment and risks associated with general anesthesia. Canadian Journal of Anaesthesia. 48 (4) 333 - 339.
- 57) Fekrat, F., Sahin, A., Yazici, K., M. and Aypar, U. (2006) Anaesthetists' and surgeons' estimation of preoperative anxiety by patients submitted for elective surgery in a university hospital. European Journal of Anaesthesiology. 23 (3) 227 - 233.
- 58) Payne, K., Elliot, R.A., Moore, J.K., Bennett, J. and Lawrence, G. (2004) Day-case anaesthesia: what would the patient prefer? Clinician in Management. 12 (3) 133 - 139.
- 59) Crockett, J.K., Gumley, A. and Longmate, A. (2007) The development and validation of the Pre-operative Intrusive Thoughts Inventory (PITI). Anaesthesia. 62 (7) 683 - 689.
- 60) Pearson, A., Richardson, M., Peels, S. and Cairns, M. (2004) The care of patients whilst in the day surgery unit: A systematic review. Health Care Reports. 2 (2) 22 - 54.
- 61) Freeman, K. and Denham, S.A. (2008) Improving patient satisfaction by addressing same day surgery wait times Journal of PeriAnesthesia Nursing 23 (6) 387 - 393.
- 62) Markovic, M., Bandyopadhyay, M., Manderson, L., Allotey, P., Murray, S. and Vu, T. (2004) Day surgery in Australia: Qualitative research report. Journal of Sociology. 40 (1) 74 - 78.
- 63) Carr, E., Brockbank, K., Allen, S. and Strike, P. (2006) Patterns and frequency of anxiety in women undergoing gynaecological surgery. Journal of Clinical Nursing. 15 (3) 341 - 352.
- 64) Khan, R.A. and Bhutiani, R.P. (2008) The walk in walk out hernia clinic. Ambulatory Surgery. 14 (2) 1 - 17.
- 65) Putnis, S., Merville-Tugg, R. and Atkinson, S. (2004) 'One-stop' inguinal hernia surgery - day-case referral, diagnosis and treatment. Annals of the Royal College of Surgeons of England. 86 (6) 425 - 427.
- 66) Shetty, A., Manimaran, N. and Reece-Smith, H. (2004) Direct access day-case hernia surgery: A logical option for reduction in waiting time. Ambulatory Surgery. 11 (1 - 2) 41 - 43.
- 67) Leff, D.R. and Bhutiani, R.P. (2006) Give the patients the choice - The 'walk in walk out' hernia clinic. Ambulatory Surgery. 12 (3) 125 - 129.

- 68) Sri-Ram, K., Irvine, T. and Ingham-Clark, C.L. (2006) A Direct Booking Hernia Service - A shorter wait and a satisfied patient. Ambulatory Surgery. 12 (3) 113 - 117.
- 69) Mottram, A. (2009) Therapeutic relationships in day surgery: a grounded theory study. Journal of Clinical Nursing. 18 (20) 2830 - 2837.